



FLY TIMES

ISSUE 61, October, 2018

Stephen D. Gaimari, editor
Plant Pest Diagnostics Branch
California Department of Food & Agriculture
3294 Meadowview Road
Sacramento, California 95832, USA
Tel: (916) 738-6671
FAX: (916) 262-1190
Email: stephen.gaimari@cdfa.ca.gov

Welcome to the latest issue of *Fly Times*! As usual, I thank everyone for sending in such interesting articles. I hope you all enjoy reading it as much as I enjoyed putting it together. Please let me encourage all of you to consider contributing articles that may be of interest to the Diptera community for the next issue, or for larger manuscripts, the Supplement series. *Fly Times* offers a great forum to report on your research activities and to make requests for taxa being studied, as well as to report interesting observations about flies, to discuss new and improved methods, to advertise opportunities for dipterists, to report on or announce meetings relevant to the community, etc., with all the associated digital images you wish to provide. This is also a great place to report on your interesting (and hopefully fruitful) collecting activities! Really anything fly-related is considered. And of course, thanks very much to Chris Borkent for again assembling the list of Diptera citations since the last *Fly Times*!

The electronic version of the *Fly Times* continues to be hosted on the North American Dipterists Society website at <http://www.nadsdiptera.org/News/FlyTimes/Flyhome.htm>. For this issue, I want to again thank all the contributors for sending me such great articles! Feel free to share your opinions or provide ideas on how to improve the newsletter. Also note, the [Directory of North American Dipterists](#) is constantly being updated. Please check your current entry and send all corrections (or new entries) to [Jim O'Hara](#) – see the form for this on the last page.

Issue No. 62 of the *Fly Times* will appear next April. Please send your contributions by email to the editor at stephen.gaimari@cdfa.ca.gov. All contributors for the next *Fly Times* should aim for 10 April 2019 (maybe then I'll get an issue out actually on time!) – but don't worry – I'll send a reminder. And articles after 10 April are OK too!

Issue 61, available online 14 November 2018

CONTENTS

NEWS

Thompson, F.C. — Final words on <i>Systema Dipteroorum</i>	1
Grimaldi, D.A. — <i>Drosophila bromeliae</i> Sturtevant, another tropical drosophilid in temperate North America	2–8
Hernández-Ortiz, V., Coates, R. — Diptera diversity: a survey of the tropical rainforest of Los Tuxtlas, México	9–13
Pollock, D.A. — Predation by Asilidae (Diptera) in eastern New Mexico and Texas: a preliminary analysis of 5000 prey records	14–20
Plakidas, J.D. — First record of the genus <i>Scythropochroa</i> Lengersdorf 1926 (Diptera: Sciaridae) from North America	21–22
Gray, R. — Pursuing mycetophilids in northern Nevada during 2018.....	23–26
Howson, E. — The Gnatwork	27

HISTORICAL DIPTEROLOGY

Evenhuis, N.L. — Revised dates of publication for Schiner’s “ <i>Fauna Austriaca. Die Fliegen (Diptera)</i> ”	28
Kotrba, M. — Gems from the H. Fischer collection: Diptera watercolors by E. H. Compton and a syntype of <i>Pegesimallus teratodes</i> (Hermann, 1906) (Asilidae).....	29–36
Evenhuis, N.L. — Reproduction of Loew’s 1859 Instructions for Collecting Diptera.....	37–45
Murphy, W.L. — Lloyd Vernon Knutson, 4 July 1934 – 10 January 2018	46–47
Gelhaus, J. — Bibliography of Papers on Crane Flies (Tipulidae s.l.) and other Diptera authored by George W. Byers	48–50

MEETING NEWS

Kirk-Spriggs, A. — The final countdown to ICD9!	51–52
Bertone, M. — Organized Meeting: North American Dipterists Society (NADS), 2018 Joint Annual Meeting of the Entomological Societies of America / Canada / British Columbia.....	53
Bertone, M. — North American Dipterists Society Code of Conduct	54
Courtney, G., Bowles, D. — 2019 Field Meeting of North American Dipterists Society, 3-7 June 2019, Bull Shoals Field Station, Missouri.....	55–57
Namayandeh, A. — Eagle Hill Institute in Maine, USA will be holding a Workshop on Chironomidae Taxonomy, June 9-15, 2019	58–59
Brown, B.V. — Fly School, 23 June–6 July 2019.....	60

OPPORTUNITIES

Dikow, T. — S.W. Williston Diptera Research Fund	61
--	----

DIPTERA ARE AMAZING!	62
----------------------------	----

BOOKS AND PUBLICATIONS	63–75
------------------------------	-------

SUBMISSION FORM, DIRECTORY OF NORTH AMERICAN DIPTERISTS	76
---	----

NEWS

Final words on *Systema Dipterorum*

F. Chris Thompson

Ponte Vedra, Florida, USA; xelaalex@cox.net

After too many years and over a quarter of million names (259,940 as today), I am retiring from the name business. Yes, I will continue to track ant fly and flower fly names (Microdontidae and Syrphidae), but I will leave the rest of *Systema Dipterorum* to Neal Evenhuis and Tom Pape.

***Drosophila bromeliae* Sturtevant, another tropical drosophilid in temperate North America**

David A. Grimaldi

Division of Invertebrate Zoology, American Museum of Natural History,
Central Park West at 79th Street New York, New York 10024-5192, USA; grimaldi@amnh.org

Summary

Drosophila bromeliae Sturtevant is reported from northern New Jersey, where it was found breeding in pumpkin flowers (*Cucurbita pepo* L.), and in central Massachusetts, both in 2018. The prior northernmost records of the species had been Cuba in 1915 and Valusia County, Florida in 1997, an extension of 13° (~1800 km) north. All species of the *D. bromeliae* group are Neotropical and, where known, breed in flowers. Observations are provided on *D. bromeliae* from New Jersey, as are the main identifying characteristics of the species. This is one of six species of Drosophilidae that have invaded the temperate Nearctic in the past decade, which are briefly reviewed. A request is made for any additional records and samples from North America.

Background

The *Drosophila bromeliae* group consists of 13 named species, the overall distribution being entirely Neotropical, from Cuba and throughout the Caribbean; Mexico (Nayarit) south through Central America and into South America to northern Bolivia (Grimaldi, 2016). Based on intensive studies by Schmitz (2010), species of the group breed in various flowers, some being polyphagous and others restricted to a particular genus of plants. The nominal species of the group, *D. bromeliae* Sturtevant, has the widest distribution, from Cuba and the Bahamas south throughout apparently the entire Caribbean, throughout Central America south of the Yucatan Peninsula, and into South America to northern Brazil (Grimaldi, 2016, Schmitz, 2010). Morphologically, species in the group can be reliably distinguished only on the basis of male genitalia, sometimes by female genitalia depending on the species (Grimaldi, 2016). Since the taxonomic revision of this group (Grimaldi, 2016), a new U.S. record of the species has been found (besides the one reported here from New Jersey), based on overlooked specimens in the AMNH collections. These are a series of 10 males and 12 females collected in Valdez, Volusia Co., Florida, V/16/1997, by Mark and Maria Minno, in flowers of Okeechobee gourd (*Cucurbita okeechobeensis*). The Florida population of this plant is a rare subspecies endemic to several areas in southern Florida; it is listed as federally endangered. It is unknown if southern Florida is part of the natural distribution of *D. bromeliae*, which would not be surprising given occurrence of the fly in Cuba and the Bahamas.

Observations

For the past 32 years the author has monitored Drosophilidae and some other flies that occur on his property in Pompton Plains, Morris County, New Jersey, and vicinity. The town is at the eastern base of the Watchung escarpment, in the northern end of the Atlantic Coastal Plain, a natural corridor extending to Florida. From early July to late August, 2018, abundant *Drosophila bromeliae* were found on blossoms of large pumpkin vines, which were growing out of piles of composted horse manure with pine shavings. Though there were only four pumpkin vines, their cumulative length was approximately 40 meters, usually with dozens of large blossoms at any one time. Other visitors to the blossoms were striped cucumber beetles, *Acalymma vittatum* (Chrysomelidae) and some unidentified bees. The bees, beetles, and flies actively consumed the abundant pollen. There was usually a small pile of pollen at the bottom of the corolla dislodged by the bees and beetles, on which flies fed. There did not appear to be any negative association between flies and beetles. Other than feeding, other fly

activities included resting and courting on the inner and upper surfaces of the petals. Their movements were slow, probably due at least in part to the fine long hairs on the lower portions of the petals. Every opened flower (Fig. 1A) harbored flies, which varied in number from 5 to a maximum of approximately 80 (Fig. 1B) (mean of 20 flies per blossom, $N = 13$ blossoms, counted 25 August, 2018). Some flies congregated on the outside surfaces of petals (Fig. 1C). Flowers that were exposed to sun had many fewer flies; most flowers were in the shade of trees. The blossoms closed in late evening through the night, and some flies (perhaps not all) cloistered in the closed flowers (they closed earlier in the day at hot temperatures); flies were most abundant in the mornings on the open flowers. On occasional cool mornings in August (ca. 16° C), opened, fresh blossoms contained no flies (the flies may have settled on the warm compost). Senescent flowers, closed and shriveled, usually had no flies, but occasionally a few and these always inside the flower.

Based on the numbers of flies in each fresh blossom, and their increase in numbers from early July to late August, the flies must have been breeding. Pumpkin yield was low (only five fruits from 40 m of vine), possibly the cause of premature flower senescence. To determine if the flies were breeding in the pumpkin blossoms, 5 senescent and 5 fresh blossoms were carefully pulled apart, especially the reproductive structures at the base. Senescent moist petals commonly were riddled with fine holes, presumably from feeding beetles. Bases of petals were thicker, with spongy tissue, and the anthers of male flowers formed a fleshy cone with a central chamber that was almost entirely closed -- ideal substrates for *Drosophila* larvae. However, no eggs or larvae were seen. To further determine if flies were breeding in the flowers, the dissected flowers and 10 additional senescent flowers were spread out in a plastic tray on top of moist, white sheets of paper toweling, under a plastic cover with a central funnel (BioQuip©). The paper toweling was used, rather than sand or vermiculite, to better visualize any puparia that would eventually develop.

From the 20 flowers placed in a container well over 100 puparia and adult flies emerged. The flowers had deliquesced, and the first puparia appeared on the paper toweling 5 days after containment; clumps of puparia appeared on corners a day later. Most adults eclosed 9 days after containment, approximately 3 days after pupariation at ambient temperature. Thus, eggs were entirely unnoticed when the dissected flowers were first inspected, perhaps having been embedded within flower tissue by the ovipositing females.

Eight female flies collected on 25 August were dissected, to check their reproductive status. The colored ovaries can be seen through the translucent pleural membrane, ovaries being light pink and in the males testes are reddish. The crop was also visible, in this case bright yellow from the ingested pumpkin pollen (only a few of the 40 or so flies preserved had not fed). Of the eight females dissected, in six the ovaries were small and not fully developed; in the other two specimens the ovaries were larger, with about 8 ovarioles, but none were gravid, i.e., with a mature egg in the lateral or common oviducts. It is possible that this was a recently eclosed cohort, not yet reproductively matured.

By 20 September, 2018 the squash plants were dying back, and among the 10 flowers left there was a very noticeable decline in the number of flies on them (an average of 15 per flower, ranging from 9 to 24).

Several individuals of *D. bromeliae* were observed in pumpkin flowers in Northampton, Massachusetts on September 15, 2018 by Isabel Novak. Voucher specimens from the NJ and MA localities are in the AMNH collection.



Figure 1. Pumpkin flower hosts of *Drosophila bromeliae* in Morris County, New Jersey (August, 2018).
A. Plant. B. Flower with about 80 flies. C. Side of same flower as in B.

Identification

Drosophila bromeliae is dark yellowish to light dusky brown, a rather typical species of *Drosophila* with clear, unmarked wing and bright red eyes. All of the New Jersey flies are distinctly darker than any of the flies from Florida or the neotropics (it's unknown if this is genetically based or due to the cooler temperatures during larval and pupal development). *Drosophila bromeliae* is easily distinguished from North American species of drosophilids by a combination of the flat facial carina (typical of *Drosophila*); well-developed oviscapt; abdominal tergites 1-5 each with a thin, diffuse, dark transverse band interrupted in the middle; the red testes and pink ovaries (seen through the pleural membrane when live or freshly preserved); and especially by the pair of prescutellar setae (enlarged pair of acrostichals) (Figs. 2A-D). The only other North American species of *Drosophila* that have prescutellar setae are the following: species in the *virilis* group (easily separated by the completely dark abdomen); *D. sigmoides* (which has a sinuous posterior crossvein and clouded wing); and some species in the *repleta* group (which have a spotted body).

Definitive separation of *D. bromeliae* from other members of the *bromeliae* species group requires male genitalia (Figs. 2E-F), especially the aedeagus, to lesser extent the surstylus and hypandrium (Grimaldi, 2016). The aedeagus of *D. bromeliae* in lateral view is arched, long and rather slender, the tip lacking subapical spines and ovoid to slightly drop-shaped in full ventral view (Fig. 2F). Eggs are also distinctive, with two very short anterior filaments (Grimaldi, 2016: fig. 6A).

Significance

It is too early to determine if *Drosophila bromeliae* has established perennial breeding populations in northeastern U.S. The summer of 2018 was particularly hot and wet, ideal for seasonal colonization of a tropical species. It also needs to be examined if the species is injurious, to either cucurbits or other fruits and vegetables. Schmitz (2010) reared the species in Brazil from flowers of morning glories (*Ipomoea*: Convolvulaceae), *Hibiscus* (Malvaceae), *Distictela* (Bignoniaceae), and *Thunbergia* (Acanthaceae) (this last genus native to Africa and Asia). I have also found *D. bromeliae* in *Ipomoea* flowers in the Dominican Republic (Grimaldi, 2016). These records and the ones involving cucurbits in the U. S. indicate that the species is quite polyphagous. It can even be maintained in the lab on standard banana-cornmeal medium and Carolina *Drosophila* Instant medium®, which is very unusual for obligate flower-breeding species in Drosophilidae, and indicating that it is quite adaptable.

Other Invasive Drosophilidae

Drosophila bromeliae joins a growing list of drosophilid species that have dispersed or were introduced to North America in the past decade or so, several of them having become very successful (Miller et al., 2017; Werner and Jaenike, 2017).

The most notorious invasive is *Drosophila (Sophophora) suzukii* (Matsumura), a native of Japan and a major pest of berries and soft fruits. It was first detected in North America from specimens captured in California in 2008 (Hauser et al., 2009; Hauser, 2011), and has since spread throughout the continent up to approximately 47°N latitude in eastern North America (Miller et al., 2017; Werner and Jaenike, 2017), as well as Europe and Asia. It is now common even in mature broadleaf forests in northeastern North America and northern Michigan (D. Grimaldi, Thomas Werner, pers. observ.), breeding in various cultivated and native fruits (Lee et al., 2015).

Like *D. suzukii*, *Zaprionus indianus* Gupta also breeds in various fresh fruits, but also decaying ones as well. It is not nearly as injurious to fruit crops as is *D. suzukii*, though there is concern that it may be out-competing native *Drosophila* in the neotropics. The natural distribution of *Z. indianus* is




Figure 2. Diagnostic features for identifying *Drosophila bromeliae* (see labels). A. Whole fly (male). B. Head, frontal. C. Dorsoapical portion of thorax. D. Abdomen, dorsal. E. Epandrium with surstyli. F. Aedeagus.

central Africa and southern Asia. It was first detected in the New World in Brazil in 1999, and in the USA in Florida in 2003 (Van der Linde et al., 2006). Since then it has spread throughout the New World tropics (D. Grimaldi, pers. observ.) and occasionally is found in New York and even southern Canada (Miller et al., 2017).

Drosophila cardini is widespread in the neotropics from Bolivia and southern Brazil to Mexico and throughout the Caribbean, including southern Florida. There are two records of it in northeastern North America: from near Guelph, Ontario, Canada and northern New Jersey, USA, collected in 2005 (Miller et al., 2017) (the New Jersey record is also from the author's backyard, collected on stinkhorn fungi). This species has a particularly broad niche, breeding in decaying fruits, vegetables, and fungi. It does not seem to be established in the northern states, although it probably is well established in Florida and U. S. Gulf states.

Two U.S. colonists that are flower-breeding like *Drosophila bromeliae* are *Drosophila gentica* Wheeler and Takada, and *Drosophila (Sophophora) flavohirta* Malloch, both established in California (Grimaldi et al., 2016). *Drosophila gentica* belongs to the Neotropical *flavopilosa* group, all of which breed in flowers, and in the case of *D. gentica* primarily on the ornamental flowering shrub *Cestrum* (Solanaceae) in both its native Central America as well as in California. It is unlikely to become an agricultural pest. *Drosophila flavohirta* is an anomalous member of the *melanogaster* species group, a native of Australia, and has also been introduced to South Africa and Madagascar. It breeds in flowers of Eucalyptus trees, which are planted throughout California. It can affect honey yield.

The most unusual introduction to the U. S. is *Phortica variegata* (Fallén), originally Palearctic in distribution. Males of *Phortica*, and in the closely related genus *Amiota*, have the peculiar habit of swarming around a person's head, like blackflies, so they are easily collected and surveyed. They appear to be attracted to eyes and lacrimal secretions, and in Europe and Asia *Phortica* species transmit thelazid nematodes to dogs, humans, and other mammals (infected cattle, horses and other ungulates have not been reported) (reviewed in Werner and Jaenike, 2017). Based on the few breeding records for *Amiota* and *Phortica*, *Phortica variegata* probably breeds in injured and decaying trees. It was first discovered in Orange County, New York in 2014 and Monroe Co., New York in 2015 (Miller et al., 2017; Werner and Jaenike, 2017), and it appears to be spreading, even abundant and nuisance in eastern Massachusetts.

Why the Spread?

There are several lines of explanation for the dramatic spread of these drosophilids in the past decade or so. One is that there has been increased commercial trade of fruits and horticultural plants, or relaxations in import regulations and surveillance, although I have no data for this. It's unlikely that this entirely explains these invasives, since they have such disparate life histories, breeding in fresh fruits, decaying fruits, fresh flowers, and trees. Another explanation may be climate change, especially since *Drosophila bromeliae* and *D. cardini* were found in New Jersey during particularly hot, wet summers, and the tropical species *Zaprionus indianus* is becoming more common there.

Request

The author would be very appreciative for obtaining information on presence/absence of *Drosophila bromeliae* in other areas of North America, particularly in the southeastern U.S., where the species is suspected to be established. Squash, pumpkin, and other cucurbit flowers may be especially sought by the flies, but any large flowers are likely to be attractive, including ornamentals (e.g., *Hibiscus*, *Datura*, etc.). Also, any samples of *Phortica* (or their *Amiota* relatives) would be appreciated (see Werner and Jaenike, 2017 and Miller et al., 2017 for external features to ID the genus). Genitalia would need to be checked here to verify the species identification. *Phortica* and *Amiota* are easily collected in hardwood forests.

References

- Grimaldi, D., P.S. Ginsberg, L. Thayer, S. McEvey, M. Hauser, M. Turelli, and B. Brown. 2015. Strange little flies in the big city: exotic flower-breeding Drosophilidae (Diptera) in urban Los Angeles. *PLoS ONE* 10(4): e0122575. doi: 10.1371/journal.pone.012257
- Grimaldi, D.A. 2016. Revision of the *Drosophila bromeliae* Species Group (Diptera: Drosophilidae): Central American, Caribbean, and Andean species. *American Museum Novitates* 3859: 1-55.
- Hauser, M. (2011). A historic account of the invasion of *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) in the continental United States, with remarks on their identification. *Pest management science*, 67(11), 1352-1357.
- Hauser, M., S. Gaimari, and M. Damus. 2009. *Drosophila suzukii* new to North America. *Fly Times* 43: 12-15.
- Lee, J. C., A. J. Dreves, A. M. Cave, S. Kawai, R. Isaacs, J. C. Miller, S. Van Timmeren, and D. J. Bruck. 2015. Infestation of wild and ornamental noncrop fruits by *Drosophila suzukii* (Diptera: Drosophilidae). *Annals of the Entomological Society of America* 108 (2): 117-129.
- Miller, M. E., S. A. Marshall, and D. A. Grimaldi. 2016. A review of the species of *Drosophila* (Diptera: Drosophilidae) and genera of Drosophilidae of northeastern North America. *Canadian Journal of Arthropod Identification* (on line): doi:10.3752/cjai.2017.31
- Schmitz, H.J. 2010. *Genética, Ecologia e Evolução de drosofilídeos (Insecta, Diptera) associados a flores*. Ph.D. Dissertation, Universidade Federal do Rio Grande do Sul. Porto Alegre, Brazil.
- Van der Linde, K., G.J. Steck, K. Hibbard, J.S. Birdsley, L. M. Alonso & D. Houle, D. 2006. First records of *Zaprionus indianus* (Diptera: Drosophilidae), a pest species on commercial fruits from Panama and the United States of America. *Florida Entomologist* 89(3), 402-404.
- Werner, T. and J. Jaenike. 2017. *Drosophilids of the Midwest and Northeast*. University of Rochester Libraries: www.rochester.edu/newscenter/fruit-flies-field-guide-jaenike-272372/

Diptera diversity: a survey of the tropical rainforest of Los Tuxtlas, México

Vicente Hernández-Ortiz¹, Rosamond Coates²,
Martha Madora-Astudillo², and José F. Dzul-Cauich¹

¹Instituto de Ecología A.C., Red de Interacciones Multitróficas,
Veracruz, México; vicente.hernandez@inecol.mx.

²Universidad Nacional Autónoma de México,
Estación de Biología Tropical Los Tuxtlas, Veracruz, México.

Introduction

The biodiversity of tropical rainforests has been threatened by an intensive loss of habitat and changes in land use policies (Gutiérrez Granados *et al.* 2018). So, the rapid disruption of tropical forests probably jeopardizes global biodiversity more than any other contemporary phenomenon. Deforestation of tropical forests has been continuously increasing in recent decades, so protected areas are a natural refuge for biodiversity. According to Laurance *et al.* (2012), tropical protected areas are often intimately linked ecologically to their surrounding habitats, and their analysis reveals great variation in reserve 'health', about half of all reserves have been effective, but the rest are experiencing an erosion of biodiversity that is often alarmingly widespread taxonomically and functionally.

Most of the insect groups that live in tropical environments of the Americas are unknown. In the last 50 years, studies of Diptera of the Neotropical region have been marginal, although in this region high biodiversity occurs. Moreover, there is a crisis in the description of new taxa, mainly due to lack of taxonomists in Latin America (Brown 2005).

In the Network of Multitrophic Interactions (INECOL, Mexico) we conducted a long-term project called "Biodiversity and Systematics of the Diptera of Mexico" (BIODIMEX), whose main goal is the knowledge of the diversity of the Mexican Diptera, either in natural and anthropic environments. We emphasize our investigation in the families of Acalypttratae, and other taxa with phytophagous habits, including taxonomic studies, and aspects of their ecology and natural history. These studies are being conducted primarily in the cloud forest and rainforest communities. By example, the study of the Diptera diversity of the canopy and the understory of coffee agroecosystems in Veracruz (Hernández-Ortiz & Dzul-Cauich 2008).

The Biosphere Reserve of Los Tuxtlas (BRLT) is located at the central region of the state of Veracruz, being the northernmost tropical rainforest in the Americas. Moreover, that area also belongs to the "Mexican Transition Zone", where flora and fauna of Nearctic and Neotropical origin converge, therefore, its biological diversity requires to be studied.

For instance, some relictual species as *Rhagoletis turpiniae* (Tephritidae) occurs in this Reserve, whose closest relatives are found in North America (Hernández-Ortiz 1993). Currently we continue discovery new species in the BRLT, as *Beebeomyia tuxtlaensis* (Richardiidae), recently described in association with aroid inflorescences (Hernández-Ortiz & Aguirre 2015), and near as 15 other new species of Tephritoidea, which are being described (Hernández-Ortiz *et al.* unpublished data).

As part of this long-term project, we are currently developing a subproject on diversity and seasonal dynamics of the Dipteran families living in the understory of a preserved area in the region of Los Tuxtlas.

The study area

In Mexico, there are few places where there are still preserved rainforest areas. The Biosphere Reserve of Los Tuxtlas (BRLT), is located at the easternmost edge of the volcanic belt that crosses Mexico, presenting a complex series of uprisings running in NW-SE direction (Dirzo & García 1992). This massif arises in the coastal plain of the Gulf of Mexico and is relatively isolated from other mountain systems (Fig. 1).

The reserve consists of three core areas, which are the Volcano San Martín Tuxtla, Volcano Santa Martha, and Volcano San Martín Pajapan (29,720 hectares), and the rest (125,402 hectares) make up the buffer zone. Within the reserve, the Tropical Biology Station Los Tuxtlas-UNAM (LTBS) contribute with 640 hectares of high evergreen forest preserved in good condition (IBUNAM 2018) (Figs. 2a, 2b).




Figure 1. Map of the Los Tuxtlas Biosphere Reserve located in the state of Veracruz, Mexico (Elaborated by Gutiérrez-García & Ricker 2011)



Figure 2. Two different aspects within the Los Tuxtlas Biological Station, UNAM (photos: Armando Aguirre)

The climate is warm and humid, with an average annual temperature between 24-26 °C, and is one of the rainiest places in Mexico, exceeding 4000 mm per year (Soto 2004, Gutiérrez-García & Ricker 2011). The area is considered a biodiversity hotspot, as it is the northernmost limit of this type of vegetation in the Neotropics and is under intense anthropogenic pressure (Gutiérrez-Granados *et al.* 2018). Historically this region has undergone extensive deforestation, which has led to a considerable reduction of natural vegetation, leaving a very fragmented landscape of high evergreen forest (Dirzo & García 1992).

Our study was conducted in a rainforest fragment of the LTBS, nevertheless, we are also conducting other comparable studies in the buffer zone. In this research we use three Malaise traps arranged in a hectare of preserved forest. These traps were monitored for 365 consecutive days, recovering every 15 days all specimens caught in each trap, which together accounted for 72 samples. All specimens were identified and quantified at family level follows Buck *et al.* (2009), and samples were preserved in 70% ethanol for subsequent studies in the Network of Multitrophic Interactions at INECOL (Xalapa, México).

Preliminary results

During an annual cycle of continuous collection in the understory of the tropical rainforest, we captured near as 68,000 specimens, belonging to 51 families of the order Diptera. The suborder Brachycera showed the greatest richness with 34 families, accounting for only 8.3% of the total abundance. In contrast, the suborder Nematocera was represented by 17 families joined together 91.7% of total captures (Table 1).

In terms of abundance, the families Cecidomyiidae and Sciaridae were the best represented by nearly 70% of all individuals. Among 10 most abundant families, we also recorded the Ceratopogonidae, Tipulidae, Mycetophilidae, Phoridae, Chironomidae, Psychodidae, Dolichopodidae and Drosophilidae. Almost one half of the families were recorded by less than 10 individuals, mostly belong to the suborder Brachycera. Despite this, it stands a single specimen of the family Inbiomyiidae, which is a first-time record for Mexico. So far, described species have been found in 8 Central and South American countries from Guatemala in the north to




Figure 3. Examples of the seasonal dynamics of three families collected along annual cycle: Dolichopodidae (top); Tipulidae (medium); Phoridae (bottom)

Bolivia and French Guiana in the south (Buck & Marshall 2006). This novel finding shows the importance of the systematic sampling conducted to assess the diptero-fauna of this region of Mexico.

We also perform analyses based on frequency-abundance indices to estimate rates of daily captures per trap (FTD). During rainy season, capture of flies was nearly four times higher compared to dry season, meanwhile, in terms of family-richness no significant fluctuation was observed, as it ranged from 60-70% between seasons.

The results of population indices exhibited several contrasting seasonal patterns, apparently depending to family assessed. For instance, the Phoridae showed two seasonal peaks during fall and spring; Dolichopodidae populations observed a single trend of growth during spring; while population indices of Tipulidae augmented between fall and winter. Such changes of the seasonal dynamics could be directly linked to certain climatic features, but it could be also explain based on specialized feeding habits for each taxon which is under analysis (Fig. 3).

Table 1. List of the families of Diptera found in the understory of the Los Tuxtlas Biological Station UNAM, Mexico.

Diptera Family	Rel-Abundance	Diptera Family	Rel-Abundance
Agromyzidae	< 10	Lygistorrhinidae	< 10
Anisopodidae	< 10	Micropezidae	11 - 100
Anthomyiidae	< 10	Milichiidae	11 -100
Asilidae	11 - 100	Muscidae	11 - 100
Bibionidae	< 10	Mycetophilidae	501 - 5000
Calliphoridae	< 10	Neriidae	< 10
Carnidae	< 10	Periscelididae	< 10
Cecidomyiidae	> 5000	Phoridae	501 - 5000
Ceratopogonidae	501 - 5000	Platystomatidae	< 10
Chironomidae	501 - 5000	Psychodidae	501 - 5000
Chloropidae	11 - 100	Rangomaramidae	101 - 500
Clusiidae	11 - 100	Rhagionidae	11 - 100
Conopidae	< 10	Sarcophagidae	11 - 100
Culicidae	11 - 100	Sciaridae	> 5000
Diadocidiidae	< 10	Simuliidae	< 10
Ditomyiidae	101 - 500	Sphaeroceridae	11 - 100
Dixidae	< 10	Stratiomyiidae	101 - 500
Dolichopodidae	501 - 5000	Syrphidae	< 10
Drosophilidae	501 - 5000	Tabanidae	< 10
Empididae	101 - 500	Tachinidae	11 - 100
Ephydriidae	< 10	Tephritidae	< 10
Fanniidae	< 10	Tipulidae	501 - 5000
Inbiomyiidae	< 10	Ulidiidae	< 10
Keroplastidae	11 - 100	Xylomyidae	11 - 100
Lauxaniidae	101 - 500	Xylophagidae	< 10
Lonchaeidae	< 10		

References

- Brown BV (2005) Malaise trap catches and the crisis in Neotropical dipterology. *American Entomologist* 51(3): 180-183.
- Buck M, Woodley NE, Borkent A, Wood DM, Pape T, Vockeroth JR, Michelsen V, & Marshall SA (2009) Key to Diptera families—adults. In, *Manual of Central American Diptera Volume 2* (Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE & Zumbado MA, Eds.). NRC Research Press, Ottawa, Canada. pp. 95-156.
- Buck M & Marshall SA (2006) Revision of the Neotropical family Inbiomyiidae (Diptera, Schizophora). *Contributions in Science* 512: 1–30.
- Dirzo R, García MC (1992) Rates of deforestation in Los Tuxtlas, a Neotropical area in southeast Mexico. *Conservation Biology* 6(1): 84-90.
- Gutiérrez-García G & Ricker M (2011) Climate and climate change in the region of Los Tuxtlas (Veracruz, Mexico): A statistical analysis. *Atmosfera*, 24(4): 347-373.
- Gutiérrez-Granados G, León-Villegas RI, Rodríguez-Moreno A, & Coates R (2018) Loss of connectivity in a highly fragmented tropical landscape: Use of ecological processes as functional indicators of biodiversity decline. *Biodiversity International Journal* 2(1): 00039 (DOI: 10.15406/bij.2018.02.00039).
- Hernández-Ortiz V (1993) Description of a new *Rhagoletis* species from tropical Mexico (Diptera: Tephritidae). *Proceedings of the Entomological Society of Washington* 95(3): 418-424.
- Hernández-Ortiz V and Dzul-Cauich JF (2008) Moscas (Insecta: Diptera). En, *Agroecosistemas cafetaleros de Veracruz: biodiversidad, manejo y conservación* (Manson RH, Hernández-Ortiz V, Gallina S y Mehlrether K, Eds.). Instituto de Ecología AC (INECOL) e Instituto Nacional de Ecología (INE-SEMARNAT), México, pp: 95-105.
- Hernández-Ortiz V, & Aguirre A (2015) A new species of the genus *Beebeomyia* (Diptera: Richardiidae) with observations of its biology on *Dieffenbachia oerstedii* (Araceae). *Journal of Natural History* 49(31-32): 1877-1889.
- IBUNAM (2018). La Estación de Biología dentro de la Reserva de la Biosfera Los Tuxtlas Semarnat. (<http://www.ibiologia.unam.mx/tuxtlas/reserva.htm>, last accessed 29 Sept 2018).
- Laurance WF, Useche DC, Rendeiro J, Kalka M, Bradshaw CJ, et al. (2012) Averting biodiversity collapse in tropical forest protected areas. *Nature* 489(7415): 290-294.
- Soto M (2004) El clima. En, *Los Tuxtlas: El paisaje de la sierra*. (S Guevara, J Laborde y G Sánchez-Ríos, Eds.). Instituto de Ecología y Unión Europea, A.C. Xalapa, pp. 195-198.

Predation by Asilidae (Diptera) in eastern New Mexico and Texas: a preliminary analysis of 5000 prey records

Darren A. Pollock

Department of Biology, Eastern New Mexico University, Portales, NM 88130

Introduction

Asilidae, or robber flies or assassin flies, are a dominant group of predatory insects, common and diverse in a variety of habitat types, but notably in open, grassland or prairie-like areas (Wood 1981, Dikow 2009). Adults are often large and conspicuous in their prey-capturing activities, which has led to many papers documenting prey choice in many asilid taxa. It is generally acknowledged that most species of robber flies are generalist predators, and will select prey items based not on taxonomic group, but rather local abundance and/or size and color (Shelly and Pearson 1978, O'Neill 1992, Cannings 2014, Londt and Dikow 2017). As mentioned by Londt (2006: 317), information on prey selection “mostly takes the form of isolated prey records that are difficult to gather for analysis and verification”. A very important resource is the “predator-prey” database (Lavigne 2003) which documents many thousands of worldwide asilid prey records, gleaned from relevant literature.

Due to a chance encounter in 2014 with an individual of the mydid fly *Phyllomydas phyllocerus* Bigot being consumed by a very large robber fly, *Proctacanthus micans* Schiner, I became interested in robber flies generally, and specifically, their prey preferences. This continues to the present, and has resulted in a sizeable collection of adult asilids pinned with their prey items. All of the 5,861 (as of 19 October 2018) predator-prey associations are housed in the insect collection of the Eastern New Mexico University Natural History Museum. In addition to these prey-associated asilids, general robber fly collecting has yielded another 10,000 or so specimens. The purpose of this paper is to present a preliminary analysis of the first 5,000 prey records and mention specific projects/studies underway, based on these data.


Geographic coverage and habitat types

The 5,000 predator-prey associations were collected “by hand”, using a standard, 12” diameter insect net, at 153 discrete, GPS-documented localities (Figs. 1–2). All associations are mounted on the same pin, with standard entomological label and a four-digit code label (Fig. 1); all data have



Figure 1. The 5,000th asilid-prey association: a female *Efferia varipes* (Williston) and *Polyphylla lanceolata* (Say) (Coleoptera: Scarabaeidae), Roosevelt Co., NM.

been entered into an Excel spreadsheet and comprise the following “fields” (columns): Cat. Number, State, County, locality (lat), locality (long), locality nickname, day, month, year, predator subfamily, predator genus, predator species, predator sex, prey order, prey family, prey genus, prey species, sex of prey, and notes. Virtually all flies have been identified to species, using the most recent taxonomic revisions and/or consultation with other asilid workers. Species of certain genera, e.g. *Machimus*, remain difficult to identify. The identifications of many of the prey items have only been made to order level; the process of getting all prey items determined to species will be a long-term task. A concerted effort was made to sample localities within 2 hours of Portales (Roosevelt Co.), NM while specimens further afield were collected opportunistically. Individual localities were visited randomly (though usually once every 1-4 weeks), and no attempt was made to quantify collecting effort; predator-prey associations were collected (or were attempted to be collected!) wherever and whenever they were observed. Approximately 98% of records were obtained from five counties in eastern New Mexico [Roosevelt (56%), Chaves (15%), Quay (11%), Lea (7%), and De Baca (4%)] and one in west Texas: Bailey (5%). Fig. 3 displays the numbers of records for each of the included counties.



Figures 2–3. Distribution of the 5,000 asilid-prey associations in eastern NM and west TX. Each circle represents a discrete GPS locality; numerals indicate number of specimens for each county. Maps created with gpsvisualizer.com (fig. 2 option “Google terrain”; fig. 3 option “county outlines”).

Almost all specimens were collected in open, grass-dominated semi-desert, or prairie-like habitats. No attempt has yet been made to analyze the botanical diversity at individual sites; some of the localities were treeless with scattered or uniform grass-diverse vegetation, with specimens collected from the ground (e.g., *Efferia* spp.) or swept from grasses (e.g., *Proctacanthus milbertii* Macquart) while others had taller shrubs, trees or cacti, on which were individuals of certain taxa of asilids (notably *Heteropogon patruelis* (Coquillett) and *H. cazieri* Wilcox, which were never seen on the ground).

Taxonomic representation of Asilidae

From 2014 to 2018, individuals representing 32 genera in seven subfamilies of Asilidae were collected in association with prey items. The top three subfamilies (numbers indicate specimen totals within each genus): Asilinae with 57% of all records (*Efferia* Coquillett, 1716; *Proctacanthus*

Macquart, 452; *Mallophora faultrix* Osten Sacken, 196), Stenopogoninae with 23% of all records (*Heteropogon* Loew, 814; genus nr. *Eucyrtopogon* Curran; 75, *Wilcoxia* James, 59), and Dasypogoninae with 15% of all records (*Diogmites* Loew, 385; *Saropogon* Loew, 330). Among Asilinae, individuals of all collected genera except for species of *Mallophora* Macquart and *Megaphorus* Bigot which were always collected on vegetation, are ground-resting, and therefore specimens with prey items are more easily seen and collected. *Efferia* is/was by far the dominant group of robber flies in the study area, with approximately 20 species collected in association with prey. The female: male sex ratio for the asilids representing the first 5,000 prey associations is 1.7:1. The complete taxonomic breakdown of the asilids involved in the 5,000 prey records is given in Table 1.

Prey selection

Spiders and insects form the bulk of robber fly diets (Wood 1981, Lavigne 2003, Dennis et al. 2012). The present study revealed that spiders and 13 orders of insects comprise the prey items; Table 1 shows the 5,000 associations, sorted to genus of Asilidae and order of prey items. Although analyses of these data are in their preliminary stages, some generalizations are possible. Over 98% of records were represented by six orders: Hymenoptera (30%), Hemiptera (24%), Diptera (22%), Coleoptera (10%), Lepidoptera (7%), and Orthoptera (6%). The rarest orders represented include Raphidioptera and Thysanoptera with one record each, Isoptera with two records, and Psocoptera and Odonata with seven and 11 records, respectively. In Londt's (2006) analysis of 2,000 Afrotropical asilid-prey associations, these same six orders comprised 96% of all records; however, the percentages represented by the specific orders were: Hymenoptera (26%), Diptera (23%), Coleoptera (15%), Orthoptera (14%), Hemiptera (10%), and Lepidoptera (8%).

At the generic level, predators such as *Efferia* spp. and other asilines seem to be generalists; however, *within* genera, individual species may show preferences for individual orders/families of prey. For example, we have approximately 20 species of *Efferia* with prey records; possible interspecies differences in prey preference are currently being analyzed. On the other hand, other genera seem to specialize on certain orders; this seems particularly true for Hymenoptera. For example, the Hymenoptera prey of *Mallophora* and *Megaphorus*, which seemingly are Hymenoptera mimics (refs?), represent 83% and 75% of all prey for these asilids, respectively. Two other genera of asilids whose species seem to prefer Hymenoptera are *Saropogon* and *Cerotainiops* Curran, with such prey representing 86% and 87% of all prey, respectively. *Cerotainiops abdominalis* (Brown) and two species of *Saropogon* have been discovered to be specialists on workers of *Pogonomyrmex* harvester ants, as mentioned below. As can be seen in Table 1, *Diogmites* spp. seem to prefer Hymenoptera (56%) and Diptera (26%). However, within *Diogmites*, one species – *D. pritchardi* – is potentially a specialist upon Coleoptera (Pollock, unpublished).

It is relatively straightforward to hypothesize on prey preferences, but given the large number of records obtained, the absences of prey orders also become obvious. For example, it is natural to assume that the absence of Orthoptera prey for species of *Atomosia* Macquart spp., with body lengths of about 10 mm, may be due to the prohibitively large size of orthopteran prey. However, for other groups of asilids, incompatibility in size cannot be relied upon as an explanation for the absence of specific groups of prey. Table 2 shows the percentages of the top six prey orders for the top five subfamilies of Asilidae, showing some trends. For example, Lepidoptera and Orthoptera comprise a significant component of prey for various genera of Asilinae (11% and 10%, respectively, of total prey associations) while in all other four subfamilies combined, they comprise only 2% and 3% of all prey. The Stenopogoninae and Stichopogoninae were associated with 58% and 48% hemipteran prey, respectively, while the overall percentage of these insects for the 5,000 asilids is only 24%.

Table 1. Details of 5,000 robber flies and their prey items, collected at various localities in eastern New Mexico and west Texas, from 2014 to mid-2018. Genera of asilid predators are listed alphabetically by, and within subfamily. Orders and sub-orders (for Hemiptera only) of prey items are given alphabetically, by 3- or 4-letter abbreviation: ARAN(ese), COL(egoptera), DIR(era), EPH(emeroptera), HET(emptera), ISO(ptera), ISO(ptera), LEP(teroptera), LEP(teroptera), ODO(X(nata), ORT(roptera), PSO(Xoptera), R, A, P(ulidoptera), and THY (sanoptera). Note: "HOM" represents the two orders Auchenorrhyncha and Sternorrhyncha

subfamily	genus	records	% total	ARAN	COL	DIP	EPH	HET	"HOM"	HYM	ISO	LEP	NEU	ODO	ORT	PSO	RAP	THY	
ASILINAE	<i>Egeria</i>	1716	34.32%	3	115	568	1	180	138	353	1	234	10	3	110				
	<i>Machimus</i>	61	1.22%	2	4	33		4	6	4		2	1		5				
	<i>Mallophora</i>	196	3.92%		10	6		12	5	163									
	<i>Megaphorus</i>	48	0.96%			9	1	1	1	36									
	<i>Procracanthella</i>	185	3.70%	1	9	44		15	37	53		9	3		14				
	<i>Procracanthus</i>	452	9.04%		50	124		4	1	108		61	1		103				
	<i>Promachus</i>	111	2.22%		10	36		9	3	39		3	1		10				
	<i>Thoria</i>	80	1.60%		0	26		1		0		11	3	4	35				
	DASYPOGONINAE	<i>Cophura</i>	2	0.04%					1		1								
		<i>Dogmites</i>	385	7.70%	2	28	100		5	23	217		5	1	4	0			
<i>Hedophylax</i>		17	0.34%		2	2				13									
<i>Lesomyia</i>		17	0.34%		1	5		5	2	4									
<i>Omatzabaurus</i>		2	0.04%			2													
<i>Saropogon</i>		330	6.60%		28	8		5	4	285									
<i>Atomosia</i>		27	0.54%		2	6		1	4	13								1	
<i>Cerastinops</i>		119	2.38%		10	1		3	1	104									
<i>Lampyris</i>		4	0.08%		3														
<i>Laphria</i>		2	0.04%		1														
<i>Pogonotoma</i>	1	0.02%																	
LAPHYSTINAE	<i>Laphystia</i>	12	0.24%					2	2	1									
STENOPOGONINAE	<i>Abaurus</i>	57	1.14%			19		21	10	2	1			1	3				
	<i>Bachomyia</i>	26	0.52%					14	12										
	<i>Heteropogon</i>	814	16.28%	25	190	20	9	307	195	51		4	9	0	4				
	<i>Holopogon</i>	11	0.22%			2		4	2	3									
	<i>Microstylus</i>	42	0.84%		1	4		14	4	4		0			19				
	nr. <i>Eucynopogon</i>	75	1.50%		4	10		35	10	12		3			1				
	<i>Ogriocerus</i>	48	0.96%		18	13		7	2	3			2		3				
	<i>Scelopogon</i>	21	0.42%		3	4												1	
	<i>Stenopogon</i>	3	0.06%								1								
	<i>Pilicornia</i>	59	1.18%		1	17		10	27	4					2				
STICHOPOGONINAE	<i>Stichopogon</i>	60	1.20%	6		18		1	28					7					
TRIGONOMIMINAE	<i>Holacophala</i>	17	0.34%	1		11		2	2										
Prey order totals		40	0.80%	490	1097	12	661	517	1475	2	332	31	11	323	7	1		1	
Prey order %				9.80%	21.94%	0.24%	13.22%	10.34%	29.50%	0.04%	6.64%	0.22%	6.46%	0.14%	0.02%	0.02%			

Table 2. Breakdown of the top six orders of insect prey (COLEoptera, DIPIera, HEMiptera, HYMenoptera, LEPidoptera, and ORThoptera), based on the top five subfamilies (each of which comprised > 1% of total records). The overall percentages for each of the orders for all 5,000 records is given on the bottom row for comparison.

subfamily	# genera	total records	COL %COL	DIP %DIP	HEM %HEM	HYM %HYM	LEP %LEP	ORT %ORT
Asilinae	8	2849	198 7%	846 30%	417 15%	756 27%	320 11%	277 10%
Dasyopogoninae	6	753	59 8%	117 16%	45 6%	520 69%	5 1%	
Laphriinae	5	153	16 10%	7 5%	9 6%	120 78%		
Stenopogoninae	10	1156	217 19%	89 8%	674 58%	76 7%	7 1%	39 3%
Stichopogoninae	1	60		18 30%	29 48%	4 7%		
overall order %			490 10%	1097 22%	1178 24%	1475 30%	332 7%	323 6%

Specific studies

Predation on Coleoptera.—Given my background as a coleopterist, this seemed a good first project; from 2014 to 2017, 293 Coleoptera were collected as prey of robber flies, representing 20 families. For three genera of Asilidae, beetles represented at least 10% of the total prey captures documented for each: *Ospriocerus* Loew (44%), *Heteropogon* (23%) and *Proctacanthus* (10%). The most popular beetle families as prey were Chrysomelidae, Scarabaeidae, Tenebrionidae, and Melyridae. The first manuscript (Pollock and Lavigne, in press) based on these asilid prey associations deals with 293 Coleoptera prey, representing 20 families; the Carabidae were not included in this paper and will be treated separately.

Predation on harvester ant workers.—Anecdotal evidence, and a few scattered literature records indicate that some species of Asilidae feed on workers of various species of *Pogonomyrmex*. We discovered this to be true in several species, notably in the genera *Cerotainiops* and *Saropogon*. In fact, of the 108 prey records known for *S. pritchardi* Bromley, 94% were harvester ant workers, despite the presence of a fairly rich pool of potential prey items from a variety of orders. For *C. abdominalis*, the bulk of the non-harvester ant prey items taken were done so on days where the temperatures were prohibitively high for the ants to be foraging (Pollock and Quinn, in prep.).

Predation of Asilidae by Asilidae.—Another interesting phenomenon made apparent by the collection of so many prey associations is the predation of robber flies on other robber flies. Of course, this should not be surprising given that many species are generalist predators and that robber flies can be abundant potential prey items. In fact, Londt (1993) found that Asilidae were the fourth most fed-upon group of prey, behind Acrididae, Formicidae and Scarabaeidae. Londt (1995) published data on 101 examples of asilid-asilid predation, with asilids identified only to genus. Our collections have produced 424 such associations, for all of which both asilids are identified to species (with a few rare exceptions), and their bulk indices (*sensu* Londt 1993) measured. These data will be used to answer questions on general prey selection in Asilidae (e.g., do larger predators select larger prey, and do females select larger prey than males?), and also intraspecific cannibalism. The same types of questions that Londt (1995) answered and also posed will be expanded upon, based on our larger data set.

Conclusion

As work continues on this collection, especially on identification of the prey items, it is expected that additional insights will be possible into ecology, behavior, and prey selection of robber flies in eastern New Mexico and western Texas. Not only will these specimens contribute to aspects of asilid biology mentioned above, but the large number of specimens will greatly add to knowledge (hitherto almost completely lacking in this geographic area) of seasonality, distribution, and basic taxonomy.

Acknowledgments

I thank various members of my family, especially my son George, who have accompanied me on many productive asilid-collecting field trips. For access to interesting collecting sites, I also am grateful to several enlightened land owners in Roosevelt County, NM (Jenny Clemmons, Marjorie Frazee, and Bill Roberts) and Bailey County, TX (Carlos Contreras and Greg Young). The unenlightened land owners shall remain nameless. For general encouragement and advice given to this neophyte dipterist, I acknowledge Eric Fisher, Rob Cannings, Torsten Dikow, Chris Cohen, Riley Nelson, and Jason Londt. Finally, I thank the many colleagues who have assisted with identifications of many of the 5,000 prey items.

Literature Cited

- Cannings, R. A. 2014. The robber flies (Diptera: Asilidae) of western Canadian grasslands. *In*: Giberson, D. J. and H. A. Cárcamo (eds.). *Arthropods of Canadian Grasslands (Volume 4): Biodiversity and Systematics Part 2*. Biological Survey of Canada. Pp. 269–297.
- Dennis, D. S., R. J. Lavigne and J. G. Dennis. 2012. Spiders (Araneae) as prey of robber flies (Diptera: Asilidae). *Journal of the Entomological Research Society* 14(1): 65–76.
- Dikow, T. 2009. Phylogeny of Asilidae inferred from morphological characters of imagines (Insecta: Diptera: Brachycera: Asiloidea). *Bulletin of the American Museum of Natural History* 319: 1–175.
- Lavigne, R. J., 2003. Predator - prey database for the family Asilidae (Hexapoda: Diptera). Modified 2016. Available from <http://www.geller-grimm.de/database/lavigne.htm>.
- Londt, J.G.H. 1993. Afrotropical robber fly (Diptera: Asilidae) predation of honey bees, *Apis mellifera* Linnaeus (Hymenoptera: Apidae). *African Entomology* 1(2):167–173.
- Londt, J.G.H. 1995. Afrotropical Asilidae (Diptera) 27: predation of Asilidae by Asilidae. *Annals of the Natal Museum* 36: 161–167.
- Londt, J.G.H. 2006. Predation by Afrotropical Asilidae (Diptera): an analysis of 2000 prey records. *African Entomology* 14(2): 317–328.
- Londt, J. G. H. and T. Dikow. 2017. Chapter 48. Asilidae (assassin flies or robber flies). *In*: Kirk-Spriggs, A. H. and B. J. Sinclair (eds.). *Manual of Afrotropical Diptera. Volume 2. Nematocerous Diptera and lower Brachycera. Suricata 5*. Pretoria: South African National Biodiversity Institute. Pp. 1097–1182.
- O’Neill, K. M. 1992. Temporal and spatial dynamics of predation in a robber fly (*Efferia staminea*) population (Diptera: Asilidae). *Canadian Journal of Zoology* 70: 1546–1552.
- Pollock, D.A. and R.J. Lavigne. In press (2019). Records of Coleoptera as prey of robber flies (Diptera: Asilidae) in eastern New Mexico and west Texas. *Proceedings of the Entomological Society of Washington*.
- Shelly, T. E. and D. L. Pearson. 1978. Size and color discrimination of the robber fly *Efferia tricella* (Diptera: Asilidae) as a predator on tiger beetles (Coleoptera: Cicindelidae). *Environmental Entomology* 7: 790–793.
- Wood, G.C. 1981. Asilidae. *In*: McAlpine, J.F., B.V. Peterson, G.E. Shewell, J.R. Vockeroth, and D.M. Wood (Coords.), *Manual of Nearctic Diptera, Volume 1*, Research Branch, Agriculture Canada Monographs 27: 549–573.


**First record of the genus *Scythropochroa* Lengersdorf 1926 (Diptera: Sciaridae)
from North America**

John D. Plakidas

111 Emerson Avenue, Pittsburgh, PA 15215. johnplakidas@verizon.net

One of the most challenging and fascinating groups of Diptera which I collect and document is the Sciaridae (Diptera), better known as the black fungus gnats. In the most recent catalog of the North American black fungus gnats (Mohrig et al. 2012), 25 genera and 166 species are presented. However the genus *Scythropochroa* Lengersdorf 1926 was not included in that catalog simply because the genus was not represented in any museum material studied. On July 1, 2018 I collected one male and several females of what I believed to be a *Scythropochroa* sp. The collection site was in a mature oak-beech forest about 5 km North of Aspinwall in Allegheny County, Pennsylvania. Dr. Menzel verified the identification for me, confirming the fact that *Scythropochroa* is now a new North American record. Adults of this genus are large, dark bodied flies typically 4-6 mm in length, making them relatively easy targets for collectors. Adults are followers of decay and I collected my specimens while at rest on a heavily rotting oak log. Here I present two photographs and two illustrations of this *Scythropochroa* sp. Some of the morphological features which define the genus are the large 1-segmented palpus as seen in Fig. 1 of the female head; the long R1 vein which joins the C beyond the M “fork” (Fig. 2); the male gonostylus with the 2-3 strong spines at the inner apical margin (Fig. 3); and the tegmen with laterally splayed apodemes and well-defined aedeagus (Fig. 4, gonostyli removed). One male and one female will be deposited in the National Museum of Natural History, Washington, D. C. for future study.





I wish to express my appreciation to Dr. Frank Menzel, Senckenberg Deutsches Entomologisches Institut (SDEI), Eberswalder Strasse 90, 15374 Müncheberg Germany, for verifying the genus *Scythropochroa*.

Reference

Mohrig, W., K. Heller, H. Hippa, P. Vilkamaa and F. Menzel 2012. Revision of the Black Fungus Gnats (Diptera: Sciaridae) of North America. *Studia Dipterologica* 19 (2012) Heft ½: 141-286.

Pursuing mycetophilids in northern Nevada during 2018

Robin Gray

P.O. Box 547, Winnemucca, Nevada 89446; sevenvalleysent@gmail.com

During 2018 I made an intensive effort to collect and rear Mycetophilidae from as many habitats as I could in Northern Central Nevada. I used Malaise, Emergence and Yellow Pan traps in this work. In addition I caught a considerable number of mycetophilids in EVS dry ice baited traps that I used for sampling mosquitoes as part of the mosquito abatement program that I am involved in. I put Malaise traps out in as large a variety of habitats as I was able. These included dry desert areas, riparian zones along the Humboldt River and a number of creeks, marshes in abandoned gravel pits, alkali flats (Fig. 1), willow thickets (Fig. 2), burned over areas, and islands of unburned vegetation in those areas, woodlands of juniper (Fig. 3) and of aspen (Fig. 4), and different sites on my own property, which sometimes yielded surprises. Elevations of these collecting sites ranged from 4200 to 9400 feet (Fig. 5). Some required taking a day off of work to hike into an area I was interested in, and do the same a week later to recover the trap. I tried to put these traps in remote and concealed locations where they would not be likely to be disturbed by humans. In fact, the only trap that was disturbed was the one I had set up on my own property, having been totally destroyed by cats (Fig. 6) who found it a wonderfully entertaining gymnasium. Both the emergence and the EVS traps made interesting contributions to the adults I collected during this past insect season. I tried to visit areas more than once during the season in the hope of being present when a short lived species was emerging.



Figures 1–2. Malaise traps set up on alkali flats (1, left), and in a willow thicket (2, right).



Figures 3–6. Malaise traps set up in a juniper grove (3, upper left), an aspen forest (4, upper right), and in the Santa Rosa Mountains at 9400 feet elevation (5, lower left). Figure 6 (lower right) was a trap on my property destroyed by cats.

In addition to this I attempted to rear adults out of every promising material I came across. These included mushrooms, various kinds of leaf litter, moldy cattail roots, decaying wood, material out of tree hollows and packrat nests. I put these into rearing chambers that I constructed (Fig. 7). They were of two types: one was a canning jar with the metal lid replaced by a square of cloth, the other a freezer container with ventilation holes lined with the same cloth so the interior could breathe. On the bottom of these containers I put about 2 centimeters of a 50/50 sand and peat moss mixture that I made myself. My rearing area was outside on the north side of a shed where it was shaded from the sun. As the season progressed and I got more and more containers I set up a table there and put the containers both on top of the table and beneath it.



Figure 7. My outdoor rearing chamber.

So far I have identified 177 adult specimens, collected mainly this year, to genus and I still have a considerable number to go. Of these 67% were taken in Malaise traps, most of the rest in EVS traps. I got no mycetophilids in Yellow Pan Traps. So far I have identified twelve genera among the adults. I am not sure how many species there are, but I plan on working on that this winter.

Of the twelve genera *Docosia* sp. made up 60% of the catch. It is present in a wide variety of habitats, including arid desert areas, from April through June. I caught two of these in an emergence trap I put over a rodent burrow. The edges of the trap were sealed over with dirt so that anything appearing in the killing jar had to have come from the burrow. After catching these two specimens I dug the burrow up, expecting to find nesting material that I could try rearing more adults of this genus out of. Instead I found that there was no nest in the burrow, just two droppings that I collect and put in a rearing chamber. The burrow dead ended and there were no branches coming off of it, it didn't appear to have been used by the animal that dug it. So I set the emergence trap over another animal burrow and caught a single specimen of *Megalopelma* sp. This burrow evidently had more than one entrance as large red spider wasps and a *Prionyx* wasp, among others, turned up in the trap. I caught one other mycetophilid, a *Rymosia* sp. in a partly flooded animal burrow in an abandoned gravel pit while dipping for mosquito larvae. This was very early in the season. I put a Malaise trap in this pit a number of times during succeeding months and caught no more *Rymosia* there. I caught one more adult mycetophilid with an emergence trap I put up over a patch of moss - a *Mycetophila* sp.

From the collecting I did this year I got the impression that the populations of some of these flies were very localized – I might catch several at one location and a hundred yards away in apparently identical habitat catch nothing. The adults of most of the genera I have seen don't seem to live very long - they will turn up in a trap one week, and not be seen again. Out of 19 *Exechia* sp. that I caught, only one was taken in a Malaise trap, the others were all collected in EVS dry ice baited traps. What is that telling me about them?

My efforts at rearing adult mycetophilids out of the various substrates I collected have so far not produced one adult fly, and I have seen no activity of any sort in any of the rearing chambers. I have no idea why my efforts on this have failed. Did I inadvertently kill whatever larvae might have been there while collecting the substrate? Did I collect the wrong substrate? But I collected everything that looked promising, neglecting nothing. It is a mystery to me where these flies are coming from. I've made a Berlese funnel and next year I am going to run prospective breeding material through it and see if I can't collect mycetophilid larvae. If I can, then I can target that material. But I haven't given up the breeding material I collected in 2018 - I will move the containers inside to a cool room over the winter and see if something comes out in spring.

This winter I will identify the rest of the adults I have collected to genus, and then move forward trying to get them to species. I will try to find keys and descriptions to do this. I don't think it will be an easy task. In the spring I will again start to collect adults and will run potential breeding material through my Berlese funnel. If I can find the time I will go out and search for adults with a net. I have not been able to get to every habitat in North Central Nevada that mycetophilids might be in, and I don't want to make assumptions about what those may or may not be because that just leads to blind spots. I think I may have missed some species because they emerge, live only a few days, and then are gone. I might pick those up if I kept a trap in one location for the entire insect season. I might try that in a promising location, and keep moving the other traps I have around.

This has been a very challenging, absorbing and interesting project and I feel I have learned a great deal about not only mycetophilids, but other insects as well. And the learning has been through direct engagement with the insects in the wild, trying to understand the clues they are giving me about their hidden, secret lives. I'm greatly looking forward to 2019.

The Gnatwork

Emma Howson

The Pirbright Institute, Ash Road, Pirbright, Woking, Surrey GU24 0NF, UK; emma.howson@pirbright.ac.uk

The Gnatwork, directed by Dr Simon Carpenter (The Pirbright Institute) and Prof Mary Cameron (LSHTM), is one of four Global Challenges Research Fund (GCRF) Networks in vector-borne disease research. This project brings together workers on blackflies (Simuliidae), sandflies (Psychodidae) and biting midges (Ceratopogonidae) from around the world to address technical issues that arise when working on small biting flies. As part of the project, we have funds available to pump-prime collaborative research projects that bring together researchers and techniques for working across the three vector groups (the next call will open late 2018). We will also be hosting a number of international training workshops aimed at early career researchers, and provide web-based resources, including re-releasing the information that was originally held at www.culicoides.net. For access to these resources and opportunities, join the Gnatwork at www.gnatwork.ac.uk.



HISTORICAL DIPTEROLOGY

Revised dates of publication for Schiner's "*Fauna Austriaca. Die Fliegen (Diptera)*"

Neal L. Evenhuis

J. Linsley Gressitt Center for Entomological Research, Bishop Museum,
1525 Bernice Street, Honolulu, Hawaii 96817, USA; neale@bishopmuseum.org

Schiner's *Fauna Austriaca. Diptera* (Schiner 1860–1864) was published in two volumes, with the title page for having the dates “1862” (for Vol. I) and “1864” (for Vol. II). However, the work was actually published in 14 Hefte (including double Hefte) between 1860 and 1864. In Evenhuis (1997), I gave publication dates as researched at that time. Recently, I came across a recording journal of Austrian book publishers (the *Österreichische Buchhändler-Correspondenz*), which gives more accurate dates for the Hefte. Most of the dates are earlier than previously recorded, and in two cases (Hefte 3/4 and 9/10), involve publication in the year previous to what was known in 1997. I here present the breakdown of each Heft and earliest known publication dates based on current research.

Table 1. Dates of Publication of Schiner's *Fauna Austriaca. Die Fliegen (Diptera)*.
(based on dates in the *Österreichische Buchhändler-Correspondenz* unless otherwise noted)

Vol.	Heft	Pages, plates	Date of publication	Source
I	1	1–72	10 Jun 1860	
	2	73–184, pl. 1	20 Sep 1860	
	3/4	185–368	1 Dec 1860	
	5	369–440	10 Aug 1861	
	6/7	441–656	20 Nov 1861	
	8 (part)	657–674, prelims, pl. 2	2 Jul 1862	(Zool.-Bot. Ges. Wien)
II	8 (part)	1–80	2 Jul 1862	(Zool.-Bot. Ges. Wien)
	9/10	81–288	1 Dec 1862	
	11/12	289–480	1 Nov 1863	
	13/14	481–658, prelims	March 1864	(Wien. Ent. Monatschr.)

References

- Evenhuis, N.L. (1997) *Litteratura Taxonomica Dipteriorum (1758–1930) being a selected list of the books and prints of Diptera taxonomy from the beginning of Linnaean nomenclature to the end of the year 1930; containing information on the biographies, bibliographies, types, collections, and patronymic genera of the authors listed in this work; including detailed information on publication dates, original and subsequent editions, and other ancillary data concerning the publications listed herein*. 2 vols. Backhuys Publishers, Leiden, x + 871 pp.
- Schiner, J.R. (1860–1864) *Fauna Austriaca. Die Fliegen. (Diptera). Nach der analytischen Methode bearbeitet von J. Rudolf Schiner. Mit der Charakteristik sämtlicher europäischer Gattungen, der Beschreibung aller in Deutschland vorkommenden Arten und dem Verzeichnisse der beschriebenen europäischen Arten*. 2 vols in 14 Hefte. C. Gerold's Sohn, Wien [= Vienna].

**Gems from the H. Fischer collection: Diptera watercolors by E. H. Compton
and a syntype of *Pegesimallus teratodes* (Hermann, 1906) (Asilidae)**

Marion Kotrba

SNSB Zoologische Staatssammlung München, Münchhausenstraße 21,
81247 München, Germany; marion.kotrba@zsm.mwn.de

The SNSB Zoologische Staatssammlung München (ZSM) recently received the natural history collection of Dr. Heinrich (“Heinz”) Fischer (* 5. iv 1911, Augsburg, Germany; † 15. iv 1991 *ibid*). Fischer was a natural history “all-rounder” interested in zoology and particularly entomology, but also ecology, geography, and archeology. He received a broad academic education in these topics at the Ludwig Maximilian University of Munich. After graduating in 1938 he was temporarily employed at the ZSM and later worked as a teacher and as a freelance scientist, all the time amassing an extensive natural history collection with a special focus on less studied groups such as Diptera. In 1983 Fischer founded a natural history museum in Königsbrunn and his collections were originally deposited there. Later most of the collection was placed into the temporary care of the Naturmuseum Augsburg. In 2016 the collection H. Fischer was finally transferred to the ZSM.

The received collection included 76 drawers of Diptera, mostly from Bavarian locations, but also with a few exotic specimens. Four of these drawers are particularly noteworthy, featuring some exceptional exemplars accompanied by beautiful illustrations in watercolor executed by E. Harrison Compton (* 11. x 1881, Feldafing, Germany; † 6. iii 1960 *ibid*). Compton is renowned for his alpine landscapes. The dipteran details demonstrate that he also excelled in this entirely different subject. The four drawers with Compton’s watercolors are illustrated here with translations of the accompanying text labels.

One of the drawers contains a well-preserved specimen of *Pegesimallus teratodes* (Hermann, 1906) (Asilidae) whose hind legs are beautifully adorned with very long, laterally compressed, feather-like bristles along the anterior and posterior margins of the much elongated hind tarsi. The impressive specimen, which is labelled as “Type” is larger and much better preserved than the type specimen of *Pegesimallus teratodes* already present in the ZSM collection, which was previously considered the unique holotype (Londt 1980). By reference to the original description by Hermann (1906) it is established that the specimen from the H. Fischer collection is indeed a syntype. Some additional information on the provenience of the type material is provided.

Drawer “C 33 Fliegen (Compton 1)” (Fig. 1, ZSM Drawer-ID: ZSM-DIPT-40/o.Nr.)

Watercolors and respective exemplars as identified on the labels: on the left, top to bottom: *Haematopoda italica* (Tabanidae, head and thorax), *Laphria flava* (Asilidae, lateral aspect of head and prothorax, tarsi of right foreleg), *Sargus cuprarius* (Stratiomyidae, lateral aspect of head and prothorax, frontal aspect of head); on the right, top to bottom: *Ogcodes gibbosus* (Acroceridae, lateral aspect), *Bombylius major* (Bombyliidae, lateral aspect).

Text labels [*translation*], top to bottom: “Die Bremsen saugen gerne Blut – und werden oft dafür erschlagen.” [*The horse flies like to suck blood – and are often slain for it*].

“Die Raubfliegen jagen andere Insekten und saugen sie aus.” [*The robber flies hunt other insects and suck them dry*].

“Die Waffenfliegen sind harmlose Tiere, obwohl sie mit zwei kräftigen Dornen am Schildchen bewehrt sind.” [The soldier flies are harmless animals, although they are adorned with two strong thorns on the scutellum].

“Die Buckelfliegen scheinen ohne Nahrungsaufnahme zu leben. Ihre Larven aber fressen Spinnen oder deren Eipakete auf.” [The hunch-back flies seem to live without food intake. Their larvae, however, devour spiders or their egg packets].

“Die Wollschweber sind äusserst gewandte Flieger. Wie Kolibris saugen sie – in der Luft stehend – den Blütenhonig. Ihre Larven schmarotzen bei der Seidenbiene (*Colletes*), fressen erst deren Blütenstaubvorrat und dann ihre Brut dazu.” [The bee flies are extremely swift fliers. Like colibri they suck – hovering in the air – the flower honey. Their larvae parasitize on the plasterer bee (*Colletes*), devouring first their pollen deposit then their brood to boot].




Figure 1. Drawer “C 33 Fliegen (Compton 1)” [flies (Compton 1)] from the H. Fischer collection.

Drawer “C 34 Harmlose, Licht u. Feuchtigkeit liebende Pflanzenbesucher“ (Fig. 2, ZSM Drawer-ID: ZSM-DIPT-232/o.Nr.)

Watercolors and respective exemplars as identified on the labels: on the left, top to bottom: *Trypetoptera punctulata* (Sciomyzidae, dorsal aspect), *Stratiomyia chamaeleon* (Stratiomyidae, dorsal aspect of head and thorax); on the right: *Limnia unguicornis* (Sciomyzidae, lateral and dorsal aspect of head and thorax).

Text labels [translation], top to bottom: “Die Hornfliegen sind meist auf Pflanzen in der Nähe von Wasser zu finden. Die Larven u. Puppen schwimmen auf der Oberfläche von stehenden oder langsam fließenden Gewässern zwischen Wasserpflanzen, von denen sie leben. Die Puppen finden sich im Genist, frei oder in Schneckenschalen.” [The horn flies are generally found on plants in the vicinity of water. The larvae and pupae float on the surface of standing or slow-flowing bodies of water between water plants, of which they live. The pupae are found in litter, free or in snail-shells].

“Eine Hornfliege.” [A snail killing fly].

“Harmlose, Licht u. Feuchtigkeit liebende Pflanzenbesucher.” [Harmless light and humidity loving visitors of plants].

“Die Waffenfliegen sind Honigsauger und Pollenfresser; sie lieben besonders Schirm- und Korbblüten und sitzen auch gerne auf sonnenbeschienenen Büschen in Wassernähe.“ [The soldier flies are honey-suckers and pollen-eaters; the particularly love umbellifer and composite flowers and also like to sit on sunlit shrubbery in the vicinity of water].

“Eine Waffenfliege.“ [A soldier fly]




Figure 2. Drawer “C 34 Harmlose, Licht u. Feuchtigkeit liebende Pflanzenbesucher” [Harmless light and humidity loving visitors of plants] from the H. Fischer collection.

Drawer “C 35 Schmarotzerfliegen und ihre Opfer” (Fig. 3, ZSM Drawer-ID: ZSM-DIPT-27/o.Nr.)

Watercolors and respective exemplars as identified on the labels: on the left: *Gymnosoma rotundata* (Tachinidae, lateral and dorsal aspect); on the right: *Echinomyia fera* (Tachinidae, lateral aspect of the head).

Text labels [translation], top to bottom:

“Schmarotzerfliegen und ihre Opfer.” [*Parasitic flies and their victims*]

“Die Kugelfliege.” [*The bullet fly*] (German vernacular name)

“Die Borstenfliege.” [*The bristle fly*] (German vernacular name)




Figure 3. Drawer “C 35 Schmarotzerfliegen und ihre Opfer” [*Parasitic flies and their victims*] from the H. Fischer collection.

“Aussergewöhnlich fruchtbar, setzen die schwarzen Borstenfliegen (1) sehr kleine gepanzerte Larven in der Nähe der Eichenspinner-Raupen (2) ab. – Die Larven lauern den Raupen auf, um sich in deren Inneres einzubohren. – Die erwachsene Raupe spinnt sich noch eine Puppenhülle (3), sie ist aber im Innern aufgefrisst. In ihrer Hülle verpuppt sich die Fliegenlarve. – An Stelle des Schmetterlings (Männchen (5); Weibchen (6)) schlüpft nun die Fliege (4). – Weil der Mensch in vielen

Schmetterlingen Schädlinge sieht, bezeichnet er die Schmarotzerfliegen als Nützlinge.”
 [Extraordinarily prolific, the black bristled flies deposit very small armor-clad larvae in the vicinity of oak egg caterpillars. – The larvae ambush the caterpillars, in order to bore into their interior. – The mature caterpillar will still construct a cocoon for itself, but it is already eaten up inside. Within its shell the fly larvae pupates. – In place of the butterfly (male (5), female (6)) the fly (4) emerges. – Because man considers many butterflies as vermins, he calls the parasitic flies beneficial].

Drawer “C 37 Fliegen aus China und Afrika” (Fig. 4, ZSM Drawer-ID: ZSM-DIPT-37/233)

Watercolors and respective exemplars as identified on the labels: on the left, top to bottom: *Diopsis nigricollis* (Diopsidae, oblique aspect of head and thorax), *Diopsis thoracica* (Diopsidae, oblique aspect of head and thorax); on the right, top to bottom: *Hermyia diabolus* (Tachnidae, two aspects of the head), *Neolaparus (Lagodias) teratodes* (Asilidae, detail of the male right hind leg tarsi).

Text labels [translation], top to bottom:

“Fliegen” – “aus China und Afrika” – “Aquarelle von E. H. Compton”. [Flies – from China and Africa – watercolors by E. H. Compton].




Figure 4. Drawer “C 37 Fliegen aus China und Afrika” [Flies from China and Africa] from the H. Fischer collection.

The red label pinned below the specimen of *Neolaparus (Lagodias) teratodes* reads “Ikuta (Africa orient.) / *Neolaparus (Lagodias) Lw* / *teratodes* Harm. ♂ / Type”. The specimen itself is labelled “Ikuta / Ost-Afrika” and “*Neolaparus* ♂ / *teratodes* Harm.”

Type material of *Pegesimallus teratodes* (Hermann, 1906)

The species was described in 1906 under the name *Lagodias teratodes* by Friedrich Hermann (* 07. iii 1859, Neapel, Italy; † 27.ii.1920, Erlangen, Germany), stating “Die Art ist durch die geradezu mōnströse federartige Beschuppung an den Tarsen der Hinterbeine auf den ersten Blick kenntlich und wird dadurch zu einer der interessantesten und auffallendsten Asilidenformen, die bisher bekannt geworden sind.” [*The species is discerned at first glance by the downright monstrous feather-like scales on the hind leg tarsi and thus becomes one of the most interesting and conspicuous forms of asilids so far recognized*].


Londt (1980) redescribed the species and included it in the genus *Pegesimallus*. Londt could study a single specimen of *teratodes* (“Holotype ♂, Kenya, Ikutha, vi.1906, Prof. Hauser, ZSM”) which he received on loan from the ZSM and considered to be the unique holotype. He notes the meagre information about the provenience of the specimen: “I have not been able to trace the locality Ikutha or anything which looks similar. As Hermann states that it is in 'englisch Ostafrika' I must assume he was referring to present-day Kenya.”



Figure 5. *Pegesimallus teratodes* (Hermann, 1906), syntype 1 (ZSM) previously considered the unique holotype. Lateral and dorsal aspect, labels.

Said specimen had been deposited at the ZSM with the collection of Friedrich Hermann in 1920. Today we find it in rather poor condition, lacking both wings, the terminal 3 tarsal segments of both hind legs and the terminal segments of both antennae (Fig. 5). The pin bears a square of clear plastic with the cleared terminalia in a drop of yellowish medium and four labels reading top to bottom: [red label] “Ikutha/ *Neolaparus*/ *teratodes*/ Type. Ham”, “Sammlung/ F. Hermann”, [red label]

“Type von/ *Neolaparus teratodes* Herm”, “*Pegesimallus/ teratodes* ♂ (Hermann 1906)/ det. J. G. H. Londt, 1978.”



Figures 6 and 7. *Pegesimallus teratodes* (Hermann, 1906), syntype 2 (H. Fischer collection). Dorsal (6A, top) and lateral (6B, middle) aspects, with labels, and illustration from Hermann (1906, “Fig. 8. *Lagodias teratodes* n. sp.”) (7, bottom).

Considering the poor condition of this supposedly unique specimen it is particularly fortunate that the specimen so prominently featured in the Fischer collection (Fig. 6) turns out to be a syntype. Not only does the label next to it declare it as a “Type” and from the type location Ikutha. Also Hermanns (1906) species description specifies that he possessed two male specimens. (“Da ich von *Lagodias teratodes* nur zwei Männchen besitze, ...” [as I only possess two males of *Lagodias teratodes*, ...]). Most importantly, his illustration “Fig. 8. *Lagodias teratodes* n. sp.” (our Fig. 7), drawn after a photograph, undeniably shows the specimen from the Fischer collection. It has to be noted, however, that the handwriting on the red type labels of the ZSM type and the red label pinned next to the specimen in the Fischer collection seem not to be from the same hand.

Some information can be added regarding the provenience of these specimens. The type locality “Ikutha, englisch Ostafrika” most likely refers to Ikutha (2°03'60"S 38°10'47"E), a semi-arid town in Kitui County, Kenya, ca. 150 km southwest of Nairobi. It is reported that the Leipzig Evangelical Lutheran Society opened a mission station at Ikutha in 1892.

Hermann (1906) states that the specimens were sent to him together with Coleoptera by a collector “Prof. Hauser” of whom, at the time of the description, he had not heard in years. In the Biographies of the Entomologists of the World Groll (2017) lists three entries for the name Hauser that fall into the appropriate time frame before Hermanns publication in 1906:

(1) Fritz Hauser (* 28. xii 1868, Vienna, Austria; † 26. viii 1910, Plomberg, Austria): An obituary by Rebel (1911)

addresses him as “Herr Fritz Hauser” [*Mr. Fritz Hauser*] and “Assistent” [*assitant*] at the physiological institute of the university. It can be ruled out that the academic title “Professor” was omitted in an obituary written in Austria at that time. This proves that Fritz Hauser never carried the title and Hermann would not have referred to him as “Professor”. Moreover Fritz Hauser's entomological activities were restricted to Lepidoptera. This renders it highly unlikely that Hermann received the type specimens from him.

(2) Friedrich Hauser (* 30. vi 1853 Nördlingen, Germany; † 00.00.1932, Erlangen, Germany), Colonel.

(3) Gustav Chr. F. Hauser (* 13. vii 1856, Nördlingen, Germany; † 30.vi.1935, Erlangen, Germany), Prof. Dr. of anatomy, “Geheimrat” [*privy councillor*].

Friedrich and Gustav Hauser were brothers and both amateur entomologists mostly interested in Lepidoptera and Coleoptera (Roß 1935). Only for Gustav Hauser it is documented that he later diversified into other Insect groups, particularly Diptera (Roß 1935). Moreover only Gustav Hauser carried the academic title “Professor”. No record was found documenting that either brother visited Africa. Groll (2017) only states that Friedrich Hauser collected in Turkestan and also occasionally dealt in insects.

In any case it appears most likely that Hermann received the specimens via “Professor” Gustav Hauser. How the one syntype finally ended up in Fischers collection is more mysterious. When Hermann died in 1920 Fischer was only 9 years old, a child unlikely to have received the precious specimen directly from the author. Also it appears quite unlikely that Hermann gave away the better one of only two type specimens he had of that species. Or that he returned it to the collector Professor Hauser, of whom he had not heard in years. It is therefore most conceivable that both syntypes were originally received by the ZSM with the Hermann collection in 1920. Fischer may have come across the spectacular specimen when he was employed at the ZSM, picked it for illustration by Compton, and kept it in the display drawer together with the watercolor. Happily, the two syntypes of what is today *Pegesimallus teratodes* are finally reunited and readily available for further scientific study in the ZSM collection.

References

- Fischer H. 1963: 83. Die Tierwelt Schwabens, 7. Teil: Raubfliegen. Bericht der Naturf. Ges. Augsburg 16: 35-46
- Groll E. K. 2017: Biographies of the Entomologists of the World. Online database, version 8, Senckenberg Deutsches Entomologisches Institut, Müncheberg; URL: sdei.senckenberg.de/biografies
- Hermann F. 1906. Beitrag zur Kenntnis der Asiliden (II) (Diptera). Zeitschrift für Hymen. und Dipterologie 1906: 129-144
- Londt J. G. 1980: Afrotropical Asilidae (Diptera) 4. The genus *Pegesimallus* Loew, 1858 (= *Lagodias* Loew, 1858; *Neolaparus* Williston, 1889), including species from other zoogeographical regions and the descriptions of the two new genera, *Brevirostrum* and *Caroncoma*. Annals of the Natal Museum 24: 233-347
- Rebel H. 1911: Obituary Fritz Hauser. Verh. Zool.-bot. Ges. Wien 61: 25-27
- Roß E. 1935: Über bedeutende Koleopterologen und den Verbleib ihrer Sammlungen. Internationale Entomologische Zeitschrift, Guben 28: 484-485

Reproduction of Loew's 1859 Instructions for Collecting Diptera

Neal L. Evenhuis

J. Linsley Gressitt Center for Entomological Research, Bishop Museum,
1525 Bernice Street, Honolulu, Hawaii 96817, USA; neale@bishopmuseum.org

In gathering information for a biography of Hermann Loew, the following was found among his many (over 270) publications I had gathered together years ago, and was initially overlooked, only because it did not contain any taxonomy. Although ~160 years old, many of the techniques are still used today — only the equipment has changed or become modernized. When rediscovering the article in his bibliography I was intrigued as to what he might have said about collecting since I had just finished writing about his 1841–1842 trip to “Asia Minor” [essentially just southwestern Turkey] where he was said to have collected thousands of specimens yet only had a few months collecting time (in between doing archeological work with his colleague August Schönborn) and these insects were all collected without traps that we use so commonly today.

This article is one of many articles on instructions for the collecting of various groups of plants and animals—each written by a specialist on the group—that were included in the *Annual Report of the Smithsonian Institution for 1858* (published 1859). The introduction to all the articles on insects (the first of the following 8 pages) is added here to explain Loew's article and put things in context for the time, i.e., the young fledgling Smithsonian Institution was just embarking on an ambitious goal of producing catalogues and inventories of the natural history of the United States. The first *Catalogue of the Diptera of North America* had just been published by Osten Sacken the year previous and the new series on the *Diptera of North America* to be headed by Loew was to begin shortly.

INSTRUCTIONS FOR COLLECTING INSECTS.

INTRODUCTORY REMARKS.

The Smithsonian Institution has undertaken the preparation of a series of reports on the insects of North America, with the view of attracting attention to and facilitating the study of a department of natural history, which, though so much neglected, is yet of great practical importance to the agriculturist and rural economist. These works will consist—first, of catalogues of all the known genera and species of each order, with reference to the places of publication ; and secondly, of monographs and detailed descriptions of both old and new species, so as to enable any one interested in the subject to identify the insects met with, and to prosecute investigations respecting them.

Towards the accomplishment of this object the Institution has already published a catalogue of the North American Coleoptera, by Dr. F. E. Melsheimer, and also one of the Diptera, by Baron Ostensacken, while similar catalogues of Neuroptera, Hymenoptera, Hemiptera, and Diurnal Lepidoptera are in an advanced state of preparation. It is now proposed to take up a very interesting and extensive group, that of the Moths or Night Butterflies, and contributions of materials towards a work on this subject are, therefore, respectfully solicited. There is no locality in the country where new species may not be readily obtained, and from which complete collections are not desirable, although novelties are most to be looked for from the regions of the west. From every region the most common and familiar species are equally desirable, with the rarest and most striking ; the object being not merely to collect all the species, but also to determine accurately their geographical distribution.

In the following pages will be found special instructions for collecting insects, furnished for this article by different entomologists, with some general remarks on the subject taken from Samouelle's Entomologist's Companion.

INSTRUCTIONS FOR COLLECTING DIPTERA.

BY H. LOEW, WITH ADDITIONS BY R. OSTENSACKEN.


The first instrument I use in a locality, the fauna of which I am not well acquainted with, is a linen bag, fourteen inches in diameter and two and a half feet long, a little rounded at the top or apex, and attached to a strong iron ring, with a hollow metallic handle, in which, if necessary, a cane or other piece of wood may be inserted. With this bag I begin to sweep, and in a short time ascertain what insects abound in the grass, on the flowers, or in moist or dry places, &c. After the last sweep, by a dexterous twist of the handle I throw the apex of the sack over the ring, so as to prevent the escape of the insects, and give them time to compose themselves at the bottom. It is then opened, and I insert my head into it to see what I have captured. They immediately begin to move upwards. The common specimens I allow to pass; but if there is one which I desire to take, I let it advance to the middle of the bag, and then gently grasp it, from the outside of the bag, with the right hand. The other captives I drive down to the bottom by blowing moderately on them, and confine them there by letting the ring fall over the right hand, which holds the insect, which is now easily seized with the left hand. This operation is repeated until everything desirable is taken out, when the bag is emptied of its contents by inversion, and the sweeping recommenced.

This is the most important of all the implements used by the Dipterologist. Its employment prevents him from wasting his time in barren localities, promptly indicates the presence of rare species, and, by furnishing this information, enables him afterwards to have an abundance of specimens.

When I have, by the use of the bag, discovered the locality of rarities, I use another instrument, which is nothing more than an ordinary insect-forceps, of the form shown in fig. 1. Here, *b* is the place for the thumb, *c* is the joint placed near the handle in order that the

forceps may be closed rapidly; the remaining portion of the arms should be made so stout that they will not spring when closing the blades; *d* is a pin to prevent lateral motion when the blades are closed;

Fig. 1.



a the oval ring must be covered with a double cotton tape tightly sewed. If the forceps are not more than nine or ten inches long the ring need scarcely be bound with tape. Its angle and straight side are useful in

capturing insects in particular situations. Then the space between the rings, on their internal side, must be covered with the best bobbinet, tightly drawn and securely sewed, so that a perfectly level surface be formed, in order to prevent the captured insect from struggling. The insect is pinned through the openings of the bobbinet, and is then easily transferred to the collecting box, or stuck into the sides or top of your hat, at pleasure. Two views of the instrument are given to illustrate its form more fully.

This instrument is not adapted for very small insects. For such, a forceps with finer bobbinet is used, but even this is superfluous, if you have become expert in employing the bag in sweeping, as well as in holding it under plants, bushes, and flowers, and, by gently striking or shaking them, allowing the insects to fall into it. Neither does it answer for the capture of very active species. It is, for the most part, useless to pursue such, and only betrays the intemperate zeal of the beginner. It is better to ascertain where such species pass the night, and thus you can easily capture that with your fingers which at other times would cost much labor. An excursion early in the morning, before the dew is off, is often very productive.

To carry more instruments with you than those just mentioned is an unnecessary burden.

Besides the capture, the *breeding* of Diptera, is of the greatest importance, and, for the most part, easy. The principal rule I follow, in order to secure the perfect insect, is, to be in no hurry about taking the imperfect one. If there is reason to presume that the larvæ will change into the pupa state at the place where it is found, I wait until then and gather the pupas. On a contrary presumption, I take the larvæ only when I observe that they are preparing for the pupa state. Decaying wood, mould in hollow trees, manure, and ground-earth under manure, stems of plants, grass, stalks of hollow weeds, dried flowers and their seed vessels, particularly of the *Compositæ*, furnish me the richest booty.

A bag full of the dried flowers of the *Compositæ*, or a bundle of dry burdocks and thistles, which I have had gathered for a few pennies, has always furnished me with something good. The pupæ of Diptera do

not require much care ; only do not let the place in which they are kept be too warm. Those I wish to separate are put in large glass jars, the tops of which are covered with paper. The others I place in a wooden box, which I keep in an unheated room, on the window-sill. When they are developed from the pupa towards spring, they naturally fly to the window, from which I take them with a glass vial, or, what is better, with a small chip-box furnished with a glass bottom.

The third point of great importance is the preparation and pinning of the specimens for the cabinet. Sulphuric ether and chloroform are but inferior means of killing them. Specimens too long exposed to these agents are apt to spoil, and, if too little exposed to them, they revive. They answer only for certain tender minute species, of which I shall speak below. The best means of killing them are kreosote and the smoke of a strong cigar. The general rule is, to pin the flies whilst living, and thus to put them into the collecting box, which should have the bottom well moistened with kreosote, and be made tight. The captives will soon die, and thus, time after time, fresh subjects may be put in as they are caught. I have had boxes in which the largest Diptera were almost immediately killed, although the kreosote had not been renewed for six days, and which were still fatal to smaller species for six months afterwards. If you have no boxes thus prepared with kreosote, or, if you wish to avoid the odor of it, then prepare a box, so that, when it is full of pinned specimens, you may blow a few strong puffs of cigar-smoke into it, and all life will soon cease.

In the pinning of specimens, the long pins are greatly to be preferred to the short ones. Five-sixths of all the entomologists now use the long pins. $m n$, fig. 2, is the normal length of pins which are used for Diptera ; the finer kinds may be of the length $m d$. The insects must be pushed high enough up the pin to enable the surface of its back to be examined with an ordinary lens without being incommoded by the pin's head. The back of the insect must, of course, be a little nearer the pin's head than the length of the focus of the glass. As smaller insects must be examined by glasses of higher power, they must be brought nearer the pin's head than larger ones.

Large flies may be pushed up so that their backs come to b , d which smaller ones should reach a . Since the beautiful magnifiers of Oberhauser, which allow the application of a lower power, have come into use, insects may be pinned at a greater distance from the head, but not much greater.

The minute and tender species, such as the *smallest Cecidomyice*, *Campylomyza*, and others, must be treated differently. They cannot be pinned in the ordinary way with safety. Van Heyden has proposed the most rational and elegant method. The smaller species are caught by the linen bag, and are then put into very porous chip pill boxes, and are killed with cigar smoke, if boxes moistened with kreosote are not preferred. When twenty or thirty are captured the box is turned up so that specimens fall into the cover. They are then transfixed from below on fine silver wire, but not so as to allow the wire to project beyond the thorax. The silver wire must previously

Fig. 2.



be cut with ordinary scissors into pieces of proper length and taken with you. The wire must be inserted into the insect by means of a very delicate pincers, and with the same instrument the other end of the wire must be stuck into a box, the bottom of which is covered smoothly with the pith of plants. Over the pith thin paper may be fastened in order to render the surface still more smooth. The top or cover of the pill box, in which the insects are placed while transfixing them, must not be too shallow, so that they may not be blown away by the wind, if the operation is performed in the field. The use of a glass vial for the capture of small insects is not to be recommended, as the moisture separates their parts and thus spoils them. The vial may be used on excursions, when you desire to take home a single living insect for the purpose of examining the organs of its mouth or its internal structure.

Those specimens which have been transfixed on the silver wire, as above indicated, are prepared for the collection in the following way: In the fall, stalks of weeds with pith not too porous, and before it is quite dry, must be collected. The best adapted for this purpose is the pith of *Artemisia*, and, still better, of *Verbascum*, (mullein.) It is cut into small




Fig. 3.  pieces, which, seen from above, have the form and size of


Fig. 4.  fig. 3, and seen from the side the thickness of fig. 4. Then

insect pins of moderate size are inserted through these

Fig. 5.  small pieces at the place indicated by the dot, (fig. 5,) and the pieces are pushed up nearly to the pin's head. As the pith is still moist it will adhere to the pin in drying, and by the slight coat of verdigris which will soon be developed, the insects, which have been previously attached to the silver wire are now inserted on the free end of the pith with the delicate pincers, where they will stick fast enough if the proper kind of pith has been used. Pith of *Sambucus*, or elder, *Helianthus*, or sunflower, and similar plants, do not answer the purpose, because it is too porous. The insect should stand a little

higher than the head of the pin, so that it may be conveniently examined from all sides. The silver wire used for this purpose should be of the very best quality so as to prevent the development of verdigris.

In regard to the *Tipularia*, and some others, whose legs easily break off, my method is the following: I capture them with the forceps and pin them as other large specimens. Before pinning, I press the thorax, so as to prevent all struggling; I then stick the pin through a piece

Fig. 6.  of paper of this form, (fig. 6,) and push it up far enough to receive the legs of the insect neatly spread out or turned under it. When the insect is dead I remove the paper, which, however, might as well be suffered to remain.

The boxes in which Diptera are kept should be about a foot square, the top should be glass, and made to fit tight to the other part by a double rab-


bet. Some persons, in addition, fasten the cover down by means of brass hooks, to prevent warping, (fig. 7.) The boxes may be made of any well seasoned wood. The bottom of the boxes should be made of wood cut *horizontally* from the trunk in plates one-fourth to one-third of an inch thick, and in this case we may use the harder kinds of wood from deciduous leaved trees, except when the rings making the last growth of the year are very hard, or else a very soft wood may be used, such as the partially decayed wood of dead linden or poplar trees; but the pieces are cut out lengthwise in the ordinary way. In both cases the finest pins can be easily inserted into the wood without risk of injuring their points and hold very well. But in the latter mentioned kind of bottom a fine pin readily takes a wrong direction, and after having been taken out it is difficult to insert it again in a perpendicular position. The use of soft material for lining the bottom of the boxes has been generally abandoned in the best collections. It is only when *perfectly* tight boxes cannot be procured that the lining is used, and then not so much with reference to the insertion of the pin as to facilitate the poisoning of the box and to exclude the *acari*, psoci, and other insects. The best material for this purpose is blotting paper spread over with mercurial ointment, and over this is laid a fine oiled silk of light color. The oiled silk may be faintly ruled with cross lines to facilitate the symmetrical arrangements of the specimens. If the oiled silk is fine enough it offers no perceptible resistance to the finest pin. But a more serious inconvenience is, that it prevents us from seeing the texture of the wood at the place where the pin is to be inserted, and hence the points of the very fine pins are liable to injury. On this account it is much better to use wooden boxes without any lining, and these when neatly made look well enough. But if it is deemed desirable to still further improve the appearance of the boxes, a sheet of thin paper may be spread over the bottom, provided the texture of the wood is very uniform or very fine pins are not to be used. This paper may be renewed from time to time, and should be fastened by glue and not by paste, which only furnishes a bait for destructive insects.

If it is deemed advisable to poison the boxes, it is most conveniently done by means of bits of felt smeared with mercurial ointment, which are fastened in the covers. During the examination of the insects the covers may be set aside, if any injury to the health is apprehended; but of this there is not the slightest danger.

In securing the insects in the collection there is one indispensable rule: nothing should touch the bottom. The labels of genera and species, as well as those on the pin, with each specimen, should be as high as possible; the first being nearly on a level in the specimen.


The labels are best made of stout paper, as they are then better kept in position on pins of moderate thickness.

Fig. 7.




172

INSTRUCTIONS FOR COLLECTING INSECTS.



The specimens should be placed at such a distance from each other that they may conveniently, and without risk, be removed and replaced—an operation which is performed with a steel pincers, having the shape indicated in the figure, (fig. 8.) The handle should be large enough to allow a firm grasp of the hand, and the branches should be kept constantly a little separated by means of a spring, as represented. The jaws of the pincers should be roughened like a file on the inside at the ends, and they may be either straight, or, still better, slightly curved, (fig. 9,) as in the figure, to admit of being passed more readily beneath the insect.

The cases used on excursions, or entomological trips, are differently constructed from those of the cabinet. The bottom must be lined with a material soft enough to allow every pin, even the finest, to be inserted easily without the aid of pincers. The best material for this end is pasteboard, which is cut in narrow strips glued against each other, so as to be in a position perpendicular to the bottom of the box.



The bottom of the box is lined with these strips, so that the pins move between the layers of the pasteboard. To prevent the pins so loosely inserted from dropping out, the lining should be sufficiently thick, and, at the same time, the box low enough to allow the head of the loosening pin to lean against the cover before dropping. (See fig. 10.)

Note by R. Ostensacken.—Collectors of Diptera should always endeavor to obtain both sexes of every species. The sexes in the families of the *Tipulidae* and of the *Asilidae* (for the greater part) are easily distinguished by the form of the abdomen, which is generally club-shaped or obtuse

in the male, and more or less pointed in the female. In other families (*Tabani*, *Syrphi*, *Stratiomyidae*, *Bombylii*, and in most of the true *Muscidae*) the eyes are close together in the male, and separated by the "front" in the female. Sometimes they are separated in both sexes, but then the front of the male is generally narrower than that of the female. If these characters fail to help in distinguishing the sexes, the size, and especially the shape, of the abdomen will be sufficient. The males are, for the most part, smaller than the females, and have the abdomen more slender.

It often happens that one of the sexes appears in great abundance, while not a single specimen of the other can be seen. In such cases

INSTRUCTIONS FOR COLLECTING INSECTS.

173

the following observations of Dr. Schiener (Wiener Entomol. Monatschrift, 1858, p. 175) may prove useful:

“From manifold and repeated experience I have arrived at the conclusion that whenever one sex of a Dipterous insect appears in great numbers, the other is always to be found somewhere in the immediate neighborhood.

“If the males soar in the air we can generally predict with certainty that the females are not far off quietly reposing on trunks of trees, as is the case with the *Anthomyzidae*, or on the under side of leaves, as with the *Syrphidae*.

“If every indication of the resting place fails, I then commence catching everything living and flying in the immediate vicinity, sweeping in the air and shaking or beating the leaves and grass, and in this way I have generally succeeded in obtaining, among one hundred specimens of the one sex, at least one of the other.

“The males soaring on the tops of mountains are generally isolated, but on descending several hundred feet we will seldom look in vain for the other sex resting on leaves or the heads of umbelliferous plants.

“I have even succeeded, by patient perseverance, in discovering single males among the numerous female horse-flies (*Tabani*) which attack cattle.”

Those who wish more detailed information about the localities where rare species of Diptera have to be looked for will find them in Bremi's excellent paper on this subject, (*Isis*, 1846, p. 163,) as well as in an article of Mr. Hoffmeister, in the “*Entomologische Zeitung*” of Stettin, 1844, p. 350.

Lloyd Vernon Knutson
4 July 1934 – 10 January 2018



William L. Murphy

Fishers, Indiana, USA; murph3000@gmail.com

Lloyd Vernon Knutson passed away on 10 January at age 83 in Latina, Italy, about 25 miles from his home in Gaeta. The cause of death was heart failure following a stroke he had suffered at home three weeks earlier.

Born in Ottawa, Illinois, Lloyd earned a B.A. in biology at Macalester College, St. Paul, Minnesota, and an M.S. in biology and a Ph.D. in limnology at Cornell University, Ithaca, New York. His Ph.D. dissertation was on the biology of European snail-killing flies (Diptera: Sciomyzidae), the group that would dominate his research for the next 55 years. He was employed for most of his career by the U.S. Department of Agriculture's (USDA) Agricultural Research Service (ARS). During his first five years, he worked as a research entomologist at the U.S. Museum of Natural History, Smithsonian Institution, eventually becoming Chief of the Systematic Entomology Laboratory. He subsequently served for almost 15 years as Chairman of the Insect Identification and Beneficial Insect Introduction Institute at the Beltsville (Maryland) Agricultural Research Center.

In 1988, Lloyd was appointed Director of the USDA/ARS Biological Control of Weeds Laboratory in Rome, Italy. Three years later, he was appointed Director of the USDA/ARS European Biological Control Laboratory in Montpellier, France, a position he held until he retired from federal service in 1997. Within a few years, he and his wife, Mara, had moved to their final home, in coastal Gaeta, Italy, about 65 miles north of Naples. There Lloyd continued to devote an enormous amount of time and energy to continuing his research, primarily on snail-killing flies but also on other organisms and on biological measurements of environmental quality. During his lifetime he authored, co-authored, edited, or co-edited at least 232 publications, with another seven manuscripts currently being completed by various coauthors. He mentored many young entomologists in the United States and in countries around the world, generously sharing his time and resources. His massive knowledge of and contributions to the knowledge of snail-killing flies likely will never be equaled.



Lloyd, 1985 (top), 2012 (bottom)

Many of you will remember Lloyd because of his service to the entomological community. He was president of the Association of Registered Professional Entomologists (1985–1986), of the Eastern Branch of the Entomological Society of America (1978–1979), and of the Entomological Society of America (1988–1989). He was vice-president of the International Organization of Biological Control (1999–2000), secretary of the Association of Systematics Collections (1982–1983), and treasurer of the Society of Systematic Zoology (1971–1974). His extensive knowledge of a broad range of disciplines was a valuable asset in his service as editor for the Biological Society of Washington (1974–1978) and for the Entomological Society of America (1972–1976). In recognition of his vast contributions to entomology, in 1995 Lloyd was elected an honorary member of the Entomological Society of America and in 1996 of the Russian Entomological Society after a visit to St. Petersburg (then Leningrad).

Gifts in memory of Lloyd may be directed to the Cornell University Entomology Excellence Fund (075113) online at giving.cornell.edu/give, by calling (800) 279-3099, or by mail to Cornell University, P.O. Box 25842, Lehigh Valley, Pennsylvania 18003-9692. Please mention the Entomology Excellence Fund (075113) in all correspondence.

A detailed obituary will be published in *Proceedings of the Entomological Society of Washington*

**Bibliography of Papers on Crane Flies (Tipulidae s.l.)
and other Diptera authored by George W. Byers**

Jon Gelhaus

The Academy of Natural Sciences of Drexel University, 1900 Ben Franklin Parkway,
Philadelphia, PA, USA 19103-1195; jkg78@drexel.edu, gelhaus@gmail.com

This bibliography of the Diptera-focused papers (primarily crane flies) written by George W. Byers is a companion to the biography and memories of George which was published earlier this year (Gelhaus et al. in Fly Times April 2018). It does not include the extensive number of papers he authored on the Mecoptera, unless this intersected with Diptera (e.g. Byers, 1989 in list following). This list was compiled from a search of literature at Oosterbroek, P. 2018. Catalogue of the Crane flies of the World (Diptera, Tipuloidea: Pediciidae, Limoniidae, Cylindrotomidae, Tipulidae), <http://ccw.naturalis.nl/index.php>. Many of these papers are available as pdf downloads from this catalogue site. In addition a Web of Science search was carried out for 1980-2018 using “Byers GW” and on ProQuest using “Byers G W” and “Byers George W” resulting in finding additional non-crane fly Diptera papers he authored.

- Rogers, J.S.; Byers, G.W. 1956. The ecological distribution, life history, and immature stages of *Lipsothrix sylvia*. Occasional Papers of the Museum of Zoology, University of Michigan 572: 1-14.
- Byers, G.W. 1958. Species recognition in immature crane flies (Diptera: Tipulidae). Proceedings of the Tenth International Congress of Entomology 1: 131-136.
- Byers, G.W. 1961a. The crane fly genus *Dolichopeza* in North America. The University of Kansas Science Bulletin 42: 665-924.
- Byers, G.W. 1961b. Biology and classification of *Chionea* (Diptera: Tipulidae). Proceedings of the Eleventh International Congress of Entomology 1: 188-191.
- Byers, G.W.; Blank, F.; Hanson, W.J.; Beneway, D.F.; Frederichson, R.W. 1962. Catalogue of the types in the Snow Entomological museum. Part III (Diptera). The University of Kansas Science Bulletin 43: 131-181.
- Byers, G.W. 1963. Type specimens of North American Tipulidae (Diptera) described by Francis Walker. Journal of the Kansas Entomological Society 36: 146-161.
- Byers, G.W. 1966. A crane fly new to the known fauna of Hawaii (Diptera: Tipulidae). Journal of the Kansas Entomological Society 39: 708-711.
- Byers, G.W. 1968. Two new species of Nearctic *Nephrotoma* (Diptera: Tipulidae). Journal of the Kansas Entomological Society 41: 395-403.
- Byers, G. W. 1969. A new family of nematocerous Diptera. J. Kansas Entomological Society 42 (4): 366-371.
- Byers, G.W. 1969. Evolution of wing reduction in crane flies (Diptera: Tipulidae). Evolution 23: 346-354.
- Byers, G. W. 1971. Type specimens of *Trichocera bimacula* and *T. gracilis* (Trichoceridae) and of *Ptychoptera metallica* (Ptychopteridae) described by Francis Walker. Ent. Scand. 7.1976.78-80.
- Byers, G.W. 1974. Larva and pupa of *Idiognophomyia enniki* Alexander (Diptera: Tipulidae). Pan-Pacific Entomologist 50: 282-287.

- Byers, G.W. 1976. Type specimens of Tipulidae described by R.W. Doane in the collection of the California Academy of Sciences. Occasional Papers of the California Academy of Sciences 124: 1-51.
- Byers, G.W. 1977. Tipulidae. In: Hurlbert, S.H. (ed.), Biota Aquatica de Sudamerica Austral, San Diego State Univ., San Diego: 259-265.
- Byers, G.W. 1978. Tipulidae. In: Merritt, R.W.; Cummins, K.W. (eds), An introduction to the aquatic insects of North America. Kendall and Hunt, Dubuque: 285-310.
- Byers, G.W. 1979a. Summer crane-flies of Lake Itasca vicinity, Minnesota. The University of Kansas Science Bulletin 51: 603-613.
- Byers, G.W. 1979b. *Tipula ferruginea* Fabricius, 1805 (Diptera; Tipulidae): Proposed conservation. Z.N.(S.) 2255. Bulletin of Zoological Nomenclature 36: 40-41.
- Byers, G.W. 1979c. *Chionea carolus*, a new species of snow crane fly from California (Diptera, Tipulidae). Journal of the Kansas Entomological Society 52: 719-722.
- Alexander, C.P.; Byers, G.W. 1981. Tipulidae. In: McAlpine, J.F. et al. (eds), Manual of Nearctic Diptera. Vol. 1. Biosystematics Research Institute, Ottawa, Ontario, Monograph 27: 153-190.
- Byers, G.W. 1981a. The crane flies (Diptera: Tipulidae) of Cocos Island, Costa Rica, with descriptions of four new species. Contributions to Science 335: 1-8.
- Byers, G.W. 1981b. Diptera. Tipulidae. In: Hurlbert, S.H.; Rodriguez, G.; Dias dos Santos, N. (eds), Aquatic Biota of Tropical South America: 231-241.
- Byers, G.W. 1982a. In memoriam Charles P. Alexander 1889-1981. Journal of the Kansas Entomological Society 55: 409-417.
- Byers, G.W. 1982b. A subapterous crane fly from Hawaii (Diptera, Tipulidae). Memoirs of the Entomological Society of Washington 10: 37-41.
- Byers, G.W. 1982c. Tipulidae. In: Hurlbert, S.H.; Villalobos-Figueroa, A. (eds), Aquatic Biota of Mexico, Central America and the West Indies: 407-414.
- Byers, G.W. 1983a. The crane fly genus *Chionea* in North America. The University of Kansas Science Bulletin 52: 59-195.
- Byers, G.W. 1983b. Tipulidae and Trichoceridae. In: Morse, J.C. et al. (eds), Aquatic insects of Upper Three Runs Creek, Savannah River Plant, South Carolina. Part II: Diptera. Journal of the Georgia Entomological Society 18: 311-313.
- Gagne, R. J. and G. W. Byers. 1984. A Remarkable New Neotropical Species of *Contarinia* (Diptera: Cecidomyiidae). Journal of the Kansas Entomological Society 57 (4): 736-738.
- Byers, G.W. 1985. A second flightless crane fly from Hawaii (Diptera: Tipulidae). International Journal of Entomology 27: 239-245.
- Byers, G.W. 1987. Synonymy of *Tipula stylifera* Dietz (Diptera: Tipulidae). Journal of the Kansas Entomological Society 60: 334-335.
- Byers, G.W. 1989. Homologies in the wing venation of primitive Diptera and Mecoptera. Proceedings of the Entomological Society of Washington 91: 497-501.
- Arnaud, P.H., Jr.; Byers, G.W. 1990. *Holorusia hespera*, a new name for *Holorusia grandis* (Bergroth) (= *Holorusia rubiginosa* Loew) (Diptera: Tipulidae). Myia 5: 1-9.
- Byers, G.W. 1991. Crane flies of the Kansas ecological reserves. Kansas Academy of Sciences Multidisciplinary Guidebook 4: 91-94.
- Byers, G.W. 1992. Crane flies - three families or one? Acta Zoologica Cracoviensia 35: 37-41.
- Byers, G.W. 1994. A third subapterous crane fly from Hawaii (Diptera: Tipulidae). Proceedings of the Entomological Society of Washington 96: 496-499.
- Gelhaus, J.K.; Byers, G.W. 1994. Tipulidae. In: Morse, J.C. et al. (eds), Aquatic insects of China useful for monitoring water quality. Hohai Univ. Press, Nanjing: 438-458.
- Byers, G.W. 1995. New species of Nearctic snow crane flies of the genus *Chionea* (Diptera: Tipulidae). Proceedings of the Entomological Society of Washington 97: 508-514.

- Byers, G. 1996. More on the Origin of Siphonaptera. *Journal of the Kansas Entomological Society*, 69 (3): 274-277.
- Byers, G.W. 1996. Tipulidae. In: Merritt, R.W.; Cummins, K.W. (eds), *An introduction to the aquatic insects of North America*. 3rd Ed. Kendall and Hunt, Dubuque: 549-570.
- Lantsov, V.I.; Byers, G.W. 1997. In memory of Prof. Evgeniy Nikolaevich Savchenko (1909-1994). *International Journal of Dipterology* 8: 55-67.
- Byers, G.W.; Robison, H.W. 1998. Crane flies of the Ouachita Highlands of Arkansas (Diptera: Tipulidae). *Transactions of the American Entomological Society* 123: 297-302.
- Byers, G.W.; Arnaud P.H.Jr. 2001. A new *Tipula* from Central California (Diptera: Tipulidae). *Myia* 6: 9-22.
- Byers, G.W. 2002. Summer Crane Flies (Tipulidae) of the Mountain Lake Vicinity, Virginia. *Banisteria* 20: 3-30.
- Byers, G.W. 2003. A new species of *Tipula* (*Lunatipula*) from Louisiana (Diptera: Tipulidae). *Journal of the Kansas Entomological Society* 76: 630-633.
- Byers, G.W.; Rossman, D.A. 2003. A new *Dactylolabis* Osten Sacken (Diptera: Tipulidae) from North Carolina. *Proceedings of the Entomological Society of Washington* 105: 70-72.
- Byers, G.W.; Rossman, D.A. 2004. Preliminary survey of the crane flies of Louisiana (Diptera: Tipulidae, Ptychopteridae). *Proceedings of the Entomological Society of Washington* 106: 884-890.
- Byers, G.W.; Rossman, D.A. 2008. A new species of *Dicranomyia* (*Idiopyga*) from Wisconsin, U.S.A. (Diptera: Tipulida). *Journal of the Kansas Entomological Society* 81: 12-14.
- Byers, G.W.; Gelhaus, J.K. 2008. Tipulidae. In: Merritt, R.W.; Cummins, K.W.; Berg, M.B. (eds), *An introduction to the aquatic insects of North America*. 4th Ed. Kendall/Hunt, Dubuque: 773-800.
- Byers, G.W.; Robison, H.W.; Crump, B. 2008. Crane flies of the Ouachita Highlands of Arkansas (Diptera: Tipulidae). II. *Journal of the Kansas Entomological Society* 81: 373-376.
- Byers, G.W. 2009. A rare *Tipula* rediscovered. In: Lantsov, V. (ed.), *Crane flies. History, taxonomy and ecology* (Diptera: Tipulidae, Limoniidae, Pediciidae, Trichoceridae, Ptychopteridae, Tanyderidae). Memorial volume dedicated to Dr. Charles Paul Alexander (1889-1981), Dr. Bernhard Mannheims (1909-1971) and Dr. Evgeniy Nikolaevich Savchenko (1909-1994). *Zoosymposia* 3: 73-75.
- Byers, G.W. 2011. A Miocene *Tipula* from Nevada (Diptera: Tipulidae). *Journal of the Kansas Entomological Society* 84: 169-173.
- Byers, G.W.; Arnaud P. H. Jr. 2011. *Tipula oleracea* Linnaeus in West-Central California (Diptera: Tipulidae). *Journal of the Kansas Entomological Society* 84: 153-154.

MEETING NEWS



The final countdown to ICD9!

Ashley Kirk-Spriggs, ICD9 chair

National Museum, 36 Aliwal Street
Bloemfontein, South Africa; Icd9@nasmus.co.za

We have now begun the final countdown to the opening of the 9th International Congress of Dipterology!

Our Organising Committee has done its level best to make this Congress an international scientific highlight for colleagues from around the world. As well as four excellent plenary speakers, the Congress will cover all major aspects of dipterology, including systematics and taxonomy, morphology, evolution, biodiversity, phylogenetics and conservation, ecology, disease vectors, forensic entomology, phytosanitary Diptera, *etc.* We are especially pleased to note that many of these have special relevance to Diptera researchers based in Africa, specifically symposia on the themes of disease vectors, Diptera pollinators, economically important fruit flies and forensic dipterology. The overall theme of the Congress is “Afrotropical dipterology”, but all themes and regions will be represented.



A total of 273 full delegates and students from 51 countries, are attending the Congress, including 61 delegates from 13 African countries. This is the largest number of African delegates ever to attend ICD.

The scientific Programme is now finalised, which includes four plenary talks, 23 symposia, 258 oral presentations and 58 poster presentations. Due to the large number of oral presentations in the Congress, we will run four parallel symposia sessions on most days and the Organising Committee has done its best to minimise conflicts among them. The Abstracts Volume is also completed, which includes 320 abstracts of oral and poster presentations submitted to the Organising Committee of

ICD9. A printed versions of the book will also be included in delegates' packages and freely downloadable PDF versions of both are available on the ICD9 website.

ICD9 is dedicated to the memory of the late Roger Ward Crosskey (1930–2017), former Honorary Member of ICD, Editor of the *Catalogue of the Diptera of the Afrotropical Region* and author of numerous papers and books on the Simuliidae and Tachinidae, in recognition of his outstanding contributions to the study of dipterology in Africa and the world.

Some special features of the Congress include launch of the 2019 “Year of the Fly”, the official launch of the *Manual of Afrotropical Diptera* and large symposia on “Tephritoidea – applied research and taxonomy”, “Calyptrate evolution and diversity” and “Advances in Diptera phylogenomics”. The Congress also highlights recent developments in “The importance of Diptera in plant-pollinator networks” and includes some novel specialised symposia, including “Diptera interactions with amphibians”. A number of contentious issues will also be raised in the symposium “Too many species, too little time: fresh approaches to “open-ended” genera”, which should lead to some lively debate.

Social events at the Congress comprise a Welcome reception and a separate evening reception for the official launch of the *Manual of Afrotropical Diptera* and happy hour sessions at the Hotel Safari and Safari Court bars. The Congress banquet will take place at the Safari Conference Centre, with a light after dinner speech by Dr Martin Hauser, entitled “Africa and me! – a dipterist’s perspective”.

ICD9 has organised a “Fly Photography Competition” with 14 entries and will award prizes and certificates for the “Best student presentation” and “Best student poster”, with excellent academic books on Diptera as prizes. As part of our public outreach programme two public lectures, by Dr Mark Benecke and Prof. Stephen A. Marshall, will be held in collaboration with the Namibia Scientific Society.

We have received active assistance from our partnering institutions: the National Museum, Bloemfontein (South Africa), the Ministry of Environment and Tourism, Gobabeb Research and training Centre and the Namibia Scientific Society (all Namibia). We have also received generous direct financial support from the National Museum, Bloemfontein, the JRS Biodiversity Foundation and the KwaZulu-Natal Museum. Delegates were supported to attend ICD9 through grants provided by the S.W. Williston Diptera Research Fund and the Belgian Development Cooperation (*via* the Royal Museum for Central Africa, Belgium), as well as other sponsorship as indicated on the back covers of the Abstracts Volume and Programme and on our website.

We look forward to welcoming delegates to Windhoek and hosting a great Congress.

**Organized Meeting: North American Dipterists Society (NADS),
2018 Joint Annual Meeting of the Entomological Societies of
America / Canada / British Columbia**

Matt Bertone

Plant Disease & Insect Clinic, North Carolina State University,
Campus Box 7613, Raleigh, NC 27695-7613; matt_bertone@ncsu.edu

As is tradition, we will be holding an informal organized meeting of the North American Dipterists Society at the 2018 Joint Annual Meeting of the ESA, ESC, and ESBC in Vancouver, British Columbia, Canada. The event will take place on Tuesday evening (November 13) from 7-9 pm, and be held in the Vancouver Convention Centre Meeting Room 215/216.

There will be introductions, announcements, and some business, along with the following presentations (only presenters listed):

- Identifying the freeloaders: Revision of nearctic *Paramyia* (Diptera: Milichiidae) - Julia Mlynarek
- Flies and flytraps: What, if any, are the relationships between Diptera and *Dionaea*? - Clyde E. Sorenson
- New adventures in ecology: Carrion-breeding Diptera associated bacteriophage discovery - Meaghan Pimsler
- Diptera of Palmyra Atoll: New survey, new discoveries - Matthew Bertone

If you are attending the annual meeting please consider joining us. For any questions please email me. Hope to see you there!

North American Dipterists Society Code of Conduct

Matt Bertone

Plant Disease & Insect Clinic, North Carolina State University,
Campus Box 7613, Raleigh, NC 27695-7613; matt_bertone@ncsu.edu

The North American Dipterists Society (NADS) is an informal group that aims to bring together individuals with a common interest in the science and appreciation of true flies. As such, inclusivity, safety and a welcoming environment for all persons regardless of ethnicity, religion, disability, physical appearance, gender, gender identity, or sexual orientation is of utmost importance to us.

The Entomological Society of America (ESA) has tasked us with adopting or creating a code of conduct to formalize our expectations of members' behavior and interactions. We fully support this endeavor. Rather than create our own, we feel the code put forth by the ESA conveys our message and so have adopted it for all meetings and NADS functions. The ESA code of conduct can be found here: <https://www.entsoc.org/conduct>. Although the code describes several types of expectations, we are especially focused on the sections concerning Harassment and Safety, Expected Behavior, and Unaccepted Behavior.

As far as inclusivity, we accept anyone with an interest in Diptera, and anyone who wants to make a presentation focused on flies in our informal meeting at the Annual Meeting of the ESA can do so; scheduled presenters are solicited openly and there is no requirement except that their talk is related to the study or appreciation of Diptera. Unfortunately, because we are an informal society we have no official means or routes for recruiting diverse members, other than word of mouth from existing members and Diptera enthusiasts. We hope current members consider encouraging people with a diversity of backgrounds and identities to become part of our society. Anyone who would like to be included need only fill out the form at the back of the *Fly Times*, or provide the information as outlined on the form.

**2019 Field Meeting of North American Dipterists Society
3-7 June 2019, Bull Shoals Field Station, Missouri**

Greg Courtney¹ & David Bowles²

¹ Department of Entomology, Iowa State University,
401 Science II, Ames, Iowa 50011, USA; gwcourt@iastate.edu

² National Park Service, Heartland Inventory & Monitoring Network
c/o Department of Biology, Missouri State University
901 South National Avenue, Springfield, MO 65897; david_bowles@nps.gov

Invitation to Attend

On behalf of the NADS, we extend a warm invitation for you to attend the 2019 field meeting. This year's meeting, at Bull Shoals Field Station in Missouri, will bring together Dipterists from throughout North America to discuss and collect flies, focusing on taxa from the Ozark Plateau. The meeting will provide a unique environment for interaction in field, laboratory, and informal settings.

Weather during the meeting is likely to be hot (85-90°F). Biting insects, ticks and chiggers, and poison ivy are abundant at the field station so use of personal protective measures is highly recommended.

Specific questions about BSFS or other accommodations can be directed to Dr. David Bowles at the email above, or at 417-836-4702.

Bull Shoals Field Station

Bull Shoals Field Station (BSFS), administered by the Department of Biology, Missouri State University, is located in Taney County, Missouri, close to the Arkansas border. The station is just south of Kirbyville, Missouri (5 miles) and approximately 10 miles from Branson, Missouri. BSFS shares property with the Drury-Mincy Conservation Area managed by the Missouri Department of Conservation. The station occupies some 5,600 acres and is located in the heart of the Ozarks ecoregion.

BSFS has modern, air conditioned, accommodations located in two residential buildings. Residences are outfitted with bunk beds, full kitchens and bathrooms. There is an education classroom with microscopes and table space that can be used for presentations and other activities, as well as a wet lab and large outdoor pavilion. Primitive on-site camping is also permitted. In addition, motor boats can be rented from a local marina, and canoes are available on-site at no cost.



For those not wishing to stay at BSFS, the city of Branson, Missouri has many reasonably priced hotels and restaurants. The following website lists available hotels in the Branson area:

<https://members.bransonchamber.com/list/ql/lodging-76>.

Please see the BSFS website for additional information:

<https://bullshoals.missouristate.edu/>.

Bull Shoals Field Station and surrounding areas offer a wide variety of ecosystems, which serve as habitat to a diverse assemblage of Diptera. The station has Ozark upland forests (oak-hickory dominant), glades, grasslands, old fields, managed food plots, as well as aquatic habitats. Aquatic habitats include naturally occurring woodland ponds, man-made ponds, streams, springs, and a large reservoir (Bull Shoals). Nearby areas include the Buffalo National River and Ozark – St. Francis National Forest to the south (Arkansas) and Mark Twain National Forest to the north and east (Missouri). These areas expand collecting option even more, to include fens and additional large springs and rivers.

The Ozark Plateau is known for its unusual plant and animal communities, including many endemic and disjunct species. The area's diverse habitats will provide many interesting flies! Note that several malaise traps will be set at BSFS a week before the meeting, so we should have plenty of fresh material to examine and identify at the start of the meeting.

Logistics

Participants who fly are advised to fly to Springfield-Branson National Airport (SGF), then arrange for a rental car. SGF is on the northwest side of Springfield and approx. 65 miles from BSFS (1-1.5 hours). There is also a small airport just south of Branson, but it serves only Frontier Airlines.



Tentative schedule**Monday, 3 June (late afternoon / evening)**

Check-in at BSFS
 Collecting near BSFS
 Dinner on your own
 Welcome and introduction to area

Tuesday, 4 June

Breakfast on your own
 Field excursions
 Dinner on your own
 Presentations and sorting the day's catch

Wednesday, 5 June

Breakfast on your own
 Field excursions
 Dinner on your own
 Presentations and sorting the day's catch

Thursday, 6 June

Breakfast on your own
 Field excursions
 Dinner (barbeque) @ BSFS
 Presentations and sorting the day's catch

Friday, 7 June

Breakfast on your own
 Check-out

Cost

There will be a small registration fee (\approx \$30) to cover the cost of using the BSFS classroom & laboratory, a Thursday night barbeque at the Drury house, and incidentals. Except for the group dinner and evening refreshments, food costs will be the responsibility of the registrant. Accommodation costs include the following options.

- 1) \$10/day/person for participants staying in either of the BSFS houses. Note again that this includes shared rooms and bunk beds, with a total of 24 beds. To make reservations, please contact Dr. David Bowles.
- 2) \$5/day/person to camp at either BSFS house or on other BSFS property. This will include access to showers and kitchens inside the house. Again, these should be arranged through Dr. David Bowles.
- 3) Non-BSFS accommodations include numerous motels in Branson.

**Eagle Hill Institute in Maine, USA will be holding a Workshop
on Chironomidae Taxonomy, June 9-15, 2019**

Armin Namayandeh

TAXANAMA Corp.
Toronto, Ontario, Canada; a.namayan@taxanama.com

**Natural History Science Programs on the Eastern Maine Coast – Seminar and Workshop:
Chironomids: Classification, Morphology, Identification, and Lifecycles**

Chironomidae midges are diverse and abundant family of freshwater insects with worldwide distribution and occurrence in variety of habitats. These attributes have proved them to be a good utility tool for environmental biomonitoring and ecological investigation of freshwaters. In most part the taxonomic and biogeographical investigation of this family emerged as a discipline to improve upon this utility. However, difficulty in identification of these taxa and lack of proper association between life stages has prevented their proper use in freshwater investigations. In this seminar I will address some of the issues involved with identification of



chironomid. We will learn the morphology of different life stages and their use in taxonomy. We will explore both macroscopic and microscopic morphological characters to sort and identify Chironomidae to different level and using different life stages. Additionally, we will learn methods such as rearing, collecting, preserving and mounting of immatures and adults.



Please follow the link below for further information:

<https://www.eaglehill.us>

About the instructor

Armin is an aquatic entomologist, focusing mainly on taxonomy and ecology of family Chironomidae (Diptera: Insecta). Armin has worked on various graduate and post graduate freshwater ecology research for over 15 years. His focus for these has been to developed ways to integrate the knowledge of taxonomy and phylogeny into the ecological investigations. Armin main research interest lies on Chironomidae of Northern Canada, especially the subpolar and polar regions. He has conducted taxonomic research on Chironomidae of Eastern Arctic, Chironomidae of Athabasca River and its tributaries and Chironomidae of Precambrian Shield in Canada. Currently, he is conducting research on taxonomy of Chironomidae in far northern Ontario and Chironomidae from Fosheim Peninsula in Ellesmere Island, Nunavut, Canada.




Fly School
23 June–6 July 2019

Brian V. Brown

Natural History Museum of Los Angeles County, 900 Exposition Boulevard,
 Los Angeles, California, 90007, USA; bbrown@nhm.org

The second offering of our fantastic Diptera course is now posted at <http://dipteracourse.com/>. Please tell your students, friends and colleagues to register early. There is a great set of instructors, special guests, and a fabulous location on the Central Coast of California. It will take place 23 June–6 July 2019. The collecting is superb at this time of the year, and we will have access to a tremendous variety of habitats from ocean shore to mountains. Please join us for two weeks of fly immersion: morphology, identification, field techniques, and more!



FLY SCHOOL

A course for Dipterists

OPPORTUNITIES

S.W. Williston Diptera Research Fund

Torsten Dikow

Department of Entomology, National Museum of Natural History, Smithsonian Institution
PO Box 37012, MRC 169, Washington, DC 20013-7012, USA; DikowT@si.edu

The S.W. Williston Diptera Research Fund is a small Smithsonian Institution administered endowment fund established for the *increase and diffusion of knowledge about Diptera*. To this day, the fund has supported the travel of graduate students and dipterists to the International Congresses of Dipterology and to the USNM for collections-based research, field work, and attendance at FlySchool.



The deadline for the 2018 award round is December 15, 2018.

E-mail the following information to Torsten Dikow as a representative of the Williston Fund committee:

1. summarize your research goals into a short proposal in PDF format (1–2 pages maximum)
2. itemize your budget (anticipated transportation costs, per diem costs for lodging and food, and any other items)
3. attach a current CV

Please consider donating to this endowment fund to support the *increase and diffusion of knowledge about Diptera* and particularly the research and travel of a new generation of dipterists.

The Williston Fund is administered by a committee of at least three members, two of whom (the majority) must be systematists actively working on Diptera, and one who must be a scientist affiliated with, but not necessarily employed by, the Smithsonian Institution (for example, a dipterist of the U.S. Department of Agriculture Systematic Entomology Laboratory (SEL)). The current committee consists of: Allen Norrbom, Woogie Kim, and Torsten Dikow.

DIPTERA ARE AMAZING!

The two syrphid photographs were submitted by Andrew Young, who recently finished his PhD with Jeff Skevington at Carleton University, and is currently a postdoc in the lab of the Fly Times editor.



Episyrphus viridaureus (Wiedemann).



Pelecocera (Chamaesyrphus) scaevoides (Fallén).

BOOKS AND PUBLICATIONS

Once again, we offer below a rundown of recent publications on Diptera. As usual if we have not included a paper that you think should have been here please feel free to pass it along to Chris (chris.borkent@gmail.com) and we will include it in the next issue. Unfortunately the online resources do not always catch everything and are a couple of months behind. We also apologize for the missing diacritics in some author's names, unfortunately this is a product of searching in Zoological Record and Web of Science, where they are removed.

- Adler P.H. 2018. Roger Ward Crosskey-The Life and Contributions of an Entomologist par Excellence (1930-2017). *Zootaxa* **4455(1)**: 35-67.
- Arillo, A., Blagoderov, V. and Penalver, E. 2018. Early Cretaceous parasitism in amber: A new species of *Burmazelmira* fly (Diptera: Archizelmiridae) parasitized by a *Leptus* sp mite (Acari, Erythraeidae). *Cretaceous Research* **86**: 24-32.
- Barraclough, D.A., Colville, J.F., Karolyi, F. and Krenn, H.W. 2018. A striking new species of *Prosoeca* Schiner, 1867 (Diptera: Nemestrinidae): An important pollinator from the Bokkeveld Plateau, Northern Cape Province, South Africa. *Zootaxa* **4497(3)**: 411-421. doi:10.11646/zootaxa.4497.3.5.
- Barriault, S., A.O. Soares, S.D. Gaimari & E. Lucas. 2018. *Leucopis glyphinivora* Tanasijtshuk (Diptera: Chamaemyiidae), a new aphidophagous biocontrol agent; Development, survival and comparison with *Aphidoletes aphidimyza* Rondani (Diptera: Cecidomyiidae). *Bulletin of Entomological Research*, 1–7, doi:10.1017/S0007485318000767
- Belqat, B., Adler, P.H., Cherairia, M. and Boudghane-Bendious, C.C. 2018. Inventory of the Black Flies (Diptera: Simuliidae) of North Africa. *Zootaxa* **4442(2)**: 201-220. doi:10.11646/zootaxa.4442.2.1.
- Bickel, D.J. and Kerr, D.S. 2018. *Scorpiurus aramoana* (Diptera: Dolichopodidae), a new species from New Zealand. *Zootaxa* **4422(1)**: 78-84. doi:10.11646/zootaxa.4422.1.4.
- Bost, A., Martinson, V.G., Franzenburg, S., Adair, K.L., Albasi, A., Wells, M.T. and Douglas Angela, E. 2018. Functional variation in the gut microbiome of wild *Drosophila* populations. *Molecular Ecology* **27(13)**: 2834-2845.
- Briem, F., Zeisler, C., Guenay, Y., Staudacher, K., Vogt, H. and Traugott, M. 2018. Identifying plant DNA in the sponging-feeding insect pest *Drosophila suzukii*. *Journal of Pest Science* **91(3)**: 985-994. doi:10.1007/s10340-018-0963-3.
- Broadley, A., Kauschke, E. and Mohrig, W. 2018. Black fungus gnats (Diptera: Sciaridae) found in association with cultivated plants and mushrooms in Australia, with notes on cosmopolitan pest species and biosecurity interceptions. *Zootaxa* **4415(2)**: 201-242. doi:10.11646/zootaxa.4415.2.1.
- Brooks, S.E., Cumming, J.M., Limeira-De-Oliveira, F. and Pollet, M. 2018. New species and records of *Cheiromyia* Dyte from Brazil and French Guiana (Diptera: Dolichopodidae). *Zootaxa* **4444(3)**: 267-282. doi:10.11646/zootaxa.4444.3.3.
- Buenaventura, E. and Pape, T. 2018. Phylogeny, evolution and male terminalia functionality of Sarcophaginae (Diptera: Sarcophagidae). *Zoological Journal of the Linnean Society* **183(4)**: 808-906. doi:10.1093/zoolinnean/zlx070.
- Burt Trevor, O. and Mengual, X. 2018. New records of *Melanostoma fumivenosum* Doesburg (Diptera: Syrphidae) with the description of the male. *Australian Entomologist* **45(2)**: 209-216.

- Busso, J.P. and Blanckenhorn Wolf, U. 2018. Disruptive sexual selection on male body size in the polyphenic black scavenger fly *Sepsis thoracica*. *Behavioral Ecology* **29**(3): 769-777.
- Busso, J.P. and Blanckenhorn, W.U. 2018. Viability selection by invertebrate predators in the polyphenic black scavenger fly *Sepsis thoracica*. *Behavioral Ecology* **29**(4): 992-1000. doi:10.1093/beheco/ary039.
- Caetano, R.L., Carrico, C., Freitas, D.M.D. and Pinto, Z.T. 2018. Ultrastructure of sensilla on antennae and maxillary palps in three Mesembrinellidae species. *Revista Chilena De Historia Natural* **91**: 9. doi:10.1186/s40693-018-0077-6.
- Camp, J.V., Bakonyi, T., Soltész, Z., Zechmeister, T. and Nowotny, N. 2018. *Uranotaenia unguiculata* Edwards, 1913 are attracted to sound, feed on amphibians, and are infected with multiple viruses. *Parasites & Vectors* **11**: 10. doi:10.1186/s13071-018-3030-2.
- Chatelain, P., Plant, A., Soulier-Perkins, A. and Daugeron, C. 2018. Diversity increases with elevation: empidine dance flies (Diptera, Empididae) challenge a predominant pattern. *Biotropica* **50**(4): 633-640.
- Chathuranga, W.G.D., Karunaratne, S.H.P.P., Fernando, B.R. and De Silva, W.A.P.P. 2018. Diversity, distribution, abundance, and feeding pattern of tropical ornithophilic mosquitoes. *Journal of Vector Ecology* **43**(1): 158-167.
- Chuncai, Y., Liu, T., Cao, W., Zhao, G. and Liu, W. 2018. A review of the Japanese *Cryptochironomus* Kieffer, 1918 (Diptera, Chironomidae). *Zookeys* **771**: 139-155.
- Chursina, M.A. and Negrobov, O.P. 2018. Phylogenetic signal in wing shape in the subfamily Dolichopodinae (Diptera, Dolichopodidae) *Zoologicheskyy Zhurnal* **97**(6): 688-700. doi:10.7868/s0044513418060041.
- Coleman, J.M., Benowitz, K.M., Jost, A.G. and Matzkin Luciano, M. 2018. Behavioral evolution accompanying host shifts in cactophilic *Drosophila* larvae. *Ecology and Evolution* **8**(14): 6921-6931.
- Craig Douglas, A., Currie, D.C. and Moulton, J.K. 2018. *Nothogreniera* new genus, for two species of Australian '*Paracnephia*' (Diptera: Simuliidae). *Zootaxa* **4444**(4): 351-380.
- Cranston Peter, S. and Tang, H. 2018. *Skusella* Freeman (Diptera: Chironomidae): new species, immature stages from Africa, Asia and Australia, and expanded distributions. *Zootaxa* **4450**(1): 41-65.
- da Silva, F.L. and Ferrington, L.C. 2018. Systematics of the new world genus *Pentaneura* Phillip (Diptera: Chironomidae: Tanypodinae): Historical review, new species and phylogeny. *Zoologischer Anzeiger* **274**: 60-89. doi:10.1016/j.jcz.2017.11.010.
- Daniels, G. 2018. Designation of a neotype for the Australian robber fly *Ommatius dimidiatus* Macquart, a new senior synonym of *Ommatius pilosus* White and *Ommatius levis* White (Diptera: Asilidae: Ommatiinae). *Zootaxa* **4450**(2): 196-202.
- Danitza Penafiel-Vinueza, A. and Violeta, R. 2018. Five new species of *Drosophila guarani* group from the Andes of southern Ecuador (Diptera, Drosophilidae). *Zookeys* **781**: 141-163.
- Dantas, G.P.S. and Hamada, N. 2018. Immature stages of *Laurotanyus travassosi* Oliveira, Messias & Silva-Vasconcelos, 1992 reveal a new synonymy in Tanypodinae (Diptera: Chironomidae). *Zootaxa* **4444**(4): 462-470. doi:10.11646/zootaxa.4444.4.6.
- De Camargo, S., Carvalho, F.D. and Esposito, M.C. 2018. The genus *Peckia* Robineau-Desvoidy (Diptera: Sarcophagidae) in the Brazilian Amazon: a new species, new records, descriptions of female terminalia and key to species. *Zootaxa* **4483**(1): 1-35. doi:10.11646/zootaxa.4483.1.1.
- De Santis, M.D. 2018. A new genus and species of Sophiini Townsend, 1936 (Diptera: Tachinidae) from Brazil, with an updated key to the genera of the tribe. *Zootaxa* **4500**(3): 433-442. doi:10.11646/zootaxa.4500.3.9.

- Deeming, J.C. 2018. A revision of the Afrotropical species of *Pachylophus* Loew (Diptera: Chloropidae) and some related Ovoviviparous genera. *Zootaxa* **4482(1)**: 1-51. doi:10.11646/zootaxa.4482.1.1.
- Dias, N.P., Zotti, M.J., Montoya, P., Carvalho, I.R. and Nava, D.E. 2018. Fruit fly management research: A systematic review of monitoring and control tactics in the world. *Crop Protection* **112**: 187-200. doi:10.1016/j.cropro.2018.05.019.
- Diaz, F., Lima, A.L.A., Nakamura, A.M., Fernandes, F., Sobrinho, I. and de Brito, R.A. 2018. Evidence for introgression among three species of the *Anastrepha fraterculus* group, a radiating species complex of fruit flies. *Frontiers in Genetics* **9**: 17. doi:10.3389/fgene.2018.00359.
- Disney, R.H.L. 2018. The hitherto unknown flightless female of *Puliciphora calix* Carles-Tolra (Diptera: Phoridae) from France. *Entomologist's Monthly Magazine* **154(3)**: 176-178.
- Du, J. and Xue, W.Q. 2018. A review of the *Delia interflua* group with descriptions of two new species (Diptera, Anthomyiidae). *Zookeys* **764**: 145-154. doi:10.3897/zookeys.764.22736.
- Ebejer Martin, J. 2018. Recent additions to the Chyromyidae of Morocco with description of a new species of *Gymnochiromyia* Hendel (Diptera: Acalyptrata). *Entomologist's Monthly Magazine* **154(3)**: 207-212.
- Eiseman, C.S., Heller, K. and Rulik, B. 2018. A new dark-winged fungus gnat (Diptera: Sciaridae) mining leaves of *Clintonia borealis* (Aiton) Raf. (Liliaceae). *Proceedings of the Entomological Society of Washington* **120(3)**: 500-507. doi:10.4289/0013-8797.120.3.500.
- El-Ahmady, A., Taha, M., Soliman, A.M. and El-Hawagry, M. 2018. A new species and new records of the genus *Sarcophaga* from Egypt, with a key to the known Egyptian species (Diptera: Sarcophagidae). *African Entomology* **26(2)**: 507-521. doi:10.4001/003.026.0507.
- El-Hawagry Magdi, S., Zatwarnicki, T. and Ebrahim, A.M. 2018. Catalogue of the Egyptian Ephydroidea (Diptera: Schizophora: Acalyptratae). *Zootaxa* **4444(3)**: 201-246.
- El-Hawagry, M.S. 2018. Catalogue of the Tachinidae of Egypt (Diptera: Oestroidea). *Egyptian Journal of Biological Pest Control* **28**: 19. doi:10.1186/s41938-018-0042-3.
- Elsayed, A.K., Matsuo, K., Kim, W., Uechi, N., Yukawa, J., Gyoutoku, N. and Tokuda, M. 2018. A new *Asphondylia* species (Diptera: Cecidomyiidae) and a eulophid wasp (Hymenoptera) inducing similar galls on leaf buds of *Schoepfia jasminodora* (Schoepfiaceae), with reference to their ecological traits and a description of the new gall midge. *Entomological Science* **21(3)**: 324-339. doi:10.1111/ens.12312.
- Epler, J.H. 2018. The genus *Cryptotendipes* Beck et Beck in Florida, with the description of a new species (Diptera: Chironomidae: Chironominae). *Zootaxa* **4433(3)**: 583-594. doi:10.11646/zootaxa.4433.3.12.
- Estrada-Villegas, S., Halczok, T.K., Tschapka, M., Page, R.A., Braendel, S.D. and Hiller, T. 2018. Bats and their bat flies: community composition and host specificity on a Pacific island archipelago. *Acta Chiropterologica* **20(1)**: 161-176.
- Evenhuis Neal, L., Perreira, W.D. and Yee, D.A. 2018. New records of two-winged flies (Insecta: Diptera) from O'ahu, Hawaiian Islands. *Bishop Museum Occasional Papers* **123**: 51-54.
- Evenhuis, N.L. and Kettani, K. 2018. Genera of Mythicomyiidae (Insecta, Diptera) new to the fauna of Morocco, with descriptions of new species. *Zootaxa* **4429(2)**: 348-356. doi:10.11646/zootaxa.4429.2.9.
- Ferro, G.B. and Marshall, S.A. 2018. A revision of the Neotropical ant-like genus *Cardiacephala* Macquart, including *Plocoscelus* Enderlein *syn. nov.* (Diptera: Micropezidae, Taeniapterinae). *Zootaxa* **4429(3)**: 401-458. doi:10.11646/zootaxa.4429.3.1.
- Fu, Y., Liu, T., Fang, X., Wang, Q. and Wang, X. 2018. Three new species of *Corynoneura* Winnertz from Oriental China (Diptera: Chironomidae: Orthoclaadiinae). *Zootaxa* **4418(1)**: 85-92.

- Fuller, L., Fuentes-Montemayor, E., Watts, K., Macgregor, N.A., Bitenc, K. and Park, K.J. 2018. Local-scale attributes determine the suitability of woodland creation sites for Diptera. *Journal of Applied Ecology* **55**(3): 1173-1184. doi:10.1111/1365-2664.13035.
- Galinskaya, T.V., Yu, O.N., Nartschuk, E.P. and Shatalkin, A.I. 2018. DNA Barcoding of Celyphidae (Diptera) from Vietnam. *Russian Journal of Genetics* **54**(8): 950-959.
- Garcia, C.D. and Urso-Guimaraes, M.V. 2018. Three new species of *Lopesia* Rubsaaen (Diptera: Cecidomyiidae) from Brazil. *Florida Entomologist* **101**(2): 203-211.
- Gardner, E.M., Gagne, R.J., Kendra, P.E., Montgomery, W.S., Raguso, R.A., McNeil, T.T. and Zerega, N.J.C. 2018. A flower in fruit's clothing: Pollination of jackfruit (*Artocarpus heterophyllus*, Moraceae) by a new species of gall midge, *Clinodiplosis ultracrepidata* sp. nov. (Diptera: Cecidomyiidae). *International Journal of Plant Sciences* **179**(5): 350-367. doi:10.1086/697115.
- Ge, Y.Q., Gao, Y.Y. and Zhang, D. 2018. Two new synonymies in the genus *Lispe* (Diptera: Muscidae). *Zootaxa* **4457**(4): 549-556. doi:10.11646/zootaxa.4457.4.4.
- Ge, Y.Q., Zhang, D. and Thomas, P. 2018. A new species of *Wohlfahrtia* Brauer & Bergenstamm (Diptera: Sarcophagidae) from northwestern China, with three new synonymies and a pictorial synopsis. *Zootaxa* **4434**(1): 130-140. doi:10.11646/zootaxa.4434.1.8.
- Gilka, W., Paasivirta, L., Gadawski, P. and Grabowski, M. 2018. Morphology and molecules say: *Tanytarsus latens*, sp nov from Finland (Diptera: Chironomidae). *Zootaxa* **4471**(3): 569-579. doi:10.11646/zootaxa.4471.3.8.
- Gillung, J.P., Winterton, S.L., Bayless, K.M., Khouri, Z., Borowiec, M.L., Yeates, D., Kimsey, L.S., Misof, B., Shing, S., Zhou, X. and others. 2018. Anchored phylogenomics unravels the evolution of spider flies (Diptera, Acroceridae) and reveals discordance between nucleotides and amino acids. *Molecular Phylogenetics and Evolution* **128**: 233-245. doi:10.1016/j.ympev.2018.08.007.
- Giunti, G., Campolo, O., Laudani, F. and Palmeri, V. 2018. Male courtship behaviour and potential for female mate choice in the black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae). *Entomologia Generalis* **38**(1): 29-46. doi:10.1127/entomologia/2018/0657.
- Gomard, Y., Cornuault, J., Licciardi, S., Lagadec, E., Belqat, B., Dsouli, N., Mavingui, P. and Tortosa, P. 2018. Evidence of multiple colonizations as a driver of black fly diversification in an oceanic island. *Plos One* **13**(8): 20. doi:10.1371/journal.pone.0202015.
- Grimaldi, D.A. 2018. First tropical american species of the "relict" genus *Litoleptis*, and relationships in Spaniinae (Diptera: Rhagionidae). *American Museum Novitates*(**3909**): 1-18.
- Grimaldi, D.A. 2018. *Hirtodrosophila* of North America (Diptera: Drosophilidae). *Bulletin of the American Museum of Natural History*(**421**): 1-75.
- Guan-Jun, C., Xu-Yuan, G., Long, X.-Z., Xian-Ru, Z., Ji-Min, L., De-Wei, W. and Yu, Y.-H. 2018. An adaptive bivariate kernel smoothing method for determining instars of *Austrosimulium tillyardianum* (Diptera: Simuliidae) larvae. *Ecological Entomology* **43**(4): 412-421.
- Haelewaters, D., Hiller, T. and Dick, C.W. 2018. Bats, Bat Flies, and Fungi: A Case of Hyperparasitism. *Trends in Parasitology* **34**(9): 784-799. doi:10.1016/j.pt.2018.06.006.
- Hancock, D.L. and Drew, R.A.I. 2018. A review of the subgenera *Asiadacus* Perkins, *Diplodacus* May, *Hemigymnodacus* Hardy, *Niuginidacus* Drew, *Papuodacus* Drew and *Sinodacus* Zia of *Bactrocera* Macquart (Diptera: Tephritidae: Dacinae). *Australian Entomologist* **45**(2): 181-208.
- Harbach, R.E. 2018. Two new species of the subgenus *Sabethinus* of *Sabethes* (Diptera: Culicidae) from Costa Rica, first confirmation of members of the subgenus in Central America. *Zootaxa* **4429**(2): 269-280. doi:10.11646/zootaxa.4429.2.3.

- Havill, N.P., Gaimari, S.D. and Caccone, A. 2018. Cryptic east-west divergence and molecular diagnostics for two species of silver flies (Diptera: Chamaemyiidae: *Leucopis*) from North America being evaluated for biological control of hemlock woolly adelgid. *Biological Control* **121**: 23-29. doi:10.1016/j.biocontrol.2018.02.004.
- Heaton, V., Moffatt, C. and Simmons, T. 2018. The movement of fly (Diptera) larvae within a feeding aggregation. *Canadian Entomologist* **150(3)**: 326-333.
- Hermansen, J.S., Starrfelt, J., Voje, K.L. and Stenseth, N.C. 2018. Macroevolutionary consequences of sexual conflict. *Biology Letters* **14(6)**: 4. doi:10.1098/rsbl.2018.0186.
- Hiller, T., Honner, B., Page, R.A. and Tschapka, M. 2018. Leg structure explains host site preference in bat flies (Diptera: Streblidae) parasitizing neotropical bats (Chiroptera: Phyllostomidae). *Parasitology* **145(11)**: 1475-1482. doi:10.1017/s0031182018000318.
- Hoffmann, C., Merkel, K., Sachse, A., Rodriguez, P., Leendertz, F.H. and Calvignac-Spencer, S. 2018. Blow flies as urban wildlife sensors. *Molecular Ecology Resources* **18(3)**: 502-510.
- Hopkins, M.C., Thomason, C.A., Brown, B.L., Kirkpatrick, L.T., Paulson, S.L. and Hawley, D.M. 2018. Experimental logging alters the abundance and community composition of ovipositing mosquitoes in the southern Appalachians. *Ecological Entomology* **43(4)**: 463-472.
- Hou, P., Li, X., Yang, D. and Zhang, C. 2018. Taxonomic study of *Gonia* Meigen (Diptera: Tachinidae) from China. *Zoological Systematics* **43(3)**: 294-308.
- Huang, J., Wang, Y.L., O'Grady, P.M., Su, Y.R. and Chen, H.W. 2018. The genus *Leucophenga* (Diptera, Drosophilidae), part VIII: twenty-one species from the Oriental region, with morphological and molecular evidence. *Zootaxa* **4503(1)**: 1-70. doi:10.11646/zootaxa.4503.1.1.
- Hussain, R.I., Walcher, R., Brandl, D., Jernej, I., Arnberger, A., Zaller, J.G. and Frank, T. 2018. Influence of abandonment on syrphid assemblages in mountainous meadows. *Journal of Applied Entomology* **142(4)**: 450-456.
- Ishihara, D. 2018. Role of fluid-structure interaction in generating the characteristic tip path of a flapping flexible wing. *Physical Review E* **98(3)**: 19. doi:10.1103/PhysRevE.98.032411.
- Jaleel, W., Lu, L.H. and He, Y.R. 2018. Biology, taxonomy, and IPM strategies of *Bactrocera tau* Walker and complex species (Diptera; Tephritidae) in Asia: a comprehensive review. *Environmental Science and Pollution Research* **25(20)**: 19346-19361. doi:10.1007/s11356-018-2306-6.
- Jaschhof, M. and Jaschhof, C. 2018. Descriptions of eighteen new species of *Glemparon*, a previously monotypic genus of Porricondyliinae (Diptera, Cecidomyiidae). *European Journal of Taxonomy* **450**: 1-38.
- Jaschhof, M. and Spungis, V. 2018. Towards reliable identification of male *Dicerura*: descriptions of three new and seven poorly known species in the Palearctic region (Diptera: Cecidomyiidae, Porricondyliinae). *Zootaxa* **4422(1)**: 85-103. doi:10.11646/zootaxa.4422.1.5.
- Jezek, J., Obona, J., Le Pont, F., Maes, J.M. and Mollinedo, S. 2018. Two new species of Bruchomyiinae (Diptera, Psychodidae) from the Neotropical Region. *Zootaxa* **4442(3)**: 469-478. doi:10.11646/zootaxa.4442.3.8.
- Jiao, K.-L., Wang, H., Jun-Hao, H., Pei-Jin, H., Long-Wa, Z., Jiao, X., Zhi-Hui, W., Zhang, J., Yuan-Hong, W., Wen-Jun, B. and others. 2018. A new species of *Contarinia* (Diptera: Cecidomyiidae) damaging inflorescence of *Carya cathayensis* (Juglandaceae) in China. *Zootaxa* **4442(1)**: 187-193.
- Jimenez-Cortes, J.G., Garcia-Contreras, R., Bucio-Torres, M.I., Cabrera-Bravo, M., Cordoba-Aguilar, A., Benelli, G. and Salazar-Schettino, P.M. 2018. Bacterial symbionts in human blood-feeding arthropods: Patterns, general mechanisms and effects of global ecological changes. *Acta Tropica* **186**: 69-101. doi: 10.1016/j.actatropica.2018.07.005.

- Jose Saravia-Pietro Paolo, M., Arce, S.i., Manzoli, D.E., Quiroga, M. and Beldomenico, P.M. 2018. Aspects of the life cycle of the avian parasite *Philornis torquans* (Diptera: Muscidae) under laboratory rearing conditions. *Canadian Entomologist* **150**(3): 317-325.
- Kato, D., Tachi, T. and Gelhaus, J. 2018. Revision of the subgenus *Dicranomyia* (*Erostrata*) Savchenko, 1976 (Diptera, Limoniidae) of Japan. *Zootaxa* **4441**(1): 181-194. doi:10.11646/zootaxa.4441.1.11.
- Katoh, T.K., Zhang, G., Toda, M.J., Suwito, A. and Gao, J.J. 2018. A revision of the subgenus *Dudaica* Strand of the genus *Drosophila* Fallen, with descriptions of six new species (Diptera, Drosophilidae). *Zookeys* **781**: 19-50. doi:10.3897/zookeys.781.27354.
- Katoh, T.K., Zhang, G., Toda, M.J., Zhang, W.X. and Gao, J.J. 2018. The *Lordiphosa denticeps* species group (Diptera: Drosophilidae) in China, with redescription of four known species and descriptions of nine new species. *Zootaxa* **4471**(1): 37-75. doi:10.11646/zootaxa.4471.1.2.
- Katoh, T.K., Zhang, G., Zhou, C.J. and Gao, J.J. 2018. Taxonomy of the *Hirtodrosophila melanderi* species group (Diptera: Drosophilidae), with descriptions of four new species from southwestern China. *Zootaxa* **4422**(3): 345-365. doi:10.11646/zootaxa.4422.3.2.
- Kaulfuss, U., Lee, D.E., Wartho, J.-A., Bowie, E., Lindqvist, J.K., Conran, J.G., Bannister, J.M., Mildenhall, D.C., Kennedy, E.M. and Gorman, A.R. 2018. Geology and palaeontology of the Hindon Maar Complex: A Miocene terrestrial fossil Lagerstätte in southern New Zealand. *Palaeogeography Palaeoclimatology Palaeoecology* **500**: 52-68.
- Kazerani, F. and Kehlmaier, C. 2018. A new species of the genus *Clistoabdominalis* Skevington (Diptera: Pipunculidae) from Iran, with a key to the Western Palearctic species of the *Clistoabdominalis ruralis* group. *Zootaxa* **4425**(3): 582-588. doi:10.11646/zootaxa.4425.3.11.
- Keresztes, L., Kolcsar, L.P., Denes, A.L. and Torok, E. 2018. Revealing unknown larvae of the *maxima* species group of the genus *Acutipula* Alexander, 1924 (*Tipula*, Tipuloidae, Diptera) using an integrative approach. *North-Western Journal of Zoology* **14**(1): 17-24.
- Kimsey Lynn, S., Kimsey, R.B. and Gaimari, S.D. 2018. Novel biology for *Tephrochlamys rufiventris* (Meigen, 1830) (Diptera: Heleomyzidae). *Proceedings of the Entomological Society of Washington* **120**(3): 543-548.
- Knop, E., Gerpe, C., Ryser, R., Hofmann, F., Menz, M.H.M., Trosch, S., Ursenbacher, S., Zoller, L. and Fontaine, C. 2018. Rush hours in flower visitors over a day-night cycle. *Insect Conservation and Diversity* **11**(3): 267-275. doi:10.1111/icad.12277.
- Knutson, L.V., Deeming, J.C. and Ebejer, M.J. 2018. The Snail-killing Flies (Diptera: Sciomyzidae) of West Africa. *Zootaxa* **4483**(1): 67-100. doi:10.11646/zootaxa.4483.1.3.
- Kodama, A., Kawai, K. and Saito, H. 2018. A new species of *Tanytarsus* van der Wulp, 1874 (Diptera: Chironomidae) and the first Japanese record of *Tanytarsus ovatus* Johannsen, 1932 with DNA barcodes. *Zootaxa* **4422**(4): 591-599. doi:10.11646/zootaxa.4422.4.9.
- Kolesik, P., Baker, G., Hill, K., Manners, A.G. and Dijkstra, E. 2018. New species of gall midge (Diptera: Cecidomyiidae) damaging flower buds of ornamental *Alstroemeria* plants. *Austral Entomology* **57**(3): 285-291.
- Kraemer, M.M.S., Delclos, X., Clapham, M.E., Arillo, A., Peris, D., Jager, P., Stebner, F. and Penalver, E. 2018. Arthropods in modern resins reveal if amber accurately recorded forest arthropod communities. *Proceedings of the National Academy of Sciences of the United States of America* **115**(26): 6739-6744. doi:10.1073/pnas.1802138115.
- Krafsur, E.S. and Maudlin, I. 2018. Tsetse fly evolution, genetics and the trypanosomiases - A review. *Infection Genetics and Evolution* **64**: 185-206. doi:10.1016/j.meegid.2018.05.033.

- Krolow, T.K., Henriques, A.L. and Pollet, M. 2017. The Tabanidae of the Mitaraka expedition, with an updated check list of French Guiana (Diptera). *Zookeys* **684**: 85-118. doi:10.3897/zookeys.684.13197.
- Kurina, O., Hippa, H. and Amorim, D.D. 2018. A contribution to the systematics of the genus *Manota* Williston (Diptera: Mycetophilidae) in Brazil. *Zootaxa* **4472(1)**: 1-59. doi:10.11646/zootaxa.4472.1.1.
- Kutty, S.N., Wong, W.H., Meusemann, K., Meier, R. and Cranston, P.S. 2018. A phylogenomic analysis of Culicomorpha (Diptera) resolves the relationships among the eight constituent families. *Systematic Entomology* **43(3)**: 434-446. doi:10.1111/syen.12285.
- Kvifte, G.M. 2018. Molecular phylogeny of moth flies (Diptera, Psychodidae, Psychodinae) revisited, with a revised tribal classification. *Systematic Entomology* **43(3)**: 596-605. doi:10.1111/syen.12288.
- Kvifte, G.M. and Ivkovic, M. 2018. New species and records of the *Pericoma trifasciata* group from Croatia (Diptera: Psychodidae). *Zootaxa* **4486(1)**: 76-82. doi:10.11646/zootaxa.4486.1.5.
- Li, L., Lan, M., Lu, W., Li, Z., Xia, T., Zhu, J., Ye, M., Gao, X. and Wu, G. 2018. De novo transcriptomic analysis of the alimentary tract of the tephritid gall fly, *Procecidochares utilis*. *Plos One* **13(8)**: 1.
- Li, X. and Yeates, D.K. 2018. Revision of the Australian bee fly genus *Pilosia* Hull, 1973 (Bombyliidae, Bombyliinae) from Western Australia. *Zootaxa* **4438(2)**: 299-312.
- Li, X.K., Rodrigues, P.F.M., Lamas, C.J.E. and Yeates, D.K. 2018. A Review of the Australian Species of *Thevenetimyia* Bigot, 1875 (Bombyliidae, Bombyliinae, Eclimini), with description of four new species and the pupal case of *T. longipalpis* (Hardy). *Records of the Australian Museum* **70(3)**: 331-375. doi:10.3853/j.2201-4349.70.2018.1678.
- Li, X.K. and Yeates, D.K. 2018. One hundred years of solitude ended: A second species of *Marmasoma* White, 1916 (Diptera, Bombyliidae, Bombyliinae, Eclimini) from Australia. *Zootaxa* **4420(1)**: 34-42. doi:10.11646/zootaxa.4420.1.2.
- Liendo Maria, C., Parreno, M.A., Cladera, J.L., Vera, M.T. and Segura, D.F. 2018. Coexistence between two fruit fly species is supported by the different strength of intra- and interspecific competition. *Ecological Entomology* **43(3)**: 294-303.
- Lima, H.I.L., Krolow, T.K. and Henriques, A.L. 2018. A new species of *Dichelacera* (*Dichelacera*) Macquart (Diptera, Tabanidae) from the Brazilian savannah. *Neotropical Entomology* **47(3)**: 380-384.
- Lin, X.L., Stur, E. and Ekrem, T. 2018. Molecular phylogeny and temporal diversification of *Tanytarsus* van der Wulp (Diptera: Chironomidae) support generic synonymies, a new classification and centre of origin. *Systematic Entomology* **43(4)**: 659-677. doi:10.1111/syen.12292.
- Liu, R.S., Bian, Y., Zhang, R.F. and Yang, D. 2018. A review of *Diaphorus* (Diptera: Dolichopodidae) from China with four new species. *Transactions of the American Entomological Society* **144(2)**: 417-429. doi:10.3157/061.144.0208.
- Londt, J., G. H. and Dikow, T. 2018. A review of the assassin-fly genus *Laphyctis* Loew, 1858 with descriptions of two new species (Diptera, Asilidae, Laphriinae). *African Invertebrates* **58(1)**: 75-106.
- Lu, J.M., Zhang, Y. and Chen, H.W. 2018. Integrative taxonomy of the genus *Pseudostegana* (Diptera, Drosophilidae) from China, with descriptions of eleven new species. *PeerJ* **6**: 41. doi:10.7717/peerj.5160.
- MacGowan, I. 2018. Definition of the *Lonchaea polyhamata* species-group (Diptera, Lonchaeidae) with a description of new species. *Zootaxa* **4446(3)**: 325-342. doi:10.11646/zootaxa.4446.3.2.
- MacGowan, I. and Compton, S.G. 2018. A new species of *Silba* (Diptera; Lonchaeidae) associated with figs. *Zootaxa* **4455(1)**: 196-200.

- Maia, V.C., Flor, I.C. and Oliveira, L.A. 2018. *Myrciamyia pterandrae* (Diptera, Cecidomyiidae, Lopesiini), a new species of gall midge associated with *Pterandra pyroidea* A. Juss. (Malpighiaceae), an endemic plant in Brazilian Cerrado. *Revista Brasileira De Entomologia* **62(3)**: 220-224. doi:10.1016/j.rbe.2018.06.003.
- Marcondes, C.B., de Mello, C.F., Bastos, A.Q., Montarsi, F., Alencar, J. and Santos-Mallet, J. 2018. Description of the egg of *Hulecoeteomyia koreica* (Edwards) (Diptera: Culicidae) using scanning electron microscopy. *Zootaxa* **4442(1)**: 194-200. doi:10.11646/zootaxa.4442.1.13.
- McCreadie, J.W., Williams, R.H., Stutsman, S., Finn, D.S. and Adler, P.H. 2018. The influence of habitat heterogeneity and latitude on gamma diversity of the Nearctic Simuliidae, a ubiquitous group of stream-dwelling insects. *Insect Science* **25(4)**: 712-720. doi:10.1111/1744-7917.12442.
- Medeiros, H.R., Hoshino, A.T., Ribeiro, M.C., Morales, M.N., Martello, F., Neto, O.C.P., Carstensen, D.W. and Menezes, A.D. 2018. Non-crop habitats modulate alpha and beta diversity of flower flies (Diptera, Syrphidae) in Brazilian agricultural landscapes. *Biodiversity and Conservation* **27(6)**: 1309-1326. doi:10.1007/s10531-017-1495-5.
- Mederos, J. 2018. New species of *Heteropterna* Skuse and *Keroplatus* Bosc (Diptera: Keroplatidae) become the second records of this family for Cuba. *Zootaxa* **4461(1)**: 57-68. doi:10.11646/zootaxa.4461.1.3.
- Mengual, X., Miranda, G.F.G. and Thompson, F.C. 2018. Unraveling *Ocyptamus* and the *Baccha* legacy (Diptera: Syrphidae): redefinition of groups and new species descriptions. *Zootaxa* **4461(1)**: 1-44. doi:10.11646/zootaxa.4461.1.1.
- Mengual, X., Stahls, G., Laska, P., Mazanek, L. and Rojo, S. 2018. Molecular phylogenetics of the predatory lineage of flower flies *Eupeodes-Scaeva* (Diptera: Syrphidae), with the description of the Neotropical genus *Austroscaeva* gen. nov. *Journal of Zoological Systematics and Evolutionary Research* **56(2)**: 148-169. doi:10.1111/jzs.12212.
- Mlynarek, J.J. and Wheeler, T.A. 2018. Phylogeny and revised classification of the tribe Elachipterini (Diptera: Chloropidae). *Zootaxa* **4471(1)**: 1-36. doi:10.11646/zootaxa.4471.1.1.
- Mohamadzade Namin, S. and Gaimari, S.D. 2018. The genus *Sapromyza* (Diptera: Lauxaniidae) in Iran with description of three new species. *Zootaxa* **4471(1)**: 154-168. doi:10.11646/zootaxa.4471.1.7.
- Mohrig, W., Kauschke, E. and Broadley, A. 2018. New black fungus gnats (Diptera: Sciaridae) from Eastern Australia. *Zootaxa* **4450(2)**: 203-241. doi:10.11646/zootaxa.4450.2.3.
- Mortelmans, J. and Pollet, M. 2018. New data and species of *Thecomyia* Perty, 1833 (Diptera: Sciomyzidae) from Mitaraka (French Guiana), with notes on the genus. In: Touroult J. (Ed.). "Our Planet Reviewed" 2015 large-scale biotic survey in Mitaraka, French Guiana. *Zoosystema* **40 (17)**: 415-423. doi: 10.5252/zoosystema2018v40a17
- Moulton John, K. 2018. Rediscovery of *Dixa repanda* Peters and description of its putative sister species, *Dixa falcata* sp nov., from the south-central United States (Diptera: Dixidae). *Zootaxa* **4459(1)**: 171-178.
- Mugasa, C.M., Villinger, J., Gitau, J., Ndungu, N., Ciosi, M. and Masiga, D. 2018. Morphological re-description and molecular identification of Tabanidae (Diptera) in East Africa. *Zookeys* **769**: 117-144. doi:10.3897/zookeys.769.21144.
- Muller, B.S. and Mostovski, M.B. 2018. A key to males of *Atherigona* s. str. (Diptera: Muscidae) from Mali, with new records and a new species for the country. *Zootaxa* **4425(2)**: 342-356. doi:10.11646/zootaxa.4425.2.9.
- Murphy, W.L., Mathis, W.N. and Knutson, L.V. 2018. Comprehensive taxonomic, faunistic, biological, and geographic inventory and analysis of the Sciomyzidae (Diptera: Acalyptratae) of the Delmarva region and nearby states in eastern North America. *Zootaxa* **4430(1)**: 1-299. doi:10.11646/zootaxa.4430.1.1.

- Muto, L., Kamimura, Y., Tanaka, K.M. and Takahashi, A. 2018. An innovative ovipositor for niche exploitation impacts genital coevolution between sexes in a fruit-damaging *Drosophila*. *Proceedings of the Royal Society B-Biological Sciences* **285(1887)**: 8. doi:10.1098/rspb.2018.1635.
- Natarajan, R., Rajavel, A.R. and Jambulingam, P. 2018. *Uranotaenia pseudostricklandi*, a new species in subgenus *Pseudoficalbia* (Diptera: Culicidae) from Kerala, India. *Zootaxa* **4429(1)**: 181-188. doi:10.11646/zootaxa.4429.1.11.
- Nedeljkovic, Z., Ricarte, A., Zoric, L.S., Dan, M., Vidakovic, D.O. and Vujic, A. 2018. The genus *Xanthogramma* Schiner, 1861 (Diptera: Syrphidae) in southeastern Europe, with descriptions of two new species. *Canadian Entomologist* **150(4)**: 440-464. doi:10.4039/tce.2018.21.
- Nicacio, G. and Juen, L. 2018. Relative roles of environmental and spatial constraints in assemblages of Chironomidae (Diptera) in Amazonian floodplain streams. *Hydrobiologia* **820(1)**: 201-213. doi:10.1007/s10750-018-3657-1.
- Okuyama, Y., Okamoto, T., Kjaerandsen, J. and Kato, M. 2018. Bryophytes facilitate outcrossing of *Mitella* by functioning as larval food for pollinating fungus gnats. *Ecology* **99(8)**: 1890-1893. doi:10.1002/ecy.2364.
- Orel, O.V. 2018. New species of the non-biting midges of Tribe Chironomini (Diptera, Chironomidae) from the Bolshekhkhtsyrsky Reserve (Russia, Khabarovsk Territory). *Zootaxa* **4461(2)**: 269-276. doi:10.11646/zootaxa.4461.2.8.
- Orel, O.V. and Bazova, N.V. 2018. New species of the genus *Microtendipes* Kieffer, 1915 (Diptera, Chironomidae) from Buryatia (Russia). *Zootaxa* **4425(1)**: 175-184. doi:10.11646/zootaxa.4425.1.11.
- Orfinger, A.B., Kelly, S.L. and Woller, D.A. 2018. Flower-feeding by two species of Mydas fly (Diptera: Mydidae) in Florida, with nNotes on tree-form *Licania michauxii* (Gopher Apple). *Southeastern Naturalist* **17(2)**: N19-N27. doi:10.1656/058.017.0208.
- Ospina-Torres, R., Mey, W. and Jaime-Murcia, P. 2018. Two new orthoclad species from Colombian Andes (Diptera: Chironomidae). *Zootaxa* **4472(2)**: 385-392.
- Ozerov, A.L. 2018. Contribution to the fauna of the genus *Meroplius* Rondani, 1874 (Diptera, Sepsidae) of the Australasian/Oceanian region. *Zootaxa* **4438(1)**: 195-200.
- Paggi, A.C. and Zilli, F. 2018. Redescription of two species of *Coelotanypus* (Kieffer) 1913 (Diptera: Chironomidae) in rivers of the Parano-Platense basin (Argentina, South America). *Zootaxa* **4486(1)**: 67-75. doi:10.11646/zootaxa.4486.1.4.
- Parchami-Araghi, M., Majnon-Jahromi, B., Gilasian, E., Withers, P., Gaimari, S.D. and Fallahzadeh, M. 2018. First Iranian record of the family Odiniidae (Diptera: Opomyzoidea), including two species new to the Middle East region. *Zootaxa* **4471(3)**: 580-584. doi:10.11646/zootaxa.4471.3.9.
- Pauli, T., Burt, T.O., Meusemann, K., Bayless, K., Donath, A., Podsiadlowski, L., Mayer, C., Kozlov, A., Vasilikopoulos, A., Liu, S.L. and others. 2018. New data, same story: phylogenomics does not support Syrphoidea (Diptera: Syrphidae, Pipunculidae). *Systematic Entomology* **43(3)**: 447-459. doi:10.1111/syen.12283.
- Perkovsky, E.E., Sukhomlin, E.B. and Zelenkov, N.V. 2018. An unexpectedly abundant new genus of black flies (Diptera, Simuliidae) from Upper Cretaceous Taimyr amber of Ugolyak, with discussion of the early evolution of birds at high latitudes. *Cretaceous Research* **90**: 80-89. doi:10.1016/j.cretres.2018.04.002.
- Pielowska, A., Sontag, E. and Szadziewski, R. 2018. Haematophagous arthropods in Baltic amber. *Annales Zoologici (Warsaw)* **68(2)**: 237-249.
- Pivar, R.J., Pinho, L.C., Klingeman, W.E., Moulton, J.K. and Sinclair, B.J. 2018. A new genus and species, and first record of the family Thaumaleidae (Diptera) from Brazil. *Canadian Entomologist* **150(4)**: 481-494. doi:10.4039/tce.2018.29.

- Podenas, S. and Byun, H.W. 2018. *Libnotes* crane flies (Diptera: Limoniidae) from Jeju Island (South Korea). *Zootaxa* **4483(2)**: 375-384. doi:10.11646/zootaxa.4483.2.9.
- Pollet, M. and Ivkovic, M. 2018. Dolichopodidae of riverbeds and springs in Croatia with an updated checklist of Croatia (Diptera). *Zootaxa* **4455(3)**: 401-428. doi:10.11646/zootaxa.4455.3.1.
- Pollet, M., Leponce, M., Pascal, O., Touroult, J. and Van Calster, H. 2018. Dipterological survey in Mitaraka Massif (French Guiana) reveals megadiverse dolichopodid fauna with an unprecedented species richness in *Paraclius* Loew, 1864 (Diptera: Dolichopodidae). In: Touroult J. (Ed.). "Our Planet Reviewed" 2015 large-scale biotic survey in Mitaraka, French Guiana. *Zoosystema* **40 (21)**: 471-491. doi; 10.5252/zoosystema2018v40a21.
- Pollet, M. and Stark, A. 2018. The quest for the identity of *Orthoceratium lacustre* (Scopoli, 1763) reveals centuries of misidentifications (Diptera, Dolichopodidae). *Zookeys* **782**: 49-79. doi:10.3897/zookeys.782.26329.
- Pont, A.C. 2018. The Muscidae (Diptera) of Armenia. *Zootaxa* **4465(1)**: 1-69. doi:10.11646/zootaxa.4465.1.1.
- Porter Sanford, D. and Plowes, R.M. 2018. Rearing and biology of the decapitating fly *Pseudacteon bifidus* (Diptera: Phoridae): a parasitoid of tropical fire ants. *Florida Entomologist* **101(2)**: 265-272.
- Radenkovic, S., Velickovic, N., Ssymank, A., Vidakovic, D.O., Djan, M., Stahls, G., Veselic, S. and Vujic, A. 2018. Close relatives of Mediterranean endemorelict hoverflies (Diptera, Syrphidae) in South Africa: Morphological and molecular evidence in the *Merodon melanocerus* subgroup. *Plos One* **13(7)**: 29. doi:10.1371/journal.pone.0200805.
- Ramage, T., Charlat, S. and Mengual, X. 2018. Flower flies (Diptera, Syrphidae) of French Polynesia, with the description of two new species. *European Journal of Taxonomy* **448**: 1-37. doi:10.5852/ejt.2018.448.
- Ricarte, A., Nencioni, A., Tubic, N.K., Grkovic, A., Vujic, A. and Marcos-Garcia, M.A. 2018. The hoverflies of an oak dehesa from Spain, with a new species and other insights into the taxonomy of the *Eumerus tricolor* group (Diptera: Syrphidae). *Annales Zoologici* **68(2)**: 259-280. doi:10.3161/00034541anz2018.68.2.005.
- Riccardi Paula, R., Bazylar, Z. and Lamas, C.E. 2018. New genus of the subfamily Oscinellinae from Brazil (Diptera: Chloropidae). *Zootaxa* **4438(2)**: 394-400.
- Roberts Derek, M. 2018. Predator feeding vibrations encourage mosquito larvae to shorten their development and so become smaller adults. *Ecological Entomology* **43(4)**: 534-537.
- Rohacek, J. 2018. European Chiropteromyzidae (Diptera): Taxonomic revision, nomenclature, classification, and preimaginal stages. *Annales Zoologici* **68(2)**: 281-316. doi:10.3161/00034541anz2018.68.2.006.
- Rosenmejer, T. and Pape, T. 2018. Five new species of the flesh fly genus *Boettcheria* (Diptera: Sarcophagidae). *Zootaxa* **4483(3)**: 579-590. doi:10.11646/zootaxa.4483.3.9.
- Roy, S., Chakraborty, S.K., Parui, P. and Mitra, B. 2018. Taxonomy of soldier flies (Diptera: Stratiomyidae) of Sunderban Biosphere Reserve, India. *Proceedings of the Zoological Society (Calcutta)* **71(2)**: 121-126.
- Rull, J., Tadeo, E., Lasa, R. and Aluja, M. 2018. The effect of winter length on duration of dormancy and survival of specialized herbivorous *Rhagoletis* fruit flies from high elevation environments with acyclic climatic variability. *Bulletin of Entomological Research* **108(4)**: 461-470.
- Runyon, J.B. and Pollet, M. 2018. *Enlinia* Aldrich, 1933 of Mitaraka, French Guiana (Diptera: Dolichopodidae). In: Touroult J. (Ed.). "Our Planet Reviewed" 2015 large-scale biotic survey in Mitaraka, French Guiana. *Zoosystema* **40 (19)**: 453-468. doi:10.5252/zoosystema2018v40a19.

- Sadeghi, M., Altan, E., Deng, X.T., Barker, C.M., Fang, Y., Coffey, L.L. and Delwart, E. 2018. Virome of > 12 thousand *Culex* mosquitoes from throughout California. *Virology* **523**: 74-88. doi:10.1016/j.virol.2018.07.029.
- Sakhvon, V.V. and Lelej, A.S. 2018. Review of the genus *Heteropogon* Loew, 1847 (Diptera: Asilidae) from Russia and Central Asia, with description of two new species. *Zootaxa* **4486(4)**: 435-450. doi:10.11646/zootaxa.4486.4.2.
- Santos, D. and Ribeiro, G.C. 2018. Systematics of the genus *Aphrophila* Edwards with description of fifteen new species (Diptera: Tipulomorpha: Limoniidae). *Zootaxa* **4473(1)**: 1-75. doi:10.11646/zootaxa.4473.1.1.
- Scaccini, D. 2018. Remarks on the biology of *Dinera ferina* (Diptera Tachinidae) as parasitoid of the two Italian *Platycerus* species (Coleoptera Lucanidae). *Bulletin of Insectology* **71(1)**: 39-43.
- Scheffer Sonja, J. and Lonsdale, O. 2018. A survey of Agromyzidae (Diptera) reared from leafmines on Long Island, New York; host associations, distribution data, and the description and host association of a new species. *Zootaxa* **4450(1)**: 77-90.
- Schirmel, J., Albrecht, M., Bauer, P.M., Sutter, L., Pfister, S.C. and Entling, M.H. 2018. Landscape complexity promotes hoverflies across different types of semi-natural habitats in farmland. *Journal of Applied Ecology* **55(4)**: 1747-1758. doi:10.1111/1365-2664.13095.
- Shelly, T.E. 2018. Sexual selection on leks: A fruit fly primer. *Journal of Insect Science* **18(3)**: 16. doi:10.1093/jisesa/iey048.
- Shen, R.R., Shi, L., Li, W.L. and Wang, J.C. 2018. A new species of subgenus *Homoneura* from Northern China, with information of 12 species newly recorded (Diptera: Lauxaniidae: Homoneura). *Zootaxa* **4418(6)**: 501-525. doi:10.11646/zootaxa.4418.6.1.
- Shults, P. and Borkent, A. 2018. Pupae of the Nearctic species of *Culicoides* Latreille subgenus *Monoculicoides* Khalaf (Diptera: Ceratopogonidae). *Zootaxa* **4504 (4)**: 451-472.
- Sinclair Bradley, J. 2018. Correct original spelling of *Hydropeza queenslandensis* Sinclair (Diptera: Empididae). *Zootaxa* **4433(2)**: 386.
- Skibinska, K. and Krzeminski, W. 2018. New species of the genus *Similinannotanyderus* (Tanyderidae, Diptera) from the Myanmar amber. *Cretaceous Research* **90**: 56-59.
- Soares, M.M.M. and Ale-Rocha, R. 2018. Two new species and new occurrences of *Syneches* Walker for Brazilian biome of Caatinga (Diptera: Hybotidae: Hybotinae). *Neotropical Entomology* **47(5)**: 634-641. doi:10.1007/s13744-017-0583-2.
- Song, C., Lin, X.L., Wang, Q. and Wang, X.H. 2018. DNA barcodes successfully delimit morphospecies in a superdiverse insect genus. *Zoologica Scripta* **47(3)**: 311-324. doi:10.1111/zsc.12284.
- Suetsugu, K. and Sueyoshi, M. 2018. Specialized pollination by fungus gnats in the introduced population of *Aspidistra elatior*. *Journal of Plant Research* **131(3)**: 497-503.
- Suh, S.J. and Kwon, Y.J. 2018. Taxonomy of the genus *Atherigona* Rondani (Diptera: Muscidae) from Korea. *Entomological Research* **48(3)**: 187-197. doi:10.1111/1748-5967.12277.
- Tadeo, E., Aluja, M. and Rull, J. 2018. Precopulatory mating and postzygotic isolation between two walnut-infesting species of *Rhagoletis* from Mexican highlands. *Entomologia Experimentalis Et Applicata* **166(9)**: 713-723. doi:10.1111/eea.12720.
- Taillefer, A.G. and Wheeler, T.A. 2018. Tracking wetland community evolution using Diptera taxonomic, functional and phylogenetic structure. *Insect Conservation and Diversity* **11(3)**: 276-293. doi:10.1111/icad.12271.
- Takaoka, H. and Huang, Y.T. 2018. A new black fly species of *Simulium* (*Simulium*) (Diptera: Simuliidae) from Taiwan. *Zootaxa* **4422(1)**: 141-150. doi:10.11646/zootaxa.4422.1.9.

- Takaoka, H., Low, V.L., Tan, T.K., Huang, Y.T., Fukuda, M. and Ya'cob, Z. 2018. A new species of the *Simulium* (*Simulium*) *argentipes* species-group (Diptera: Simuliidae) from Taiwan, and its phylogenetic relationships with four related species. *Journal of Medical Entomology* **55**(4): 884-892. doi:10.1093/jme/tjy028.
- Takaoka, H., Srisuka, W., Low, V.L. and Saeung, A. 2018. A new species and a new record of the *Simulium* (*Gomphostilbia*) *gombakense* species-group (Diptera: Simuliidae) from Thailand. *Acta Tropica* **185**: 156-166. doi:10.1016/j.actatropica.2018.05.011.
- Takaoka, H., Srisuka, W., Van Lun Low, W. and Atiporn, S. 2018. Five new species of the *Simulium decuplum* subgroup of the *Simulium* (*Gomphostilbia*) *batoense* species-group (Diptera: Simuliidae) from Thailand and their phylogenetic relationships. *Acta Tropica* **182**: 271-284.
- Takaoka, H., Ya'cob, Z. and Sofian-Azirun, M. 2018. Classification, annotated list and keys for the black flies (Diptera: Simuliidae) of Peninsular Malaysia. *Zootaxa* **4498**(1): 1-65.
- Tanasak, C., Bhusri, B., Poonyapat, S., Thekhawet, W., Eakanan, N., Theeraphap, C. and Ruangsittichai, J. 2018. Species identification of horse flies (Diptera: Tabanidae) in Thailand using DNA barcoding. *Veterinary Parasitology* **259**: 35-43.
- Theischinger, G., Billingham, Z.D. and Grouns, I. 2018. *Ozeoura*-a new genus of Chioneinae (Insecta: Diptera: Tipuloidea: Limoniidae) from Australia. *Records of the Australian Museum* **70**(5): 447-468. doi:10.3853/j.2201-4349.70.2018.1714.
- Thompson, F.C. 2018. A new flower fly synonym (Diptera: Syrphidae). *Proceedings of the Entomological Society of Washington* **120**(3): 651.
- Tillier, P. and Quindroit, C. 2018. Diptera Ptychopteridae - a new national inventory. *Insectes* **189**: 32-34.
- Touroult, J., Pollet, M. and Pascal, O. 2018. Overview of Mitaraka survey: research frame, study site and field protocols. In: Touroult J. (Ed.) "Our Planet Reviewed" 2015 large-scale biotic survey in Mitaraka, French Guiana. *Zoosystema* **40** (13): 327-365. doi: 10.5252/zoosystema2018v40a13.
- Ventura, C., Briones-Roblero, C.I., Hernandez, E., Rivera-Orduna, F.N. and Zuniga, G. 2018. Comparative analysis of the gut bacterial community of four *Anastrepha* fruit flies (Diptera: Tephritidae) based on pyrosequencing. *Current Microbiology* **75**(8): 966-976. doi:10.1007/s00284-018-1473-5.
- Viviani, V.R., Amaral, D.T., Bevilaqua, V.R. and Falaschi, R.L. 2018. Orfelia-type luciferin and its associated storage protein in the non-luminescent cave worm *Neoditomyia* sp (Diptera: Keroplatidae) from the Atlantic rainforest: biological and evolutionary implications. *Photochemical & Photobiological Sciences* **17**(10): 1282-1288. doi:10.1039/c8pp00207j.
- Vosteen, I., Gershenzon, J. and Kunert, G. 2018. Dealing with food shortage: larval dispersal behaviour and survival on non-prey food of the hoverfly *Episyrphus balteatus*. *Ecological Entomology* **43**(5): 578-590.
- Vujic, A., Radenkovic, S. and Likov, L. 2018. Revision of the Palearctic species of the *Merodon desuturinus* group (Diptera, Syrphidae). *Zookeys* **771**: 105-138. doi:10.3897/zookeys.771.20481.
- Wagner, R. 2018. Remarkable new Australian mothflies (Diptera, Psychodidae). *Zootaxa* **4457**(2): 285-295.
- Wahlberg, E. and Johanson, K.A. 2018. Molecular phylogenetics reveals novel relationships within Empidoidea (Diptera). *Systematic Entomology* **43**(4): 619-636. doi:10.1111/syen.12297.
- Wakie, T., Yee, W.L. and Neven, L.G. 2018. Assessing the Risk of Establishment of *Rhagoletis cerasi* (Diptera: Tephritidae) in the United States and Globally. *Journal of Economic Entomology* **111**(3): 1275-1284. doi:10.1093/jee/toy054.
- Wang, M. and Yang, D. 2018. Three new species of *Argyra* from China (Diptera, Dolichopodidae, Diaphorinae). *Zookeys* **772**: 141-151.

- Wee, S.L., Tan, S.B. and Jurgens, A. 2018. Pollinator specialization in the enigmatic *Rafflesia cantleyi*: A true carrion flower with species-specific and sex-biased blow fly pollinators. *Phytochemistry* **153**: 120-128. doi:10.1016/j.phytochem.2018.06.005.
- Whitmore, D., Buenaventura, E. and Pape, T. 2018. Odd, outsized, and obscure: *Sarcophaga (Hadroxena) karakoncolos* sp n. (Diptera: Sarcophagidae) from Turkey. *Zootaxa* **4422(3)**: 385-394. doi:10.11646/zootaxa.4422.3.4.
- Xi, Y. and Yin, X. 2018. Four new species of *Phyllomyza* Fallen (Diptera: Milichiidae), with an updated key from China. *Zoological Systematics* **43(3)**: 309-316.
- Xi, Y.Q., Yang, D. and Yin, X.M. 2018. The genus *Phyllomyza* Fallen from China, with descriptions of three new species (Diptera, Milichiidae). *Zookeys* **760**: 143-157. doi:10.3897/zookeys.760.22595.
- Xu, W.T., Zhang, D. and Pape, T. 2018. Biology of *Eumacronychia* Townsend, with a redescription of *E. persolla* Reinhard, 1965 (Diptera, Sarcophagidae). *Zookeys* **783**: 55-65. doi:10.3897/zookeys.783.28057.
- Ya'cob, Z., Takaoka, H., Low, V.L. and Sofian-Azirun, M. 2018. Characterization of *Simulium (Simulium) hackeri* Edwards (Diptera: Simuliidae) from Malaysia: Morphological description of the pupa and larva, and DNA barcoding. *Acta Tropica* **185**: 110-114. doi:10.1016/j.actatropica.2018.04.029.
- Ya'cob, Z., Takaoka, H., Van Lun Low, H. and Sofian-Azirun, M. 2018. A new species of *Simulium (Simulium)* (Diptera: Simuliidae) from Genting Highlands, Malaysia. *Acta Tropica* **182**: 1-3.
- Yan, C.C., Liu, T., Cao, W., Zhao, G.J. and Liu, W.B. 2018. A review of the Japanese *Cryptochironomus* Kieffer, 1918 (Diptera, Chironomidae). *Zookeys* **771**: 139-155. doi:10.3897/zookeys.771.24220.
- Yau, T. and Marshall, S.A. 2018. A revision of the genus *Bromeloecia* Spuler (Diptera: Sphaeroceridae: Limosiniinae). *Zootaxa* **4445(1)**: 1-115. doi:10.11646/zootaxa.4445.1.1.
- Yeo, D., Puniamoorthy, J., Ngiam, R.W.J. and Meier, R. 2018. Towards holomorphology in entomology: rapid and cost-effective adult-larva matching using NGS barcodes. *Systematic Entomology* **43(4)**: 678-691. doi:10.1111/syen.12296.
- Ying-Qiang, G., Yun-Yun, G. and Zhang, D. 2018. Two new synonymies in the genus *Lispe* (Diptera: Muscidae). *Zootaxa* **4457(4)**: 549-556.
- Ying-Qiang, G., Zhang, D. and Pape, T. 2018. A new species of *Wohlfahrtia* Brauer & Bergenstamm (Diptera: Sarcophagidae) from northwestern China, with three new synonymies and a pictorial synopsis. *Zootaxa* **4434(1)**: 130-140.
- Yoshizawa, S. and Tachi, T. 2018. Taxonomic review of the genus *Pygophora* Schiner of Japan (Diptera: Muscidae). *Zootaxa* **4418(5)**: 449-468. doi:10.11646/zootaxa.4418.5.3.
- Zatwarnicki, T. 2018. Solving the puzzle of taxonomic position of the petroleum fly by resurrection of *Diasemocera* Bezzi from *Psilopa* Fallen (Diptera: Ephydriidae) with proposed specific and generic synonymies. *Annales Zoologici* **68(3)**: 527-552. doi:10.3161/00034541anz2018.68.3.012.
- Zhang, Y., Feng, S.G., Zeng, Y.Y., Ning, H., Liu, L.J., Zhao, Z.H., Jiang, F. and Li, Z.H. 2018. The first complete mitochondrial genome of *Bactrocera tsuneonis* (Miyake) (Diptera: Tephritidae) by next-generation sequencing and its phylogenetic implications. *International Journal of Biological Macromolecules* **118**: 1229-1237. doi:10.1016/j.ijbiomac.2018.06.099.
- Zhou, J.L., Shamshev, I., Kwon, Y. and Yang, D. 2018. Species of the subgenus *Empis (Xanthempis)* from South Korea (Diptera, Empididae). *Zookeys* **769**: 145-155. doi:10.3897/zookeys.769.24545.

SUBMISSION FORM

DIRECTORY OF NORTH AMERICAN DIPTERISTS

For those who have not yet sent in a synopsis of their interests for the *Directory of North America Dipterists*, the following form is provided. Please restrict yourselves to no more than 20 words when listing the titles of your major projects and the animals you work with. Should any of you like to expand or modify your entries from the last list, use the form to indicate the changes.

The information can be emailed, or the form completed and faxed or mailed to the following address:

Dr. James O'Hara
 Canadian National Collection of Insects
 Agriculture & Agri-Food Canada
 K.W. Neatby Building, C.E.F.
 Ottawa, Ontario, CANADA, K1A 0C6
 Tel.: (613) 759-1795
 FAX: (613) 759-1927
 Email: james.ohara@canada.ca

Full name: _____

Address: _____

_____ **Telephone:** _____

FAX: _____ **Email:** _____

Projects and taxa studied: _____
