



US 20070065461A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2007/0065461 A1**
Guerry et al. (43) **Pub. Date: Mar. 22, 2007**

(54) **IMMUNOGENIC CAPSULE COMPOSITION
FOR USE AS A VACCINE COMPONENT
AGAINST CAMPYLOBACTER JEJUNI**

Related U.S. Application Data

(60) Provisional application No. 60/722,086, filed on Sep. 21, 2005.

(76) Inventors: **Patricia Guerry**, Silver Spring, MD
(US); **Shahida Baqar**, Olney, MD
(US); **Mario Artur Monteiro**, Guelph
(CA)

Publication Classification

(51) **Int. Cl.**
A61K 39/02 (2006.01)
(52) **U.S. Cl.** **424/234.1**

Correspondence Address:
**NAVAL MEDICAL RESEARCH CENTER
ATTN: (CODE 00L)
503 ROBERT GRANT AVENUE
SILVER SPRING, MD 20910-7500 (US)**

(57) **ABSTRACT**

An immunogenic composition, and method of using the composition, composed of a capsule polysaccharide polymer from one or more strains *Campylobacter jejuni*. The composition is either used alone or is conjugated to a carrier molecule, such as CRM₁₉₇. An aspect of the invention is that the immunogenic composition induces an immune response without the induction of Guilliam Barre Syndrome.

(21) Appl. No.: **11/524,057**

(22) Filed: **Sep. 20, 2006**




FIG. 1




FIG. 2




FIG. 3




FIG. 4

US 2007/0065461 A1

Mar. 22, 2007

1

IMMUNOGENIC CAPSULE COMPOSITION FOR USE AS A VACCINE COMPONENT AGAINST CAMPYLOBACTER JEJUNI**CROSS-REFERENCES TO RELATED APPLICATIONS**

[0001] This application claims priority to U.S. Provisional application 60/722,086, filed Sep. 21, 2005 and is hereby incorporated by reference.

FIELD OF INVENTION

[0002] The inventive subject matter relates to an immunogenic composition capable of conferring protection against diarrhea caused by *Campylobacter jejuni* and a method of inducing an immune response to said composition.

BACKGROUND OF INVENTION

[0003] *C. jejuni* is a leading cause of diarrheal disease worldwide and a documented threat to US military personnel (Taylor, 1992; Tauxe, 1992). The symptoms of campylobacter enteritis include diarrhea, abdominal pain, fever and often accompanied by vomiting. Stools usually contain mucus, fecal leukocytes and blood, although watery diarrhea is also observed (Cover and Blaser 1999). However, despite the importance of this organism to human disease, there are no licensed vaccines against *C. jejuni*.

[0004] Because of the medical importance of *C. jejuni*, considerable research is dedicated toward understanding this pathogen. However, notwithstanding this effort, there is surprisingly little understanding about how *C. jejuni* causes human disease. The genome of one strain, NCTC 11168 (Parkhill, et al., 2000) revealed several unusual aspects about the biology of *C. jejuni*. One striking feature is the presence of an unexpectedly high number of genes encoding putative enzymes involved in sugar and/or polysaccharide synthesis (Parkhill et al., 2000). The sequence, and resulting research fostered primarily by the availability of the sequence, has revealed that these genes fall into 4 main functional clusters that underscore the importance of some unusual carbohydrate structure to the biology of *C. jejuni*. These clusters include Lipooligosaccharide (LOS) synthesis, genetic control of flagellin glycosylation, genetic control of N-linked glycosylation and the control of the biosynthesis and assembly of capsule.

[0005] Vaccine strategies against *C. jejuni* have been largely limited due to the molecular mimicry between lipooligosaccharide (LOS) cores of many strains of *C. jejuni* and human gangliosides (Moran, et al., 1996). This mimicry is thought to be a major factor in the strong association of *C. jejuni* infection with Guillain Barre Syndrome (GBS), a post-infectious polyneuropathy (Allos, 1997). Thus, antibodies generated against LOS cores result in an autoimmune response to human neural tissue. It has been estimated that as many as 1/3000 cases of campylobacter enteritis results in GBS. Therefore, the possibility of developing GBS could be associated with any whole cell vaccine against *C. jejuni* that includes ganglioside mimicry.

[0006] LOS synthesis in *Campylobacter* is controlled by a number of genes, including genes encoding enzymes involved in biosynthesis of sialic acid for incorporation into

LOS. Thus, *C. jejuni* is one of a limited number of bacteria that can endogenously synthesize sialic acid, a 9 carbon sugar that is found in many mammalian cells. This is consistent with the observed molecular mimicry of LOS and human gangliosides important in GBS (Aspinall et al., 1993, 1994 (a and b); Salloway et al., 1996).

[0007] Although glycosylation of proteins was once considered to be a eukaryotic trait, there is an increase awareness of prokaryotic protein glycosylation (Power and Jennings, 2003). The best characterized and most extensively glycosylated bacterial protein is campylobacter flagellin. Flagellin from strain 81-176 is glycosylated at 19 serine or threonine sites by an O-linkage to pseudaminic acid and derivatives of pseudaminic acid (Thibault et al., 2001). Pseudaminic acid is an unusual 9 carbon sugar that resembles sialic acid, but which is highly immunogenic, unlike sialic acid. Moreover, mutants that are unable to glycosylate flagellin cannot assemble a flagellar filament (Goon et al, 2003). Since flagella are indispensable virulence determinants of *C. jejuni*, glycosylation is therefore also a key virulence determinant.

[0008] One of the most unusual aspects of *C. jejuni* is the presence of a general system for N-linked glycosylation of numerous proteins (Szymanski et al., 1999; reviewed in Szymanski et al., 2003). This system, which includes an oligosaccharide transferase similar to that found in the eukaryote *Saccharomyces cerevisiae*, attaches a glycan which has recently been shown to be a heptasaccharide composed of one bacillosamine residue (an unusual deoxy sugar), one D-glucose, and five D-GalNAc residues (Young et al., 2002). The glycosylation appears to occur on numerous periplasmic, and perhaps, surface exposed proteins in *C. jejuni* (Young et al., 2002). The unusual glycan, again, appears to be highly immunogenic and is recognized during human infection (Szymanski et al., 1999, 2003).

[0009] An interesting recent revelation regarding the Campylobacter genome sequence was the presence of a complete set of capsule transport genes similar to those seen in type II/III capsule loci in the *Enterobacteriaceae* (Parkhill et al., 2000; Karlyshev et al., 2000). Subsequent genetic studies in which site-specific mutations were made in several capsule transport genes indicated that the capsule was the serodeterminant of the Penner serotyping scheme (Karlyshev et al., 2000; Bacon et al., 2001). The Penner scheme (or HS for heat stable) is one of two major serotyping schemes of campylobacters and was originally thought to be based on lipopolysaccharide O side chains (Moran and Penner, 1999).

[0010] Currently it is believed that all of the structures previously described as O side chains are, in fact, capsules. The chemical structures of the capsule/O side chains of several Penner serotypes have been determined, and these structures include several unusual sugar structures, as summarized in Table 1. Thus, the capsule of the genome strain, NCTC 11168, contains a heptopyranose as a L-gluco conformer, which is the first report of such a structure in nature (St. Michael et al., 2002). The capsule of the type strains HS23 and HS36 contain the same carbohydrates in different ratios, and include a mixture of 4 unusual altro-heptoses (6-deoxy- α -D-altro-heptose, D-glycero- α -D-altro-heptose, 6-deoxy-3-Me- α -D-altro-heptose, and 3-Me-D-glycero- α -D-altro-heptose (Aspinall et al., 1992).

TABLE 1

Structure of some capsular polysaccharides of <i>C. jejuni</i> strains.		
Strain	Structure	Reference
HS3	$[->4)-\alpha-D\text{-galactose-(1-3)-3 hydroxypropanoyl)-L-}\alpha\text{-D-ido-heptose)-(1->)}_n$	Aspinall et al., 1995
HS19	$\beta\text{-D-glucuronic acid amidated with 2-amino-2-deoxyglycerol linked to GluNAc}$	Aspinall et al., 1994 a, b
HS23, HS36	$\text{GlcNAc-Gal-D-glycero-}\alpha\text{-D-heptose)}_n$, where the D-glycero- $\alpha\text{-D-heptose}$ is a mixture of 4 altro-heptoses	Aspinall et al., 1992
81116	Two polysaccharides at a ratio of 3A:1B:A = glucose, glucuronic acid, and mannose that is O-acetylated; B = glucose, galactose, and GlcNAc.	Muldoon et al. (2002)
NCTC 11168	Uronic acid amidated with 2-amino-2-deoxyglycerol at C6 and 6-O-methyl-D-glycero- $\alpha\text{-L-gluco-heptopyranose}$ as a side branch.	St. Michael et al. (2002)

[0011] There are several examples of highly effective capsular vaccines. *S. pneumoniae* has 83 different capsular types, but the current *S. pneumoniae* vaccine contains a mixture of the 23 most prevalent serotypes in the US and Europe. *N meningitidis* has fewer serogroups, thus potentially simplifying vaccine development, and, in fact, serogroups A, B and C are responsible for >90% of cases of meningococcal meningitis (Jennings, 1990). However, the polysaccharide from serotype B is poorly immunogenic in man, likely because it mimics human tissues. Capsular vaccines have also been developed against *H. influenzae* and Group B *Streptococcus*.

[0012] As previously mentioned, there currently are no licensed vaccines against *Campylobacter*, due greatly to the molecular mimicry between LOS cores of many strains of *C. jejuni* and human gangliosides (Moran, et al., 1996). However, vaccine formulations incorporating bacterial capsules have been developed against a number of pathogens. In general, capsule vaccines are immunogenic in humans and non-toxic (Jennings, 1990). One of the general problems associated with capsule vaccines is the poor immunogenicity of all polysaccharides in infants, and the fact that many of the capsular vaccines are directed at diseases that are particularly threatening to the pediatric population. Based on murine studies, pure polysaccharide antigens are considered to be T cell independent, and capable of inducing only IgM type responses. Adult humans, in contrast, are able to generate IgG, in addition to IgM and IgA antibodies against polysaccharides. Responses in infants to vaccines against type B *H. influenzae* (Schneerson et al 1980; Anderson, 1983; Marburg, 1986), group A, B and C *Neisseria meningitidis* (Jennings and Lugowski, 1981 and 1983; and type 6A *Streptococcus pneumoniae* (Chu et al., 1983) have all improved following conjugation to proteins.

SUMMARY OF INVENTION

[0013] Vaccines are the preferred method of conferring anti-diarrhea protection in populations that are potentially exposed to diarrheagenic bacteria. However, currently there are no licensed effective vaccines against *Campylobacter jejuni*.

[0014] An object of this invention is an anti-*C. jejuni* immunogenic composition, composed of a capsule polysaccharide polymer, capable of inducing an immune response against important pathogenic strains *C. jejuni* without concomitantly inducing GBS.

[0015] Another object of the invention is an anti-*C. jejuni* prophylactic formulation with enhanced T-cell dependent immunity to important pathogenic strains of the bacteria by conjugating the capsule of *C. jejuni* to T-dependent carrier molecules, such as CRM₁₉₇.

[0016] Another object of the invention is a method of administering the carrier conjugated or unconjugated anti-*C. jejuni* capsule polysaccharide composition in order to induce an immune response.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] FIG. 1. Schematic diagram showing conjugation of purified capsules of *C. jejuni* strains 81-176 and CG 8484 to CRM₁₉₇. In *C. jejuni* 81-176 capsule, position 0-2 of Gal unit is partially (approximately 50-75%) substituted by O-methyl-phosphoramidate.

[0018] FIG. 2. ELISA results showing antibody response following subcutaneous administration of *C. jejuni* strain 81-176 capsule conjugated to CRM.

[0019] FIG. 3. ELISA results of study illustrating longevity of immune response to capsule conjugated to CRM.

[0020] FIG. 4. ELISA results of serum titer of capsule-specific antibody illustrating IgG response in mice protected from challenge by *C. jejuni*.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0021] *C. jejuni* capsular moieties are important in serodetermination. However, despite over 60 Penner serotypes having been identified, most *Campylobacter* diarrheal disease is caused by *C. jejuni* expressing only a limited number of serotypes. Because of the importance of capsule structure in serodetermination, it is postulated that they are highly immunogenic structures. Additionally, they are also unlikely to exhibit the unwanted autoimmune induction caused by immuno-mimicry observed by lipooligosaccharides. Therefore, capsules or capsular components would be highly useful in anti-*C. jejuni* vaccines. *C. jejuni* capsule, as defined in this application, is a generic term for capsular polymers, which are composed of repeating polysaccharide structures. The repeating structures can be homopolymers, defined as a repeating single sugar moiety, or repeating disaccharides or trisaccharides. A number of species of capsular repeating polysaccharide polymers have been identified. To illustrate the genus of capsular polysaccharide structures, Table 2 lists known capsular polysaccharide structures for *Campylobacter* strains.

[0022] The chemical composition of *C. jejuni* capsules were analyzed by first growing *C. jejuni* and then purifying the capsule using water-phenol extraction, ultra-centrifugation and gel permeation chromatography. The specific carbohydrate structures were determined by gas-liquid chromatography (GLC), and GLC-mass spectrometry, and fast atom bombardment-mass spectrometry (FAB-MS). Anomeric configuration of the sugars was determined by nuclear magnetic resonance (NMR) spectrometry.

TABLE 2

Strain/HS type	Structure	Reference
HS3 type strain	$\rightarrow 4\text{-}\alpha\text{-D-Gal-(1}\rightarrow 3\text{)(3-hydroxypropanoyl)-L-glycero-}\alpha\text{-D-ido-Hep-(1}\rightarrow$	Aspinall et al. (1995)
HS19 type strain	$\rightarrow 4\text{-}\beta\text{-D-GlcA-(1}\rightarrow 3\text{)-}\alpha\text{-D-GlcNAc-(1}\rightarrow$ (the GlcA units are present as amides of 2-amino-2-deoxyglycerol)	Aspinall et al., 1994 a and b
HS23/36 type strain	Four closely-related polysaccharides: $\rightarrow 3\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow 3\text{)-}\alpha\text{-D-Gal-(1}\rightarrow 2\text{)-6d-}\alpha\text{-D-altro-Hep-(1}\rightarrow$; $\rightarrow 3\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow 3\text{)-}\alpha\text{-D-Gal-(1}\rightarrow 2\text{)-6d-3-O-Me-}\alpha\text{-D-altro-Hep-(1}\rightarrow$; $\rightarrow 3\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow 3\text{)-}\alpha\text{-D-Gal-(1}\rightarrow 2\text{)-D-glycero-}\alpha\text{-D-altro-Hep-(1}\rightarrow$; $\rightarrow 3\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow 3\text{)-}\alpha\text{-D-Gal-(1}\rightarrow 2\text{)-3-O-Me-D-glycero-}\alpha\text{-D-altro-Hep(1}\rightarrow$.	Aspinall et al., 1992
81116 (HS6)	Two polysaccharides at a ratio of 3A:1B, where <div style="text-align: center;"> $\begin{array}{ccc} \text{OAc (30\%)} & & \text{OAc (20\%)} \\ \downarrow & & \downarrow \\ 3 & & 6 \end{array}$ </div> A = $\rightarrow 3\text{-}\beta\text{-D-Glc-(1}\rightarrow 2\text{)-}\alpha\text{-D-GlcA-(1}\rightarrow 3\text{)-}\alpha\text{-D-Man-(1}\rightarrow 3\text{)-}\alpha\text{-D-Glc-(1}\rightarrow$ B = $\rightarrow 3\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow 6\text{)-}\alpha\text{-D-Glc-(1}\rightarrow 4\text{)-}\alpha\text{-D-Gal-(1}\rightarrow$ <div style="text-align: center;"> $\begin{array}{c} 3 \\ \uparrow \\ \beta\text{-D-GlcNAc-(1} \\ \downarrow \\ 6\text{-O-Me-D-L-}\alpha\text{-L-glc-Hepp-(1} \end{array}$ </div>	Muldoon et al. 2002
NCTC 11168 (HS2)	$\rightarrow 2\text{-}\beta\text{-D-Ribf-(1}\rightarrow 5\text{)-}\beta\text{-D-GalfNAc-(1}\rightarrow 4\text{)-}\alpha\text{-D-GlepA6(NGro)-(1}\rightarrow$ (Here, Glucuronic acid is amidated with 2-amino-2-deoxyglycerol at C-6)	St. Michael et al., 2002
HS41 type strain	Two closely related polysaccharides: $\rightarrow 2\text{-}\beta\text{-L-Araf-(1}\rightarrow 2\text{)-}\beta\text{-D-6d-altro-Hepf-(1}\rightarrow 2\text{)-}\beta\text{-L-6d-altrof-(1}\rightarrow$ (75%); and $\rightarrow 2\text{-}\beta\text{-L-Araf-(1}\rightarrow 2\text{)-}\beta\text{-D-6d-altro-Hepf-(1}\rightarrow 2\text{)-}\alpha\text{-D-Fucf-(1}\rightarrow$ (25%)	Hannify et al., 1999
HS30 (<i>C. coli</i>) Type strain	$\rightarrow 5\text{-Ribitol-1-P}\rightarrow$ <div style="text-align: center;"> $\begin{array}{c} 2 \\ \uparrow \\ 6\text{d-}\beta\text{-D-talo-Hep-(1}\rightarrow 4\text{)-}\beta\text{-D-GlcNAc-(1} \end{array}$ </div>	Aspinall et al. 1993
HS1 type strain	$\rightarrow 4\text{-}\beta\text{-D-Gal-(1}\rightarrow 2\text{)-(R)-Gro-(1-P}\rightarrow$ (with two branches at C-2 and C-3 of Gal of $\beta\text{-D}$ fructofuranoses that are further substituted at C-3 with O-methyl phosphoramidate groups)	McNally et al., 2005
HS4 (strain 8486)	$\rightarrow 6\text{-d-}\alpha\text{-D-ido-Hep-(1}\rightarrow 4\text{)-beta-D-GlcNAc-(1}\rightarrow$	unpublished

[0023] Based on these analyses, the capsule of *C. jejuni* strain 81-176 was determined to be a repeating polymer of three carbohydrates, illustrated by the formula: $\alpha\text{-D-Gal(1-2)-}\alpha\text{-6-deoxy-3-Me-D-altro-heptose(1,3)}\beta\text{-D-acetylglucosamine}([\rightarrow 3\text{-}\alpha\text{-D-Gal-(1}\rightarrow 2\text{)-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3\text{)-}\beta\text{-D-GlcNAc-(1}\rightarrow]_n$. Position O-2 of Gal unit is partially (approximately 50 to 75%) substituted by O-methyl-phosphoramidate.

[0024] Additionally, the capsule of strain CG8486 was analyzed and shown to be composed of a similar structure but of a repeating disaccharide illustrated by the formula: $\rightarrow 3\text{-}\alpha\text{-6-deoxy-D-ido-Heptose-(1}\rightarrow 4\text{)-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

[0025] Therefore, an aspect of this invention is a vaccine formulation comprised of one or a plurality of species of the genus of capsule polymers of repeating polysaccharides of *C. jejuni*.

EXAMPLE 1

Immune Response Induced by Anti-*C. jejuni* Composition Containing *C. jejuni* Strain 81-176 Capsule

[0026] In order to demonstrate the immunogenicity of *C. jejuni* capsule, BALB/c mice were immunized subcutaneously with either phosphate buffer saline (PBS) or purified capsule from *C. jejuni* 81-176. The ensuing immune reaction was measured by enzyme-linked immunosorbent assay (ELISA) using bovine serum albumin (BSA) as a control. In this analysis, antigen was purified capsule.

[0027] In the study, immunization with PBS or 3 doses of 25 μg of capsule did not result in any immune response (secretory IgA in feces, serum IgG or serum IgM).

[0028] Immunization with 3 doses of 100 μg or 2 doses of 400 μg of capsule (see below) also resulted in no sIgA

response. However, 1/5 animals that received 3 doses of 100 µg showed a serum IgG response (titer of 1:160) and 3/5 animals showed a serum IgM response (titers of 1:4000-1:6000). After 2 doses of 400 µg 3/5 animals had a serum IgG response (titers ranging from 1:640-1:1280) and 5/5 had a serum IgM response (titers ranging from 1:6000-1:10,000). This group did not receive a third immunization due to the strong immune response after only 2 immunizations.

[0029] Although the capsule immunogen in this example was administered subcutaneously without adjuvant, subcutaneous administration with adjuvant is contemplated, including but not limited to LTR192G, Aluminum hydroxide, RC529E, QS21, E294, oligodeoxynucleotides (ODN), CpG-containing oligodeoxynucleotides, aluminum phosphate, MPL® (GlaxoSmithKline, Middlesex, UK) or combinations of these or other potential adjuvants. Additionally, although the example illustrates subcutaneous administration, it is also contemplated that administration may be intranasally, with or without adjuvant.

EXAMPLE 2

Immunity to Capsule—CRM₁₉₇ Conjugate

[0030] Although significant IgG response was observed in response to purified capsule, in mice, immunity to polysaccharides are often associated with T-cell independent immune responses. Therefore, children are typically only capable of mounting an IgM response in the face of polysaccharide antigens with adults capable of generating an IgG, IgA and IgM response.

[0031] In order to potentially further improve the response to capsule moieties, the immunogenicity of *C. jejuni* capsule was evaluated as a conjugate to T-dependent carrier proteins. A preferred carrier is CRM₁₉₇, a nontoxic version of diphtheria toxin that has been successfully used in pneumococcal conjugate vaccines (Anderson, 1983). Referring to FIG. 1, capsule-CRM conjugate was prepared by oxidizing purified *C. jejuni* strain 81-176 capsule with periodate and linking the oxidized product to CRM. In FIG. 1, conjugation was conducted on capsule polysaccharide moiety [→3)-α-D-Gal-(1→2)-6d-α-D-altro-Me-Hep-(1→3)-β-D-GlcNAc-(1→)] or to the polysaccharide moiety α-6-deoxy-ido-Hep-tose (1-4)-β-GlcNAc.

[0032] Immunogenicity of the capsule-CRM conjugate was evaluated in Balb/c mice. Five groups of mice were immunized subcutaneously with 3 doses in 14-day intervals as illustrated in Table 3. Blood was collected on days: -1, 14, 28, 56 and 92. A determination whether the animals had seroconverted to capsule was made by measuring IgM and IgG responses by enzyme-linked immunosorbent assay (ELISA). Endpoint titers for IgM and IgG, shown in FIG. 2, was >56 and >32, respectively. Endpoint titer data is presented as loge. The percent of responding mice to capsule is illustrated in Table 4.

TABLE 3

Group	Capsule-CRM ₁₉₇ (µg)	CRM ₁₉₇ (µg)
A	—	—
B	—	12
C	—	50
D	25	—
E	100	—

[0033]

TABLE 4

CPS-CRM dose (µg)	Ig	% Responders to CPS on study day*				
		14	28	42	56	92
25	M	80	80	80	60	40
	G	0	20	80	80	80
100	M	100	80	100	100	80
	G	0	0	100	20	0

[0034] As demonstrated in Table 4 and FIG. 2, conjugated capsule induced a strong antibody response to capsule. No seroconversion to capsule was seen in animals receiving phosphate buffered saline only. Although this example illustrates the invention using capsule conjugate from *C. jejuni* 81-176, the invention also contemplates the use of capsule from other strains of *C. jejuni* in the production of anti-diarrheal vaccines.

EXAMPLE 3

Longevity of Immune Response in Mice to Capsule-CRM Conjugate

[0035] In order to evaluate the longevity of the immune response induced by capsule conjugated to CRM₁₉₇, serum titers were monitored in immunized mice over an extended period. In the study, mice (n=10) were immunized subcutaneously with 3 doses of phosphate buffered saline (PBS)(0, 5, 10, 25 or 50 µg of *C. jejuni* capsule conjugated to CRM₁₉₇ at 14 day intervals (i.e., study days 0, 14, and 28). Blood samples were collected at intervals before immunization through study day 304 and anti-capsule IgG was determined by enzyme-linked immunosorbent assay (ELISA).

[0036] Referring to FIG. 3, in mice immunized with PBS, negligible anti-capsule specific antibodies were detected, which remained unchanged during the course of the study. After 2 doses a low, but easily detectable level, of IgG was detected by ELISA. This response was enhanced following the 3rd dose. Although a robust immune response was detected, the study did not yield a discernable vaccine dose response, possibly indicating that dosage levels were initially higher than required. Serum IgG levels peaked at approximately day 90-150 and remained sustained during the experiment. A slight decline in IgG levels was evident after day 196.

EXAMPLE 4

Efficacy of Capsule—CRM Conjugate

[0037] Evaluation of the efficacy of *C. jejuni* capsule—CRM₁₉₇ formulation was evaluated in mice. In this study, mice were immunized with 3 doses of 5, 10 or 25 µg of capsule—CRM₁₉₇ conjugate at 14 day intervals (i.e., days 0, 14 and 28). Control animals received PBS. All animals (n=7) were challenged on day 120, with 4×10⁹ cfu of *C. jejuni* 81-176. Following challenge, animals were followed for six consecutive days for the development of infection associated illness. Based on the severity of sickness, a score was assigned to each animal as follows:

[0038] 0=no apparent illness; 1=ruffled fur;

[0039] 2=ruffled fur and hunched back; 3=dead.

[0040] Daily sickness index and group average indices were calculated. In addition, before challenge and after challenge, loss in body weights were determined. Vaccine efficacy based on illness and loss in body weights was calculated based on the following formula:

$$\frac{(\text{control}-\text{vaccinated})/(\text{control})\times 100}{}$$

[0041] Referring to FIG. 4, similar capsule-specific serum IgG was seen following immunization with 5, 10 or 25 µg of capsule—CRM₁₉₇. As illustrated in Table 5, a protective immune response was observed with a dose response in terms of protective efficacy. The sickness index was lowest in the group immunized with 25 µg of capsule—CRM₁₉₇ and highest in group immunized with 5 µg of capsule—CRM₁₉₇ conjugate. This dose response was also seen when changes in % body weight were used to evaluate efficacy.

[0042] Further characterization of humoral immune responses, capsule-specific serum IgG subclasses (i.e., IgG1, IgG2a and IgG2b) were determined on a limited number of samples. Animals which showed at least 50% protection tended to have higher levels of IgG2b than IgG2a, with a 2b/2a ratio of 4.3+/-1.8 for 9 protected animals and 2.6+/-0.8 for non-protected animals. Higher IgG2b levels are an indication of a Th1 type immune response.

TABLE 5

CPS_CRM µg	C. jejuni challenge outcome:		% efficacy based on:	
	Sickness Index	Body weight	Sickness Index	Body weight
0	1.19 ± 0.28	14.5 ± 5.9	n/a	n/a
5	0.69 ± 0.33	10.1 ± 2.9	42.1	30.3
10	0.73 ± 0.31	9.1 ± 4.4	38.9	37.3
25	0.38 ± 0.19	6.6 ± 3.0	68.0	54.4

EXAMPLE 5

Prophetic Example of Induction of Immunity to Capsule in Humans

[0043] An aspect of this invention is the ability of *C. jejuni* capsule to induce a vigorous and efficacious immune response in humans but not induction of contraindicating Guillain Barre Syndrome. Optimal methods for inducing protective immunity in humans are preceded by studies in animals such as in mice and monkeys. For each vaccine formulation containing capsules from a single or mixtures of *C. jejuni* strains, a limited amount of experimentation is required to ascertain the optimal effective dose ranges. However, a prophetic method for the induction of anti-*C. jejuni* medicated diarrheal protective immunity contains the following steps:

[0044] a. priming is by administration of an immunogenic formulation containing *C. jejuni* capsule from a single *C. jejuni* or multiple strains of *C. jejuni* bacteria. Alternatively or in addition to purified capsule, the formulation can contain *C. jejuni* capsule conjugated to a T-dependent carrier molecule, such as CRM₁₉₇. The vaccine formulation can be administered orally,

nasally, subcutaneously, intradermally, transdermally, transcutaneously intramuscularly, or rectally. Depending on the route of administration, the vaccine formulation can be administered with or without any of a number of adjuvants, including but not limited to LTR 1 92G, Aluminum hydroxide, RC529E, QS21, E294, oligodeoxynucleotides (ODN), CpG-containing oligodeoxynucleotides, aluminum phosphate, MPL® (GlaxoSmithKline, Middlesex, UK) or combinations of these or other potential adjuvants. The range of a unit dose of immunogen is 0.1 µg to 10 mg of immunogen in a range of buffer solutions.

[0045] b. Subsequent to a priming dose, 2 to 4 boosting doses can also administered with unit dose range of 0.1 µg to 10 mg of immunogen in a buffered aqueous solutions with or without adjuvant.

REFERENCES

[0046] 1. Allos, B. M. 1997. Association between *Campylobacter* infection and Guillain Barre Syndrome. J. Infect. Dis. 176(Suppl 2):S125-128.

[0047] 2. Anderson, P. W. 1983. Antibody responses to *Haemophilus influenzae* type b and diphtheria toxin induced by conjugates of oligosaccharides of the type b capsule with the non toxic protein CRM₁₉₇. Infect. Immun. 39:233-238.

[0048] 3. Aspinall, G. O., S. Fujimoto, A. G. MacDonald, H. Pang, L. A. Kuryanczyk and J. L. Penner. 1994a. Lipopolysaccharides from *Campylobacter jejuni* associated with Guillain Barre Syndrome patients mimic human gangliosides in structure. Infect. Immun. 62:2122-2125.

[0049] 4. Aspinall, G. O., McDonald, A. G., and Pang, H. 1992. Structures of the O chain from lipopolysaccharides of *Campylobacter jejuni* serotypes O:23 and O:36. Carbohydr. Res. 231:13-20.

[0050] 5. Aspinall, G. O., A. G. MacDonald, H. Pang, L. A. Kuranczyk, and J. L. Penner. 1994b. Lipopolysaccharides of *Campylobacter jejuni* serotype O:19: structures of core oligosaccharide regions from the serostrain and two bacterial isolates from patients with Guillain Barre Syndrome. Biochem. 33:241-249.

[0051] 6. Aspinall, G. O., A. G. MacDonald, T. S. Raju, H. Pang, L. A. Kuryanczyk, J.L.Penner and A. P. Moran. 1993. Chemical structure of the core region of *Campylobacter jejuni* serotype O:2 lipopolysaccharide. Eur. J. Biochem. 213:1029-1037.

[0052] 7. Aspinall, G. O., C. M. Lynch, H. Pang, R. T. Shaver, A. P. Moran. 1995. Chemical structures of the core region of *Campylobacter jejuni* O:3 lipopolysaccharide and an associated polysaccharide. Eur. J. Biochem. 1995. 231(3): 570-578.

[0053] 8 Baqar, S., Bourgeois, A. L., Applebee, L A., Mourad, A. S., Kleinosky, M. T., Mohran, Z., and J. R. Murphy. 1996. Murine intranasal challenge model for the study of *Campylobacter* pathogenesis and immunity. Infect. Immun. 64:4933-4939.

[0054] 9. Chu, C., Schneerson, R., Robbins, J. B. and Rastogi, S. C. 1983. Further studies on the immunoge-

US 2007/0065461 A1

Mar. 22, 2007

6

- nicity of *Haemophilus influenzae* type b and pneumococcal type 6A polysaccharide-protein conjugates. *Infect. Immun.* 40:245-256.
- [0055] 10. Cover, T. L. and M. J. Blaser. 1999. The pathobiology of *Campylobacter* infections in humans. *Ann. Rev. Med.* 40:269-185.
- [0056] 11. Goon, S., J. F. Kelly, S. M. Logan, C. P. Ewing, P. Guerry. 2003. Pseudaminic acid, the major modification of *Campylobacter* flagellin, is synthesized via the Cj1293 gene. *Mol. Microbiol.* 50(2):659-671.
- [0057] 12. Hanniffy, O. M., A. S. Shashkov, A. P. Moran, M. M. Prendergast, S. N. Senchenkova, Y. A. Knirel, and A. V. Savage. 1999. Chemical structure of a polysaccharide from *Campylobacter jejuni* 176.83 (serotype O:41) containing only furanose sugars. *Carbohydr. Res.* 319:124-132.
- [0058] 13. Jennings, H. J. Capsular polysaccharides as vaccine candidates. 1990. *Curr. Top. Microbiol. Immunol.* 150:97-127.
- [0059] 14. Jennings, H. J. and Lugowski, C. 1981. Immunochemistry of groups A, B and C meningococcal polysaccharides-tetanus toxoid conjugates. *J. Immunol.* 127:1011-1018.
- [0060] 15. Jennings, H. J., C. W. Lugowski, F. E. Ashton, J. A. Ryan. 1983. The structure of the capsular polysaccharide obtained from a new serogroup (L) of *Neisseria meningitidis*. *Carbohydr. Res.* 112(1):105-111.
- [0061] 16. Karlyshev, A. V., Linton, D., Gregson, N. A., Lastovica, A. J. and Wren, B. W. 2000. **Genetic and biochemical evidence of a *Campylobacter jejuni* capsular polysaccharide that accounts for Penner serotype specificity.** *Mol. Microbiol.* 35:529-541.
- [0062] 17. Marburg, S., Jom, D., Tolman, R. L., Arison, B., McCauley, J., Kniskern, P. J., Hagopian, A. and Vella, P. P. 1986. Biomolecular chemistry of macromolecules: synthesis of bacterial polysaccharide conjugates with *Neisseria meningitidis* membrane protein. *J. Am. Chem. Soc.* 108:5282-5287.
- [0063] 18. McNally, D. J., H. C. Jarrell, J. Li, N. H. Khieu, E. Vinogradov, C. M. Szymanski, and J. R. Brisson. 2005. The HS: 1 serostrain of *Campylobacter jejuni* has a complex teichoic acid-like capsular polysaccharide with nonstoichiometric fructofuranose branches and O-methyl phosphoramidite groups. *FEBS J.* 272:4407-4422.
- [0064] 19. Moran, A. P., B. J. Appelmelk, and G. O. Aspinall. 1996. Molecular mimicry of host structures by lipopolysaccharides of *Campylobacter* and *Helicobacter* spp.: implications in pathogenesis. *J. Endotox. Res.* 3(6):521-531.
- [0065] 20. Moran, A. P. and J. L. Penner. 1999. Serotyping of *Campylobacter jejuni* based on heat-stable antigens: relevance, molecular basis and implications in pathogenesis. *J. Appl. Microbiol.* 86:361-377.
- [0066] 21. Muldoon, J., A. S. Shashkov, A. P. Moran, J. A. Ferris, S. N. Senchenkova, and A. V. Savage. 2002. Structures of two polysaccharides of *Campylobacter jejuni* 81116. *Carbo. Res.* 337:2223-2229.
- [0067] 22. Parkhill, J., B. W. Wren, K. Mungall, J. M. Ketley, C. Churcher, D. Basham, T. Chillingworth, R. M. Davies, T. Feltwell, S. Holroyd, K. Jagels, A. V. Karlyshev, S. Moule, M. J. Pallen, C. W. Penn, M. A. Quail, M. A. Rajandream, K. M. Rutherford, A. H. M. van Vliet, S. Whitehead, and B. G. Barrell. 2000. The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable tracts. *Nature* 403:665-668.
- [0068] 23. Power, P. M. and Jennings, M. P. 2003. The genetics of glycosylation in gram negative bacteria. *FEMS Microbiol. Lett.* 218:211-222.
- [0069] 24. Russell, R. G., M. J. Blaser, J. I. Sarmiento and J. Fox. 1989. Experimental *Campylobacter jejuni* infection in *Macaca nemestrina*. *Infect. Immun.* 57:1438-1444.
- [0070] 25. Russell, R. G., M. O'Donnoghue, D. C. Jr. Blake, J. Zulty and L. J. DeTolla. 1993. Early colonic damage and invasion of *Campylobacter jejuni* in experimentally challenged infant *Macaca mulatta*. *J. Infect. Dis.* 168:210-215.
- [0071] 26. Salloway, S., L. A. Mermel, M. Seamans, G. O. Aspinall, J. E. Nam Shin, L. A. Kurjanczyk, and J. L. Penner. 1996. Miller Fisher Syndrome associated with *Campylobacter jejuni* bearing lipopolysaccharide molecules that mimic human ganglioside GD₃. *Infect. Immun.* 64:2945-2949.
- [0072] 27. Schneerson, R., Barrera, O., Sutton, A. and Robbins, J. B. 1980. Preparation, characterization and immunogenicity of *Haemophilus influenzae* type b polysaccharide protein conjugates. *J. Exp. Med.* 152:361-376.
- [0073] 28. St. Michael, F., C. M. Szymanski, J. Li, K. H. Chan, N. H. Khieu, S. Larocque, W. W. Wakarchuk, J. -R. Brisson, and M. A. Monteiro. 2002. The structures of the lipooligosaccharide and capsule polysaccharide of *Campylobacter jejuni* genome sequenced strain NCTC 11168. *Eur. J. Biochem.* 269:5119-5136.
- [0074] 29. Szymanski, C. M., Yao, R., Ewing, C. P., Trust, T. J., and Guerry, P. (1999) Evidence for a system of general protein glycosylation in *Campylobacter jejuni*. *Mol Microbiol* 32: 1022-1030.
- [0075] 30. Szymanski, C. M., Logan, S. M., Linton, D., and Wren, B. W. 2003. *Campylobacter*-a tale of two protein glycosylation systems. *Trends Microbiol.* 11:233-238.
- [0076] 31. Tauxe, R. V. (1992) Epidemiology of *Campylobacter jejuni* infections in the United States and other industrialized nations. In *Campylobacter jejuni: Current status and future trends* (Edited by Nachamkin I., Blaser M. J. and Tompkins L. S.), p. 9. American Society for Microbiology, Washington, D.C..
- [0077] 32. Taylor, D. N. (1992) *Campylobacter* infections in developing countries. In *Campylobacter jejuni: Current status and future trends* (Edited by Nachamkin I., Blaser M. J. and Tompkins L. S.), p. 20. American Society for Microbiology, Washington, D.C..
- [0078] 33. Thibault, P., Logan, S. M., Kelly, J. F., Brisson, J. -R., Ewing, C. P., Trust, T. J., and Guerry, P. (2001)

US 2007/0065461 A1

Mar. 22, 2007

7

Identification of the carbohydrate moieties and glycosylation motifs in *Campylobacter jejuni* flagellin. J Biol Chem 276: 34862-34870.

[0079] 34. Young, N. M., Brisson, J. -R., Kelly, J., Watson, D. C., Tessier, L., Lanthier, P. H., Jarrell, H. C., Cadotte, N., St. Michael, F., Aberg, E., and Szymanski, C. M. 2002. Structure of the N linked glycan present on multiple glycoproteins in the gram negative bacterium, *Campylobacter jejuni*. J. Biol. Chem. 277:42530-42539.

What is claimed is:

1. An immunogenic composition, wherein said composition is composed of *Campylobacter jejuni* capsule polysaccharide polymer from one or more *Campylobacter jejuni* strains.

2. The immunogenic composition of claim 1, wherein said polysaccharide is composed of a repeating structure consisting of a homopolymer, disaccharide or trisaccharide.

3. The immunogenic composition of claim 2, wherein said polysaccharide polymer is a repeating trisaccharide structure having the formula $[\rightarrow 3)\text{-}\alpha\text{-D-Gal-(1}\rightarrow 2)\text{-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow]_n$.

4. The immunogenic composition of claim 2, wherein said polysaccharide polymer is a repeating disaccharide structure having the formula $\rightarrow 3)\text{-}\alpha\text{-6-deoxy-D-ido-Heptose (1-4)-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

5. An immunogenic composition, wherein said composition is composed of *Campylobacter jejuni* capsule polysaccharide polymer from one or more strains of *Campylobacter jejuni* conjugated to a carrier molecule.

6. The immunogenic composition of claim 5, wherein said capsule is composed of a polysaccharide polymer that is a repeating structure consisting of a homopolymer, disaccharide or trisaccharide.

7. The immunogenic composition of claim 5, wherein said carrier molecule is CRM₁₉₇.

8. The immunogenic composition of claim 7, wherein said carrier molecule is conjugated by reductive amination.

9. The immunogenic composition of claim 5, wherein said capsule polysaccharide is composed of a repeating structure having the formula $[\rightarrow 3)\text{-}\beta\text{-D-Gal-(1}\rightarrow 2)\text{-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow]_n$.

10. The immunogenic composition of claim 5, wherein said capsule polysaccharide is composed of a repeating structure having the formula $\rightarrow 3)\text{-}\alpha\text{-6-deoxy-D-ido-Heptose (1}\rightarrow 4)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

11. The immunogenic composition of claim 7, wherein said capsule polysaccharide is composed of a repeating structure having the formula $[\rightarrow 3)\text{-}\alpha\text{-D-Gal-(1}\rightarrow 2)\text{-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow]_n$.

12. The immunogenic composition of claim 7, wherein said capsule polysaccharide is

13. composed of a repeating structure having the formula $\rightarrow 3)\text{-}\alpha\text{-6-deoxy-D-ido-Heptose (1}\rightarrow 4)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

14. A method of producing anti-*Campylobacter jejuni* immunity comprising the steps:

a. administering the immunogenic composition of claim 1 containing said *C. jejuni* capsule polysaccharide polymer from one or more *Campylobacter jejuni* strains with or without adjuvant at a dose range of 0.1 μ g to 10 mg per dose;

b. administering a boosting dose of said immunogenic composition with or without adjuvant at a dose range of 0.1 μ g to 10 mg per dose.

15. The method of claim 14, wherein said capsule polysaccharide is conjugated to a carrier.

16. The method of claim 14, wherein said carrier is CRM₁₉₇.

17. The method of claim 14, wherein said capsule polysaccharide polymer is composed of a repeating structure consisting of a homopolymer, disaccharide or trisaccharide.

18. The method of claim 14, wherein said capsule polysaccharide is composed of a repeating structure having the formula $[\rightarrow 3)\text{-}\alpha\text{-D-Gal-(1}\rightarrow 2)\text{-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3)\text{-}\beta\text{-D-GlcNAc (1}\rightarrow]_n$.

19. The method of claim 14, wherein said capsule polysaccharide is composed of a repeating structure having the formula $\rightarrow 3)\text{-}\alpha\text{-6-deoxy-D-ido-Heptose (1}\rightarrow 4)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

20. The method of claim 15, wherein said capsule polysaccharide is composed of a repeating structure having the formula $[\rightarrow 3)\text{-}\alpha\text{-D-Gal-(1}\rightarrow 2)\text{-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3)\text{-}\beta\text{-D-GlcNAc (1}\rightarrow]_n$.

21. The method of claim 15, wherein said capsule polysaccharide is composed of a repeating structure having the formula $\rightarrow 3)\text{-}\alpha\text{-6-deoxy-D-ido-Heptose (1}\rightarrow 4)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

22. The method of claim 16, wherein said capsule polysaccharide is composed of a repeating structure having the formula $[\rightarrow 3)\text{-}\alpha\text{-D-Gal-(1}\rightarrow 2)\text{-6d-}\alpha\text{-D-altro-Me-Hep-(1}\rightarrow 3)\text{-}\beta\text{-D-GlcNAc (1}\rightarrow]_n$.

23. The method of claim 16, wherein said capsule polysaccharide is composed of a repeating structure having the formula $\rightarrow 3)\text{-}\alpha\text{-6-deoxy-D-ido-Heptose (1}\rightarrow 4)\text{-}\beta\text{-D-GlcNAc-(1}\rightarrow$.

24. The method of claim 14, wherein said adjuvant is selected from the group consisting of LTR 192G, Aluminum hydroxide, RC529E, QS21, E294, oligodeoxynucleotides (ODN), CpG-containing oligodeoxynucleotides, aluminum phosphate.

* * * * *