Sudden Death in Grower-Finisher Pigs (>20 kg)

If <40 kg each, may include up to 3 pigs per case

Fresh tissues: 1 sample per Whirl-Pak bag, per pig – label with stickers provided.

Formalin-fixed tissues: 1 sample jar per pig

TISSUE SAMPLING LIST - from EACH pig:

Tissue	Fresh (include lesions)	In Formalin (include lesions)
Tonsil	X	X
Thyroid gland		X
Trachea		X
Thymus		X
Lung	X 3	X
Heart		X
Liver	X	X
Spleen	X	X
Kidney	X	X
Adrenal gland		X
Pancreas		X
Mesenteric lymph node	X	X
Gastrohepatic lymph node	X	X
Stomach		X
Duodenum		X
Jejunum		X
lleum		x
Spiral colon		X
Diaphragm		x
Skeletal muscle		x
Brain (or boad)	X half, sagitally sectioned	X half, sagitally sectioned
(or head)	(OR entire head – fresh)	(OR entire head – fresh)



Sudden Death in Grower-Finisher Pigs (>20 kg)

If <40 kg each, may include up to 3 pigs per case

Fresh tissues: Use a separate Whirl-Pak bag for each sample –label bags with the stickers provided (tissue name), and add a pig designation (A, B, C – if >1 pig).

Formalin-fixed tissues: Use 1 sample jar per pig. For solid organs, samples should be approximately 1-2 cm². For intestine, take approximately 3 cm long segments and partially open longitudinally with scissors (to allow formalin to contact the mucosal surface).

Test list (for AHL use):

Virology: PRRSV PCR prtqn, influenza A virus PCR inflpcr: lung (pooled)

Bacterial culture *bsetup*, *cults*: lung, spleen (pooled by tissue).

Histopathology hist

Freeze and hold: tonsil /spleen, lung, kidney/ liver, lymph nodes, brain.



