



Agriculture and
Agri-Food Canada
Research
Branch


Agriculture et
Agroalimentaire Canada
Direction générale
de la recherche

***Diptera types in the
Canadian National
Collection of Insects***

***Part 4
Tachinidae***

***Types de Diptères de la
Collection nationale
des insectes du Canada***

***Quatrième partie
Tachinidae***



Left Blank Page
Page laissée blanche

Diptera types in the Canadian National Collection of Insects

Part 4 Tachinidae

Bruce E. Cooper and James E. O'Hara

Biological Resources Division, Centre for Land and Biological Resources Research, Research Branch, Agriculture and Agri-Food Canada, Ottawa, Ontario K1A OC6

Research Branch
Agriculture and Agri-Food Canada

Publication 1918/B
1996

Types de Diptères de la Collection nationale des insectes du Canada

Quatrième partie Tachinidae

Bruce E. Cooper et James E. O'Hara

Division des ressources biologiques, Centre de recherches sur les terres et les ressources biologiques, Agriculture et agro-alimentaire Canada, Direction générale de la recherche, Ottawa, Ontario K1A OC6

Direction générale de la recherche
Agriculture et agro-alimentaire Canada

Publication 1918/B
1996

©Minister of Supply and Services Canada 1996
Cat. No. A53-1918/1996
ISBN 0-662-62254-5
Printed 1996

Cover illustration
Tachina florum Walker

©Approvisionnements et Services Canada 1996
No de cat.A53-1918/1996
ISBN 0-662-62254-5
Imprimé en 1996

Illustration de la couverture
Tachina florum Walker

CONTENTS**TABLE DES MATIÈRES**

Abstract/ Résumé	7
Introduction	7
Acknowledgments/ Remerciements	9
Tachinidae	11
Literature cited/ Bibliographie	80
Index	87

Left Blank Page
Page laissée blanche

ABSTRACT

The primary type material of the family Tachinidae (Diptera) housed in the Canadian National Collection of Insects, as of 31 December 1994, is catalogued. The Tachinidae collection contains 897 holotypes, 239 associated allotypes, 35 lectotypes (no associated allolectotypes), 2 neotypes, and 117 species that are represented by syntypes. Reference to the original description of each taxon is cited. Label data associated with the specimens are quoted fully, except for paratypes and paralectotypes. A bibliography and an index to named taxa are provided. Certain lectotype fixations, particularly by Mesnil for nominal species described by Villeneuve, are recognized for the first time.

INTRODUCTION

This work is the fourth, and last, in a series of catalogues on the Diptera types in the Canadian National Collection of Insects (CNC). The first, published in 1991, dealt with the Nematocera. The second, published in 1993, documented the types of brachyceran Diptera exclusive of the Schizophora. The third, currently in production, will cover the Schizophora exclusive of the Tachinidae. A brief history of the Diptera collection in the CNC is given in Part 1 of this series.

It is estimated that the Canadian National Collection of Insects houses approximately 250,000 specimens of the family Tachinidae. A large portion of these were collected by past members of the Diptera Unit, particularly A.R. Brooks, C.H. Curran, J.F. McAlpine, G.E. Shewell, J.R. Vockeroth, and D.M. Wood. The collection has been enhanced significantly by the addition of the personal tachinid collections of H.J. Reinhard (acquired 1968; specimens collected mostly from southwestern United States and Mexico), L.P. Mesnil (acquired 1970–1986; Old World specimens, including many types of J. Villeneuve), and W.R. Thompson

RESUME

Ce catalogue répertorie tous les types primaires de la famille des Tachinidae (Diptères) qui étaient conservés dans la Collection nationale canadienne d'insectes au 31 décembre 1994. La collection de Tachinidae comprend 897 holotypes, 239 allotypes associés, 35 lectotypes (aucun allolectotype), deux néotypes et 117 espèces représentées par des syntypes. Pour chaque taxon, nous indiquons la référence à la description originale. Les données inscrites sur les étiquettes de chaque spécimen sont citées intégralement, sauf pour les paratypes et les paralectotypes. Nous incluons une liste bibliographique ainsi qu'un index de tous les taxons mentionnés. Certaines désignations de lectotype, particulièrement celles de Mesnil pour les espèces nominales décrites par Villeneuve, sont reconnues pour la première fois.

INTRODUCTION

Cet ouvrage est le quatrième et le dernier d'une série de catalogues sur les types de diptères dans la Collection nationale canadienne d'insectes (CNC). Le premier, publié en 1991, portait sur les Nématocères. Le second, paru en 1993, documentait les types de Diptères brachycères à l'exclusion des Schizophores. Le troisième, qui englobe les Schizophores hormis les Tachinidae, devrait paraître sous peu. La Partie I de cette série inclut également un bref historique de la collection des diptères de la CNC.

On estime à 250,000 le nombre de spécimens de la famille des Tachinidae retrouvés dans la CNC. Une grande partie de ce matériel fut collectionnée par d'anciens membres de l'Unité des Diptères, dont, en particulier, A.R. Brooks, C.H. Curran, J.F. McAlpine, G.E. Shewell, J.R. Vockeroth et D.M. Wood. L'importance de la collection s'est accrue considérablement suite à l'acquisition des collections personnelles de tachinides de H.J. Reinhard (obtenue en 1968; spécimens provenant surtout du sud-ouest américain et du Mexique), L.P. Mesnil (acquise

(acquired 1969; Trinidad specimens), and through purchase of South American specimens from such private collectors as F. Plaumann and L.E. Peña. Holdings of Tachinidae in the CNC are largest for the Nearctic and Neotropical regions, though all zoogeographic regions of the world are represented.

The number of primary types of Tachinidae in the CNC, as of 31 December 1994, totals 1051. More than 90% of these represent nominal species described by seven authors, namely H.J. Reinhard (267 primary types), L.P. Mesnil (250), J. Villeneuve (178), W.R. Thompson (118), C.H. Curran (91), A.R. Brooks (58), and J.D. Tothill (22).

The format of this work is very similar to that in previous Parts of the series. Entries are arranged alphabetically by genus and species, using the original combinations and spellings. Subgeneric names are included if they were used in the original description of a new species, and are placed in alphabetical order following names without subgenera. Each entry includes the following information: name of taxon and author, year of publication, abbreviated reference (volume number and page where description begins), and current name of country of origin of holotype. Label data associated with each primary type and allotype are cited in full, with diagonal lines separating information given on separate labels (a diagonal line not followed by a space indicates that the diagonal line is present on the label). The number of paratypes or paralectotypes is given along with their country or province/state of origin. The sex of each specimen is given. Notes about nomenclature, genitalic dissections, major damage to primary types, etc., are included under Remarks. For additional information about the names of Mesnil, see "The tachinid taxa of Louis P. Mesnil, with notes on nomenclature (Insecta: Diptera)" (O'Hara, in press). Certain lectotype fixations, particularly by Mesnil for nominal species described by Villeneuve, are recognized for the first time.

entre 1970-1986; matériel de l'ancien monde, incluant plusieurs types de Villeneuve), et W.R. Thompson (acquise en 1969; spécimens de Trinidad). L'achat de spécimens sud-américains de collectionneurs privés tels que F. Plaumann et L.P. Peña a également contribué à rehausser la collection de diptères. La CNC compte la plus importante collection de Tachinidae des régions néarctiques et néotropicales, et toutes les régions zoogéographiques de la terre y sont représentées.

Au 31 décembre 1994, on dénombrait 1051 types primaires de Tachinidae à la CNC. Plus de 90% de ces types représentent des espèces nominales décrites par sept auteurs, soit H.J. Reinhard (267 types primaires), L.P. Mesnil (250), J. Villeneuve (178), W.R. Thompson (118), C.H. Curran (91), A.R. Brooks (58) et J.D. Tothill (22).

La présentation de ce catalogue est analogue à celle des parties précédentes de cette série. On a dressé une liste alphabétique des noms des genres et des espèces, selon leur combinaison et orthographe originales. Les noms des sous-genres employés dans la description originale d'une nouvelle espèce sont également inclus et paraissent par ordre alphabétique à la suite des noms ne comportant pas de sous-genres. Chaque entrée comprend les renseignements suivants: le nom du taxon et de l'auteur, l'année de publication, la référence abrégée de la revue où a paru la description originale (incluant le volume et la page où figurent le nom et la description), ainsi que le nom actuel du pays d'origine de l'holotype. Pour chacun des types primaires et des allotypes, les données des étiquettes correspondantes sont retranscrites intégralement. Une barre oblique sépare les informations provenant de différentes étiquettes (une barre oblique qui n'est pas suivie d'un espace indique que cette barre est présente sur l'étiquette même). On note le nombre et le pays ou province/état d'origine des paratypes ou des paralectotypes. Le sexe de chaque spécimen est indiqué. Des notes sur la nomenclature, les dissections des génitalia, les dommages importants aux types primaires, etc. sont

ACKNOWLEDGMENTS

We would like to thank J.M. Cumming (Centre for Land and Biological Resources Research, Agriculture and Agri-Food Canada), and T.A. Wheeler (Department of Natural Resource Sciences, McGill University, Macdonald Campus) for reviewing the manuscript. We wish to express our sincere appreciation to L. Dumouchel (CLBRR) for the French translation of the introductory sections. We would also like to thank L. Speers (CLBRR) for assistance with the dBase® database upon which this manuscript is based.

ajoutées sous la section "Remarks".

Pour des informations additionnelles sur les noms de Mesnil, voir "The tachinid taxa of Louis P. Mesnil, with notes on nomenclature (Insecta: Diptera)" (O'Hara, sous presse). Certaines désignations de lectotype, particulièrement celles de Mesnil pour les espèces nominales décrites par Villeneuve, sont reconnues pour la première fois.

REMERCIEMENTS

Nous tenons à remercier J.M. Cumming (Centre de recherches sur les terres et les ressources biologiques, Agriculture et Agro-alimentaire Canada), ainsi que T.A. Wheeler ("Department of Natural Resource Sciences", Université McGill, Campus MacDonald) pour la révision critique qu'ils ont faite du manuscrit. Nous exprimons également notre reconnaissance à L. Dumouchel (CRTRB) pour la traduction française des sections d'introduction. Finalement, nous aimerais remercier L. Speers (CRTRB) de l'aide fournie avec la base de données dBase à partir de laquelle cet ouvrage a été élaboré.

Left Blank Page
Page laissée blanche

Acemyia favilla Reinhard, 1974, Can. Ent. 106: 1155. USA.

Holotype ♀: Bondurant, Wyo. VIII-4-1949 L. D. Beamer/ HOLOTYPE *Acemyia favilla* Reinhard/ *Acemyia favilla* R Rnh. Paratypes: 1♀, Illinois; 1♀, South Dakota; 1♀, Ohio.

Acemyia indica Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 183. INDIA.

Holotype ♂: Coll. on wing from paddy nursery/ Lucknow. U.P. June 1963/ 33/ *Acemyia indica* Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Paratype: 1♂, India.

Achaetoneura stilla Reinhard, 1943b, J. Kans. ent. Soc. 16: 17. USA.

Holotype ♂: College Station 9-27.21Texas/ H J Reinhard Collector/ HOLOTYPE *Achaetoneura stilla* Reinhard/ Frontiniella parancilla Tnsd. Det. J. O'Hara 1992.

Allotype ♀: College Station 6.27-17Texas/ H J Reinhard Collector/ ♀/ ALLOTYPE *Achaetoneura stilla* Reinhard/ Frontiniella parancilla Tnsd. Det. J. O'Hara 1992.

Paratypes: 1♂, New York; 1♂, Utah; 1♀, Ohio.

Acroglossa vetula Reinhard, 1964b, Acta zool. mex. 7: 20. MEXICO.

Holotype ♀: Minatitlan, Ver., Mex. 8/26-IX/1/61 R&K Dreisbach/ HOLOTYPE *Acroglossa vetula* Reinhard/ *Acroglossa vetula* R Rnh. Paratypes: 2♀♀, Mexico.

Actia alipes Villeneuve, 1942b, Bull. Soc. ent. Fr. 47: 134. EGYPT.

Holotype ♀: Assuan 44592. II./ *Actia alipes* n.sp./ Peribaea alipes Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ Peribaea palaestina Vill. ♀ B. Herting det.

Actia brunnescens Villeneuve, 1921, Annls Soc. ent. Belg. 61: 46. GERMANY.

Holotype ♂: ♂/ Ndr.Oderwitz 9.5.08 [handwritten word illegible] H.Krammer C./ *Actia brunnescens* Typ. Villen./ Asiphona brunnescens Vill L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Actia brunnipalpis Villeneuve, 1921, Annls Soc. ent. Belg. 61: 45. GERMANY.

Holotype ♂: Harz 4.VII.06/ *Actia brunnipalpis* Typ. Villen./ Asiphona brunneipalpis [sic] Vill. L.P. Mesnil det., 1970/ TYPE/ Asiphona siphonoides Str. L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970/ HOLOTYPE ♂ *Actia brunneipalpis* [sic] Villeneuve det. Stig Andersen/ Ceranthisia ♂ siphonoides (Strobl) det. Stig Andersen 88.

Remarks: Holotype terminalia are stored in glycerine in plastic tubing pinned below specimen.

Actia cibdela Villeneuve, 1913, Revue zool. afr. 3: 35. NIGERIA.

Lectotype ♂: Oshogbe, S.Nigeria. Dr.T.F.G.Mayer, 1-13.X. 1910/ *Actia cibdela* Typ. Villen./ Ceromyia cibdela Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ LECTOTYPE *Actia cibdela* Villeneuve O'Hara designation Selected 1986.

Remarks: For lectotype designation see O'Hara (1989: 54).

Actia comitata Villeneuve, 1936f, Bull. Annls Soc. ent. Belg. 76: 416. UGANDA.

Syntype: 1♂.

♂: Uganda Kampala 3. VI. 1936 H. Hargreaves./ *Actia comitata* Cotyp. Villen. ♂/ *Actia comitata* Villen. L.P. Mesnil det., 1970/ CNC Syntype *Actia comitata* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Actia dasymyia O'Hara, 1991, Can. Ent. 123: 761. CANADA.

Holotype ♂: N.W.T.-21 m.e. Tuktoyaktuk 20-25.VI.71 D.M. Wood/ HOLOTYPE *Actia dasymyia* O'Hara CNC No. 21105.

Allotype ♀: Same data as holotype except with allotype label and collected "2-5.VII.71".

Paratypes: 23♂♂, 14♀♀, Northwest Territories; 4♂♂, 2♀♀, Yukon; 1♀, Colorado.

Actia diffidens Curran, 1933, Am. Mus. Novit. 614: 5. CANADA.

Holotype ♂: *Actia* TYPE ♂ *diffidens* Curran. No./ Kentville,N.S. Jun.17 1924 R.P.Gorham/ HOLOTYPE *Actia diffidens* No 3641 Curran. Allotype ♀: St.Peters,N.S. 10-VII-1930 M.L.Preble/ 195/ *Actia* TYPE *diffidens* Curran. ♀ No./ ALLOTYPE *Actia diffidens* No. 3641 Curran.

Actia dimorpha O'Hara, 1991, Can. Ent. 123: 755. USA.

Holotype ♂: USA:GA.McIntoshCo. Sapelo Island 27.VIII-7.IX.1987 BRC HYM TEAM, MT shrubs, sand dunes/ HOLOTYPE *Actia dimorpha* O'Hara CNC No. 21106.

Allotype ♀: USA:GA,McIntoshCo. Sapelo Island 9-21.IX .1987 BRC HYM. TEAM Savannah, FIT/ ALLOTYPE *Actia dimorpha* O'Hara CNC No. 21106.

Paratypes: 22♂♂, 2♀♀, Georgia; 2♂♂, 2♀♀, South Carolina; 1♂, Florida; 5♂♂, 15♀♀, Ontario.

Actia exsecta Villeneuve, 1936f, Bull. Annls Soc. ent. Belg. 76: 416. UGANDA.

Syntype: 1♂.

♂: Uganda Kampala 3. VI. 1936 H. Hargreaves/ *Actia exsecta* Cotyp. Villen. ♂/ *Actia exsecta* Villen. L.P. Mesnil det., 1970/ CNC Syntype *Actia exsecta* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Actia flaviseta Villeneuve, 1921, Annls Soc. ent. Belg. 61: 45. GERMANY.

Lectotype ♀: Samara 57934/ *Actia flaviseta* Typ. Villen. Ceromyia flaviseta Villen. L.P. Mesnil det., 1970/ CNC Syntype *Actia flaviseta* Villeneuve Label affixed 1994/ Lectotype fixation

- by Mesnil, 1963: 836. Label affixed 1995/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Paratype: 1♂, Germany.
- Remarks: We regard Mesnil's (1963b: 836) reference to a holotype as a lectotype fixation for this species. Lectotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.
- Actia jocosa* Villeneuve, 1942b, Bull. Soc. ent. Fr. 47: 134.
ALGERIA.
Holotype ♂: Algier IV. 52286/ *Actia jocosa* n.sp./ Ceranthis jocosa Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Actia jocularis* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 47.
JAPAN.
Holotype ♂: 23-IX-1950 TOKURA H.HASEGAWA/ 296/ *Actia jocularis* Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Actia languidula* Villeneuve, 1913, Revue zool. afr. 3: 36.
NIGERIA.
Syntype: 1♂.
♂: Oshogbo, S.Nigeria. Dr.T.F.G.Mayer, 1-15 X 1910/ *Actia languidula* Typ. Villen/ Ceromyia languidula Villen. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype *Actia languidula* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Actia nigrohalterata* Villeneuve, 1921, Annls Soc. ent. Belg. 61: 45. DENMARK.
Holotype ♂: Danemark/ How 12.5.12/ *Actia nigrohalterata* Typ. Villen./ Ceromyia nigrohalterata Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ HOLOTYPE ♂ *Actia nigrohalterata* Villeneuve 1921 det. Stig Andersen/ Entomophaga ♂ nigrohalterata (Vill.) det. Stig Andersen.
- Actia ontario* Curran, 1933, Am. Mus. Novit. 614: 4. CANADA.
Holotype ♀: *Actia* TYPE *ontario* ♀ Curran No./ Norway Point Lake of Bays,Ont. 31. VII. 1919 J.McDunnough/ HOLOTYPE *Actia* *ontario* No.Curran 3640/ ♀.
- Actia radialis* O'Hara, 1991, Can. Ent. 123: 763. CANADA.
Holotype ♂: Gatineau Park,QUE. 28.VII.1967 W.G.Matthewman K-391 Ex Fomes fomentarius/ 68-143/ A47/ HOLOTYPE *Actia radialis* O'Hara CNC No. 21107.
Paratype: 1♂, Ontario.
Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.
- Actia samarensis* Villeneuve, 1921, Annls Soc. ent. Belg. 61: 46.
RUSSIA.
Holotype ♀: Samara 57934/ *Actia samarensis* Villen./ Asiphona samarensis Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Actia starkei* Mesnil, 1952c, Bull. Annls Soc. ent. Belg. 88: 155.
GERMANY.
- Holotype ♂: ♂/ *Gymnop. versiwt* Gr. Hennersdt. 31. 5.02./ *Asiphona starkei* Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Remarks: Two females in the CNC may be paratypes, but they are not labeled as such.
- Actia sternalis* O'Hara, 1991, Can. Ent. 123: 765. CANADA.
Holotype ♂: Mi. 51, Y.T. Dempster Hwy. 18-27.VII.1973 G.& D.M.Wood/ HOLOTYPE *Actia sternalis* O'Hara CNC No. 21108.
Paratypes: 6♂♂, Yukon.
- Actia sufferta* Villeneuve, 1942b, Bull. Soc. ent. Fr. 47: 133.
GERMANY.
Holotype ♂: *Thryptocera* latifrons Mg/ *Actia sufferta* n.sp./ Ceromyia sufferta Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ HOLOTYPE ♂ *Actia sufferta* Villeneuve 1942 det. Stig Andersen/ Entomophaga ♂ sufferta (Vill.) det. Stig Andersen.
- Actia tenuipalpis* Villeneuve, 1921, Annls Soc. ent. Belg. 61: 46.
RUSSIA, GERMANY.
Syntypes: 2♂♂.
♂: Zlatoust Vi.2409 J. Schnabl/ *tenuipalpis* Typ. Villen./ Ceranthis tenuipalpis Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ LECTOTYPE ♂ *Actia tenuipalpis* Villeneuve det. Stig Andersen 1988/ Ceranthis ♂ tenuipalpis (Vill.) det. Stig Andersen/ CNC Syntype *Actia tenuipalpis* Villeneuve Label affixed 1994.
♂: Berlin Pichelsberg 16. 6. 19/ Ceranthis tenuipalpis Vill. L.P. Mesnil det., 1970/ CNC Syntype *Actia tenuipalpis* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: Terminalia of Zlatoust syntype are stored in glycerine in plastic tube pinned below specimen. Lectotype designation has not been published.
- Actia (Euchaetactia) montium* Villeneuve, 1921, Annls Soc. ent. Belg. 61: 47. FRANCE.
Holotype ♂: Lautaret 57053./ Euchaetactia montium Typ. Villen./ TYPE/ HOLOTYPE ♂ *Goniocera (Euchaetactia) montium* Villeneuve det. Stig Andersen/ EX L.-P. MESNIL COLLECTION 1970/ *Goniocera montium* Villen. L.P. Mesnil det., 1970.
Remarks: Holotype terminalia are stored in glycerine in plastic tubing pinned below specimen.
- Actinochaetopteryx japonica* Mesnil, 1970, Mushi 44: 117. JAPAN.
Holotype ♀: Nukabira Hokkaido 16. VII. 1963 S. Takano/ 376/ *Actinochaetopteryx japonica* Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Actinodoria argentata* Reinhard, 1974, Can. Ent. 106: 1156.
MEXICO.
Holotype ♀: Jalapa, Ver. Mex. VIII/1-6/61 R&K Dreisbach/ HOLOTYPE *Actinodoria argentata* Rnh. CNC No.

- Actinodoria argentea* Thompson, 1964, Studia ent. 7: 126.
TRINIDAD.
Holotype ♂: HOLOTYPE *Actinodoria argentea* Thomp. CNC No. 9821/ Melajo R. III.18.60
Trinidad,BWI./ *Actinodoria argentea* n.sp./ HOLOTYPE CNCNo.
Allotype ♀: ALLOTYPE *Actinodoria argentea* Thomp. CNC No. 9821/ Arima V. TRINIDAD W.I. III. 9.
1961 W.R.Thompson/ 1169 III.10.61/ *Actinodoria*.
Paratypes: 2♂♂, 1♀, Trinidad.
- Admontia badiceps* Reinhard, 1958d, J. Kans. ent. Soc. 31: 279.
USA.
Holotype ♂: Puyallup, Wn. [Washington] III-23.1933
WmWBaker,Coll/ HOLOTYPE *Admontia* *badiceps* Reinhard/ *Admontia badiceps* Rnh.
Allotype ♀: Puyallup, Wn. IV-25. 1935 J. Wilcox,
Coll./ ALLOTYPE *Admontia badiceps* Reinhard.
- Admontia dubia* Curran, 1927b, Can. Ent. 59: 297. CANADA.
Holotype ♀: Holdfast Sask. 7.8.1925 Kenneth M.King/
Admontia TYPE *dubia* ♀ Curran No. 2469.
- Admontia nigrita* Thompson, 1968, Mem. ent. Soc. Can. 56: 123.
TRINIDAD.
Holotype ♀: HOLOTYPE *Admontia nigrita* Thomp. CNC No. 9869/ Simla 1964 Trinidad, WI./
Admontia.
Paratype: 1♀, Trinidad.
- Admontia pollinosa* Curran, 1927b, Can. Ent. 59: 296. CANADA.
Holotype ♂: *Admontia* TYPE *pollinosa* Curran No.
2470/ Man. Binscarth H.A.Robertson 9 VIII 1923.
Allotype ♀: *Admontia* ALLOTYPE *pollinosa* ♀ Curran
No. 2470/ Aweme, Man. N. Criddle 7 X 1924.
Paratype: 1♀, South Dakota.
- Admontia rufochaeta* Curran, 1927b, Can. Ent. 59: 296. USA.
Holotype ♂: GreenBayWis NFHoward 7-5-15/
Admontia TYPE *rufochaeta* Curran No. 2468.
- Adoryphorophaga luxatura* Reinhard, 1974, Can. Ent. 106: 1156.
MEXICO.
Holotype ♂: K107, Pueb. Sept 5,47/ F. A. Cowan M. R.
Wheeler Collectors/ HOLOTYPE
Adoryphorophaga luxatura Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratype: 1♀, Mexico.
- Adoryphorophaga secutoris* Reinhard, 1974, Can. Ent. 106: 1157.
MEXICO.
Holotype ♀: Jalapa, Ver., Mex. VIII/1-6/61 R&K
Dreisbach/ HOLOTYPE *Adoryphorophaga*
secutoris Reinhard/ *Adoryphorophaga secutoris* R
Rnh.
- Akosempomyia caudata* Villeneuve, 1932a, Bull. Annls Soc. ent.
Belg. 71 (1931): 244. TAIWAN.
Syntype: 1♂.
♂: Formosa, Toyenmongai/ *Akosempomyia*
caudata, Type Villen.
- Allophorocera lapponica* Wood, 1974b, Can. Ent. 106: 670.
FINLAND.
Holotype ♂: Saana,Kilpisjärvi FINLAND 900m.
16-17.VII.1969 G.C.&D.M.Wood/ HOLOTYPE
Allophorocera lapponica Wood CNC No. 13245.
Allotype ♀: Same data as holotype except with allotype
label, and additional label "IDEMA illustration".
- Alloprosopaea efflatouni* Villeneuve, 1923a, Bull. Soc. r. ent.
Égypte 7 (1922): 90. EGYPT.
Holotype ♀: Dashor 28.4.21/ Dashor Egypt 28. 4. 1921/
Alloprosopaea efflatouni Villen. L.P. Mesnil det.,
1970/ TYPE.
- Alophorellopsis argentifrons* Brooks, 1945a, Scient. Agric. 25: 675.
CANADA.
Holotype ♂: Oliver,B.C. 28 V.1923 C.B.Garrett/
HoloTYPE ♂ *Alophorellopsis argentifrons* No.
Brks 5592.
Remarks: Date of collection of holotype was published
as 28 June 1923.
- Alsomyia rufipes* Villeneuve, 1937e, Bull. Annls Soc. ent. Belg. 77:
407. INDIA.
Lectotype ♂: LECTO-TYPE/ Coimbatore Parasite on
Clania crameri 7.IX.13. T.B.F. Coll/ TYPE/
Nealsomyia rufipes Vill. L.P. Mesnil det., 1964/
Alsomyia rufipes Villeneuve LECTOTYPE ♂
designated by R.W.Crosskey 1971.
Remarks: For lectotype designation see Crosskey
(1976: 264).
- Amazohoughia flavipes* Thompson, 1964, Studia ent. 7: 120.
TRINIDAD.
Holotype ♀: HOLOTYPE *Amazohoughia flavipes*
Thomp. CNC No. 9820/ I.C.T.A. Trinidad,BWI.
DEC-1-1952./ Collector F.J.Simmonds/
Amazohoughia flavipes n.sp.
Allotype ♂: ALLOTYPE *Amazohoughia flavipes*
Thomp. CNC No. 9820/ U. Manzanilla I.26.61.
Trinidad,BWI.
Paratype: 1♀, Trinidad.
- Amesioclea cincta* Villeneuve, 1936a, Bull. Mus. r. Hist. nat. Belg.
12(4): 1. SOUTH AFRICA.
Syntype: 1♂.
♂: Wint-hoek [Windhoek] Tulbagh 3600 f/
Amesioclea cincta Type n.sp./ *Amesioclea cincta*
Villen. L.P. Mesnil det., 1970/ TYPE/ Prodexilla
posio Walker L.P. Mesnil det., 1970/ CNC
Syntype *Amesioclea cincta* Villeneuve Label
affixed 1994/ EX L.-P. MESNIL COLLECTION
1970.
- Amphicestonia dispar* Villeneuve, 1939a, Bull. Annls Soc. ent.
Belg. 79: 349. TUNISIA.
Holotype ♂: Kroumirie/ *Amphicestonia dispar* Type
Villen./ TYPE/ EX L.-P. MESNIL COLLECTION
1970.
- Anacamptomyia pallida* var. *obscurella* Mesnil, 1950a, Revue Zool.
Bot. afr. 43: 24. ZAIRE.
Syntype: 1♂.

- ♂:* Congo - beige Eala III -1936 J.Ghesquière 2289/ Parasite de Ropalia/ R. Mus. Hist. Nat. Belg. No 4 8 2/ Anacamptomyia obscurella Mesn. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Anacamptomyia pallida var. obscurella Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Remarks: Holotype right hind leg is glued to double mount strip. The CNC also has two males (one with same data as specimen labeled as type except collected on 14 April 1936, the other with only a male symbol) and one female (one label with female symbol), which may be syntypes.
- Anaeogmena setinervia* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 15. JAPAN.
- Holotype ♀: V-9-1953 Ryukyu Is. T. SHIRAKI/ 291/ Aneogmena [sic] setinervia Mesnil L.P. Mesnil det., 1969/ TYPE/ HOLOTYPE ♀ of *Anaeogmena setinervia* Mesnil Examined 1970 R.W.Crosskey.
- Remarks: Date of collection of holotype was published as 3 May 1953.
- Angustiopsis saginata* Reinhard, 1959a, Ent. News 70: 231. USA.
- Holotype ♂: Presidio Co. VII-13-40 Tx./ HOLOTYPE Angustiopsis saginata Reinhard.
- Allotype ♀: Same data as holotype except with allotype label.
- Paratype: 1♀, Texas.
- Anoxynops aurifrons* Thompson, 1968, Mem. ent. Soc. Can. 56: 57. TRINIDAD.
- Holotype ♂: HOLOTYPE Anoxynops aurifrons Thomp. CNC No. 9852/ Santa Cruz V 2.27.60 Trinidad,BWI./ 1411 II.27.64.
- Anoxynopoidea grisea* Thompson, 1968, Mem. ent. Soc. Can. 56: 61. TRINIDAD.
- Holotype ♂: I.C.T.A. Trinidad,BWI. FEB.8.1954/ Collector F.J.Simmonds/ 62/ F/ HOLOTYPE Anoxynopoidea grisea Thomp. CNC No. 9853.
- Allotype ♀: ALLOTYPE Anoxynopoidea grisea Thomp. CNC No. 9853/ MOUNT HOPE Trinidad,BWI. JAN.20.1953/ Collector F.J.Simmonds/ 62/ 1010.2 V.26.59.
- Anthomyiopsis plagioderae* Mesnil, 1972, Fliegen palaearkt. Reg. 64g: 1109. SWITZERLAND.
- Holotype ♂: Feldmeilen 12. 5. 1954/ Anthomyiopsis plagioderae Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Paratype: 1♀, Switzerland.
- Antillicola trinitatis* Thompson, 1963a, Can. J. Zool. 41: 422. TRINIDAD.
- Syntype ♀: HOLOTYPE Antillicola trinitatis Thomp. CNC No. 9778/ Brazil II.2.61 Trinidad,BWI/ Antillicola trinitatis n.sp./ CNC Syntype Antillicola trinitatis Thompson Label affixed 1994.
- Antistasea fimbriata* Bischof, 1904, Verh. zool.-bot. Ges. Wien 54: 83. SOUTH AFRICA.
- Holotype ♂: Algoa bay Capland Dr.Brauns 11 11 95/ Antistasea fimbriata Typ. Bisch./ Antistasea fimbriata Bisch. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Antistasea mutans* Mesnil, 1970, Mushi 44: 106. SOUTH AFRICA.
- Holotype ♂: Mfongosi Zulu L. W E Jones/ Antistasea mutans Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Anurophylla setosa* Villeneuve, 1938a, Bull. Annls Soc. ent. Belg. 78: 413. ALGERIA.
- Holotype ♂: Anurophylla setosa Type Villen./ La Calle algérie ♂/ A. setosa Vill. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Aphantorhapha atoma* Reinhard, 1947, J. Kans. ent. Soc. 20: 19. USA.
- Holotype ♂: College Station Apr.18,1943 Tex/ H J Reinhard Collector/ HOLOTYPE Aphantorhapha atoma Reinhard.
- Allotype ♀: Same data as holotype except with allotype label.
- Paratypes: 23♂♂, 5♀♀, Texas.
- Aphantorhapha pulla* Reinhard, 1974, Can. Ent. 106: 1157. MEXICO.
- Holotype ♀: Cuernavaca Mor., Mex. 7-29-61 7000' R&K Dreisbach/ HOLOTYPE Aphantorhapha pulla Reinhard/ Aphantorhapha pulla R Rnh.
- Aphelogaster (Aphelogaster) compressa* Mesnil, 1974, Fliegen palaearkt. Reg. 64g: 1291. CHINA.
- Holotype ♂: Mandeschurei Charbin 7.1952 V. Alin leg.,vend. Eing. Nr. 6, 1953/ Aphelogaster compressa Mesn. L.P. Mesnil det., 1973/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Aphria klapperichi* Mesnil, 1967. Mushi 41: 49. CHINA.
- Holotype ♂: Shaowu - Fukien (500m) J.Klapperich 29.6 .1937/ Aphria klapperichi Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Aphria (Plagiopsis) rubida* Mesnil, 1973a, Fliegen palaearkt. Reg. 64g: 1153. TUNISIA.
- Holotype ♂: TUNISIE BOU HEDMA C. DUMONT 1929/ APRIL/ Aphria rubida Mesnil L.P. Mesnil det., 1971/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Aplomyia latimana* Villeneuve, 1934a, Revue Zool. Bot. afr. 25: 409. UGANDA.
- Holotype ♀: R. [Ruwenzori] 1800m 11.IV.14/ tarses ant. dilatés/ Aplomyia latimana Typ. Villen./ Aplomyia latimana Villen. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Aplomyiopsis angustipennis* Reinhard, 1964a, J. Kans. ent. Soc. 37: 42. USA.
- Holotype ♂: Amherst, Ohio 6-20 1939/ H J Reinhard Collector/ HOLOTYPE Aplomyiopsis angustipennis Reinhard/ Aplomyiopsis angustipennis R Rnh.

Paratype: 1♂, Ohio.

Aporotachina latifrons Jacentkovski, 1944, Sb. ent. Odd. nár. Mus. Praze 21-22: 394. CZECH REPUBLIC.
Holotype ♂: Lovice 6.VI.42/ Aporotachina latifrons n.sp./ TYPE/ Aporotachina latifrons Jacent. L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970.

Archiclops carthaginiensis mageritensis Mesnil, 1956a, Fliegen palaearkt. Reg. 64g: 510. SPAIN.
Holotype ♂: El Pardo J. GIL/ Archiclops mageritensis Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Archytas bennetti Thompson, 1963a, Can. J. Zool. 41: 366.
TRINIDAD.
Holotype ♀: HOLOTYPE Archytas bennetti Thomp. CNC No. 9768/ Simla TRINIDAD W.I. III.16.61 W.R.Thompson/ Archytas bennetti n.sp/ species H./ HOLOTYPE CNCNo./ 1175 III.17.61.
Remarks: Date of collection of holotype was published as 17 March 1961.

Archytas caroniensis Thompson, 1963a, Can. J. Zool. 41: 380.
TRINIDAD.
Syntypes: 1♂, 1♀.
♂: 2. 17. 60 Brazil Trinidad,BWI./ Type ♂ [underside of label] genit. ♂ Type II/ CNC Syntype Archytas caroniensis Thompson Label affixed 1994.
♀: HOLOTYPE Archytas caroniensis Thomp. CNC No. 9772/ 2. 17. 60 Brazil Trinidad,BWI./ 1126 IX.22.60/ allotype ♀/ CNC Syntype Archytas caroniensis Thompson Label affixed 1994.

Archytas dissimiloides Thompson, 1963a, Can. J. Zool. 41: 369.
TRINIDAD.
Holotype ♀: HOLOTYPE Archytas dissimiloides Thomp. CNC No. 9770/ Archytas D Det. n.sp./ 1007 ♀./ Archytas dissimiloides n.sp/ Collector F.J.Simmonds/ MARACAS V. Trinidad,BWI. NOV.28.1952.

Archytas hiemalis Thompson, 1963a, Can. J. Zool. 41: 368.
TRINIDAD.
Syntype: 1♀.
♀: HOLOTYPE Archytas hiemalis Thomp. CNC No. 9769/ SANTA CRUZ V. 3. 5. 60 Trinidad,BWI./ 1076 III.4.60/ Archytas ♀/ Archytas hiemalis n.sp./ CNC Syntype Archytas hiemalis Thompson Label affixed 1994.

Archytas instabilis Curran, 1928b, Can. Ent. 60: 224. USA.
Holotype ♂: Archytas TYPE instabilis ♂ Curran No.2828/ Hewitt, N.J. VI.18.1904./ Wm.T.Davis Collection.

Archytas nivalis Curran, 1928b, Can. Ent. 60: 254. USA.
Holotype ♂: Washington, D.C. X.29.26 N.K.Bigelow/ Archytas TYPE nivalis ♂ Curran No.2823.
Allotype ♀: Great Falls Va./ Wm.T.Davis July 15,1913/ SLIDE OS64 No./ Archytas AlloTYPE nivalis ♀ Curran No.2823/ Slide No. 0015 FWR.

Archytas piarconensis Thompson, 1963a, Can. J. Zool. 41: 387.
TRINIDAD.

Syntype ♀.
♀: HOLOTYPE Archytas piarconensis Thomp. CNC No. 9774/ PIARCO Trinidad,BWI. OCT.27.1952/ Collector F.J.Simmonds/ 1180C III.29.61./ CNC Syntype Archytas piarconensis Thompson Label affixed 1994.

Archytas purseglovei Thompson, 1963a, Can. J. Zool. 41: 364.
TRINIDAD.

Syntypes: 1♂, 1♀.
♂: HOLOTYPE Archytas purseglovei Thomp. CNC No.9767/ MARACAS V Trinidad,BWI. NOV.12-1952/ Collector F.J.Simmonds/ HOLOTYPE CNCNo./ allotype ♂/ CNC Syntype Archytas purseglovei Thompson Label affixed 1994.
♀: ALLOTYPE CNCNo./ Type ♀/ S. Cruz V. IV.8.60/ 1108 IV.8.60/ CNC Syntype Archytas purseglovei Thompson Label affixed 1994.

Archytas rufiventris Curran, 1928b, Can. Ent. 60: 280. USA.
Holotype ♂: Archytas TYPE rufiventris Curran No./ Royal Palm Park Fla. Apr. 12-18, 1923 F 4671.S.

Allotype ♀: Miami 12.XI Fla/ CHTTownsend Collector/ Archytas AlloTYPE ♀ rufiventris Curran No.
Paratype: 1♂, Florida.

Archytas sanctaecrucis Thompson, 1963a, Can. J. Zool. 41: 386.
TRINIDAD.

Syntype: 1♀.
♀: HOLOTYPE Archytas sanctaecrucis Thomp. CNC No. 9773/ Santa Cruz V. 3.5.60 Trinidad,BWI./ 1080 III.7.60/ Archytas ♀ sanctaecrucis/ CNC Syntype Archytas sanctaecrucis Thompson Label affixed 1994.

Archytas trinitatis Thompson, 1963a, Can. J. Zool. 41: 374.
TRINIDAD.

Syntype: 1♂.
♂: HOLOTYPE Archytas trinitatis Thomp. CNC No. 9771/ St. Augustine, Trinidad,BWI. OCT.1 1952./ Collector F.D. Bennett/ Type ♂. [underside of label] VII./ CNC Syntype Archytas trinitatis Thompson Label affixed 1994.

Archytas vulgaris Curran, 1928b, Can. Ent. 60: 276. USA.

Holotype ♂: HoloTYPE ♂ Archytas vulgaris 1434. No.Curran/ C.H. Curran. Lawrence,Ks. 17.VI.1922.

Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 22♂♂, 23♀♀, Kansas; 2♂♂, 1♀, Mississippi; 1♂, Tennessee.

Arctophyo algens Curran, 1926c, Can. Ent. 58: 215. CANADA.

Holotype ♂: Arctophyo HoloTYPE ♂ algens Curran. No. 1944/ LowBush Ont. Lake Abitibi VII-19-1925 N.K.Bigelow.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 15♂♂, 28♀♀, Ontario.

Arctophyto marginalis Curran, 1924d, Can. Ent. 56: 302.

CANADA.

Holotype ♂: HoloTYPE Arctophyto marginalis Curr No. 750/ Revelstoke Mt., B.C. Elevation 6000 ft. 12.VIII.1923 E.R.Buckell.

Allotype ♀: Same data as holotype except with allotype label and collected by P.N. Vroom.

Paratypes: 1♂, 1♀, Alberta; 1♂, British Columbia.

Argentoepalpus rufipes Brooks, 1949, Can. Ent. 81: 22. CANADA.

Holotype ♂: Robson, B.C. 7.IV.1947 H.R. Foxlee/ Argentoepalpus Holo TYPE ♂ rufipes Brks No. 5771.

Allotype ♀: Same data as holotype except with allotype label and collected "21.IV.1947".

Paratypes: 6♂♂, 8♀♀, British Columbia.

Argyrophylax atricauda Mesnil, 1967, Mushi 41: 38. JAPAN.

Holotype ♂: Unzen Kyushu 31. V. 1964 S.Takano/ 411/ Argyrophylax atricauda Mesn. L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Argyrophylax bisetosa Thompson, 1963b, Studia ent. 6: 351.

TRINIDAD.

Syntypes: 4♀♀.

♀: HOLOTYPE Argyrophylax bisetosa Thomp. CNC No. 9792/ MARACAS V. Trinidad, BWI. NOV.7.1952/ Collector F.J.Simmonds/

Argyrophylax bisetosa n sp./ CNC Syntype

Argyrophylax bisetosa Thompson Label affixed 1994.

♀: PARATYPE Argyrophylax bisetosa Thomp. CNC No. 9792/ I.C.T.A. Trinidad, BWI.

OCT.23.1952/ Collector F.J.Simmonds/

Argyrophylax Det. ♀ ♀/ CNC Syntype

Argyrophylax bisetosa Thompson Label affixed 1994.

♀: PARATYPE Argyrophylax bisetosa Thomp. CNC No. 9792/ I.C.T.A. Trinidad, BWI. MARCH.

1954./ Collector F.J.Simmonds/ 121/ CNC

Syntype Argyrophylax bisetosa Thompson Label affixed 1994.

♀: PARATYPE Argyrophylax bisetosa Thomp. CNC No. 9792/ Valaqr. XI.25.60/ Trinidad, BWI.

1231 VII.20.62/ CNC Syntype Argyrophylax bisetosa Thompson Label affixed 1994.

Argyrophylax nova novella Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 14. JAPAN.

Holotype ♂: Obihiro S.Takano/ 260/ Argyrophylax novella Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Argyrophylax triangulifera Thompson, 1963b, Studia ent. 6: 352.

TRINIDAD.

Syntype: 1♀.

♀: HOLOTYPE Argyrophylax triangulifera Thomp. CNC No. 9793/ ARIPO V. Trinidad, BWI.

FEB.11.1954/ COLLECTOR F.D.BENNETT/

Hind legs missing/ CAR/ 1229 V.25.62/

Argyrophylax triangulifera nsp./ CNC Syntype

Argyrophylax triangulifera Thompson Label affixed 1994.

Arrhinomyia nudicosta Mesnil, 1970, Mushi 44: 116. JAPAN.

Holotype ♂: Oshitata Spa Mt. Bandai 18.VII.1966 S.Takano/ 418/ Arrhinomyia nudicosta Mesnil L.P. Mesnil det., 1969/ TYPE.

Atactosturmia vittata Thompson, 1963b, Studia ent. 6: 381.

TRINIDAD.

Holotype ♀: HOLOTYPE Atactosturmia vittata Thomp. CNC No. 9796/ St. Augustine, Trinidad, BWI. OCT.31.1952/ Collector F.J.Simmonds/ 91/ Sp. 11 11/ Atactosturmia vittata n sp.

Athanatus knowltoni Reinhard, 1947, J. Kans. ent. Soc. 20: 16.

USA.

Holotype ♂: S.L. Wood Logan Dry Cany. Ut. June 18 1944/ HOLOTYPE Athanatus knowltoni Reinhard/ Athanatus knowltoni R. Rnh.

Atylomyia albifrons Villeneuve, 1911, Wien. ent. Ztg 30: 86.

EGYPT.

Lectotype ♀: 13/1 25/2 1910/ Helouan (Egypte)/ Atylomyia ♀ argentifrons type Villen./ Atylomyia argentifrons Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Atylomyia albifrons Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1962: 776. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Although the label reads *argentifrons*, the name was published as *albifrons*. We regard Mesnil's (1962: 776) reference to a holotype as a lectotype fixation for this species.

Atylomyia rungsi Mesnil, 1962, Fliegen palaearkt. Reg. 64g: 778. MOROCCO.

Holotype ♂: S.Maroc Ait Melloul 10.XI.1948 Ch. Rungs/ Atylomyia rungsi Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Aubaeanetia anorbitalis Brooks, 1945b, Can. Ent. 77: 96.

CANADA.

Holotype ♂: Holotype ♂ Aubaeanetia anorbitalis No. Brks 5522/ Aylmer, Que. 18. VII. 1924 C.H.Curran/ No fronto-orbitalis is this not Paradexodes albifacies Tns./ Has Aubaeanettia [sic] spinules below squamae as in Paradexodes?/ Aubaeanettia assimilis A. Tns.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 1♂, Quebec; 1♀, Maryland.

Aulicomyia invulnerata Reinhard, 1943c, Bull. Brooklyn ent. Soc. 38: 81. USA.

Holotype ♀: Amherst, Ohio July, 1933/ H J Reinhard Collector/ HOLOTYPE Aulicomyia invulnerata Reinhard/ Aulicomyia invulnerata R. Rnh/ Neothelaira aurifrons R. Coq./ Aporotachina aurifrons R (Coq.).

Paratype: 1♀, Ohio.

Bactromyia dasyops Mesnil, 1969, Entomophaga 13: 204. NEW CALEDONIA.

Holotype ♂: Noumea Aux Vata XI. 1964/ Ex Lepidopt. Terminalia catappa L. [underside of

label] Combretacée identif. B.M./ Bactromyia dasyops Mesn. L.P. Mesnil det., 1966/ TYPE/ HOLOTYPE ♂ of Bactromyia dasyops Mesnil Examined 1970 R.W.Crosskey/ Genus PARADRINO Mesn. R.W.Crosskey det. 1972/ Paradrino laevicula? det. H. Shima, 1988.
Remarks: Holotype puparium is glued to a paper strip below specimen.

Bactromyia delicatula Mesnil, 1953a, Fliegen palaearkt. Reg. 64g: 265. TAIWAN.
Holotype ♂: Kankau (Koshun) Formosa H.Sauter 1912/ 7.VII./ Bactromyia delicatula Mesn. L.P. Mesnil det., 1952/ TYPE/ HOLOTYPE ♂ of Bactromyia delicatula Mesnil Examined 1970 R.W.Crosskey.
Remarks: Holotype is missing terminalia.

Baeomyia juniperi O'Hara, 1984, Can. J. Zool. 62: 1395. CANADA.
Holotype ♂: Williams Lake, B.C. Em. 21-V-58 F.I.S./ 57-7276-03 Ex. Semiothisa triviata/ 63-145 192/ 11/ HOLOTYPE Baeomyia juniperi ♂ O'Hara.
Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.
Holotype puparium is pinned below specimen.

Baeomyia sonorensis O'Hara, 1984, Can. J. Zool. 62: 1393. USA.
Holotype ♂: USA Ariz. Graham Co. 2.4km. W. on Hwy 366 from Hwy. 666 1160m. 7-9.VIII.1983 J.E. O'Hara/ HOLOTYPE Baeomyia sonorensis ♂ O'Hara.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 7♂♂, 2♀♀, Arizona.

Baeomyia xanthogaster O'Hara, 1984, Can. J. Zool. 62: 1394. CANADA.
Holotype ♂: No 50.2985B Date 28.IV F.I.S. 1951/ S. granitata Vermillion [sic] B.C./ HOLOTYPE Baeomyia xanthogaster ♂ O'Hara.
Allotype ♀: Bridesville B.C. 27.VII.1949. F.I.S./ BC48-3753/ Host S. granitata/ ALLOTYPE Baeomyia xanthogaster ♀ O'Hara.
Paratypes: 12♂♂, 11♀♀, British Columbia.
Remarks: Holotype puparium is pinned below specimen.

Baumhaueria (Aguilarina) microps Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 18. YUGOSLAVIA.
Holotype ♂: Zemun Yugoslavia 9.4.1960/ Baumhaueria microps Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Bessa obliquata Mesnil, 1944b, Revue fr. Ent. 11: 16. GUINEA.
Holotype ♂: Guinée Haut 07/ Bessiola obliquata Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Besseria excavata Herting, 1979, Stuttg. Beitr. Naturk. (A) 323: 8. MADAGASCAR.
Holotype ♂: Madagascar, Tananarive, Lambertson 1912/ Besseria fossulata Bezzi/ Besseria excavata n. sp., B. Herting det., Type.

Besseria incompleta Curran, 1926d, Can. Ent. 58: 84. CANADA.
Holotype ♀: Macdiarmid Ont. VI-27-22 NKBIGELOW/ Besseria TYPE incompleta Curran No. 1869./ Besseria anthophila Loew B. Herting det./ Besseria incompleta Det. Curran. C.H.Curran.

Billaea agrianomei Mesnil, 1969, Entomophaga 13: 206. NEW CALEDONIA.
Lectotype ♂: Yahou New Caledonia 4.XII.1965/ Ex Agrionome fairmairei Cerambycidae/ Billaea agrianomei Mesn. L.P. Mesnil det., 1966/ EX L.-P. MESNIL COLLECTION 1970/ LECTOTYPE ♂ *Billaea agrianomei* Designated by D.A. Barraclough.
Remarks: For lectotype designation see Barraclough (1992: 1225). See comments by O'Hara (in press).

Billaea kolomyetzi Mesnil, 1970, Mushi 44: 121. POLAND.
Holotype ♂: Bialowieska Od.407.MTodn.
Pin.hypn.qnezc. 23.VII. 1938 J. Pawlowicz/
Billaea kolomyetzi Mesn L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Billaea latimargo Villeneuve, 1937b, Bull. Mus. r. Hist. nat. Belg. 13(13): 2. "LAPLAND".
Holotype ♀: Billaea latimargo ♀ typ. Villen./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Billaea nipigonensis Curran, 1926d, Can. Ent. 58: 89. CANADA.
Holotype ♂: Billaea HoloTYPE ♂ nipigonensis Curran No. 1794/ Macdiarmid Ont. VII-13-22 NK BIGELOW.

Billaea ringdahli Villeneuve, 1937b, Bull. Mus. r. Hist. nat. Belg. 13(13): 2. SWEDEN.
Syntype: 1♂.
♂: Visby 21.7-36/ (Suède)/ Billaea Ringdahli Typ. Villen./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Billaea trochanterata Mesnil, 1970, Mushi 44: 122. ARMENIA.
Holotype ♂: Erivan/ Billaea trochanterata Mesnil L.P. Mesnil det., 1969/ TYPE/ comme Riedel mais chète ant. long+. velu, 2st. -tp autrement sinué Phorostoma sp.
Remarks: The country was published as Georgia, in error.

Billaea (Homalogaster) vitripennis Mesnil, 1950c, Bull. Annls Soc. ent. Belg. 86: 116. ZIMBABWE.
Syntype: 1♂.
♂: GOTA GOTA URUNGWE Dept. Agric. S. Rhodesia 10/9/1938/ W. L. Williams Collector/ 4714/ Coll. 8646/ [blank red label]/ TYPE/ Billaea vitripennis Mesn. L.P. Mesnil det., 1977/ CNC Syntype Billaea (Homalogaster) vitripennis Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.
Remarks: Date of collection of syntype was published as 10 September 1936.

Bithia nova Mesnil, 1973a, Fliegen palaearkt. Reg. 64g: 1146. TURKMENISTAN.

- Holotype ♂: C. AHNGER Tedjen 6.IV 905/ Bithia nova Mesnil L.P. Mesnil det., 1971/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Blepharella nigra* Mesnil, 1967, Mushi 41: 41. JAPAN.
Holotype ♂: Nukabira Hokkaido 24.VII.1964 S. Takano/ 367/ Blepharella nigra Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Blepharella) fuscipennis* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 235. ZAIRE.
Holotype ♂: Penghe 19.II.14/ Blepharella fuscipennis Mesn. L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Blepharella) oldi* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 235. TANZANIA.
Holotype ♂: Nyasaland. Dr.J.E.S.Old. Aluona Ruo Dist X 1912/ Micropodomyia oldi ♂ Typ. Villen./ Blepharella oldi Villeneuve L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Blepharella) setigera* var. *carbonata* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 235. ZAIRE.
Holotype ♂: Escpm. Kabasha: Chambi X-1933/ Blepharella carbonata Mesn. L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Blepharella) setigera* var. *confusa* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 235. SOUTH AFRICA.
Holotype ♂: Metanastria I 3 Kw nap?/ ex Lasiocampid metanastria spec./ Dec. 1915/ Blepharella confusa Mesn. L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Blepharella) setigera* var. *ruficauda* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 235. SOUTH AFRICA.
Holotype ♂: Sydenham marley 6/1915 C.E.P/ Blepharella ruficauda Mesn. L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Blepharellina) picta* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 234. NIGERIA.
Holotype ♀: [underside of round label with "? Caught in ? grass 2.VII.10 ?"]/ Oshogbo, S.Nigeria. Dr.T.F.G.Mayer./ Blepharellina picta Mesn. L.P. Mesnil det., 1966/ TYPE.
- Blepharella (Congochrysosoma) atricauda* Mesnil, 1970, Mushi 44: 97. ZIMBABWE.
Holotype ♂: GOTA GOTA URUNGWE S. Rhodesia. Dept. Agric. 29/7/1938/ Blepharella atricauda Mesn. L.P. Mesnil det., 1970/ TYPE.
- Blepharella (Congochrysosoma) erebiae* Mesnil, 1970, Mushi 44: 96. MALAWI.
Holotype ♀: Mt. Mlanje, Nyasaland, 10.14. 1913. S. A. Neave./ Blepharella erebiae Mesnil L.P. Mesnil det., 1969/ TYPE.
- Blepharella (Congochrysosoma) fallaciosa* Mesnil, 1970, Mushi 44: 96. UGANDA.
Holotype ♂: Uganda Entebbe 14.VIII 1911 C.C.Gowdey./ Blepharella fallaciosa Mesn. L.P. Mesnil det., 1969/ TYPE.
- Blepharella (Congochrysosoma) lodosi* Mesnil, 1968b, Stuttg. Beitr. Naturk. 187: 1. GHANA.
Holotype ♂: TAFO GHANA 13:8:66 LODOS/ Congochrysosoma lodosi Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: Holotype puparium is mounted on paper strip below specimen.
- Blepharella (Congochrysosoma) neglecta* Mesnil, 1968b, Stuttg. Beitr. Naturk. 187: 2. ZAIRE.
Holotype ♂: Walikale I.1915/ Congochrysosoma neglecta Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Blepharella (Congochrysosoma) pellucida* Mesnil, 1970, Mushi 44: 98. ZAIRE.
Holotype ♂: MUSÉE DU CONGO Lulanga :Tsekum-baka - VIII - 1927 J. Ghesquière/ [underside of label] IMP. INST. ENT. COLL.NO. 10520/ R. DET. I. 5209/ Blepharella pellucida Mesn. L.P. Mesnil det., 1969/ TYPE.
- Blepharella (Congochrysosoma) perfida* Mesnil, 1970, Mushi 44: 96. ZAIRE.
Holotype ♂: Katanga : Kafakumba 16 - V - 1925 G. F. Overlaet/ Blepharella perfida Mesnil L.P. Mesnil det., 1969/ TYPE.
- Blepharella (Congochrysosoma) xanthaspis* Mesnil, 1970, Mushi 44: 97. SOUTH AFRICA.
Holotype ♂: Zululand ESHOWE 24.VI.37 L. P. Regnard/ Blepharella xanthaspis Mesn. L.P. Mesnil det., 1969/ TYPE.
- Blepharella (Parapodomyia) claripennis* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 236. ZAIRE.
Holotype ♂: Congo - belge Eala - X - 1953 J.Ghesquière 953/ Parasite chenille de Limacodide/ R. Mus. Hist. Nat. Belg. 10482/ TYPE/ Parapodomyia claripennis Mesnil L.P. Mesnil det., 19/ HOLOTYPE ♂ of Blepharella (Parapodomyia) claripennis Mesn. Examined 1970 R.W.Crosskey.
- Blepharipoda schineri* Mesnil, 1939a, Monogr Sta. Lab. Rech. agron. 7: 32. FRANCE.
Syntypes: 3♂♂, 1♀.
♂: La Celle St Cloud Nr. Versailles 21.5.1939/ Crossocosmia schineri Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ Blepharipa sericariae (Rond.) det. H. Shima, 1992/ CNC Syntype Blepharipoda schineri Mesnil Label affixed 1994.
♂: La Celle St Cloud Nr. Versailles 21.5.1939/ Crossocosmia schineri Mesnil L.P. Mesnil det., 1969/ CNC Syntype Blepharipoda schineri Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

- ♂: La Celle St Cloud Nr. Versailles 28.5.1939/
Crossocosmia schineri Mesnil L.P. Mesnil det.,
1969/ CNC Syntype Blepharipoda schineri Mesnil
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.
- ♀: LesLoges in Josas Nr. Versailles 11.6.1938/
Crossocosmia schineri Mesnil L.P. Mesnil det.,
1969/ CNC Syntype Blepharipoda schineri Mesnil
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.
- Blepharomyia foliacea* Mesnil, 1975b, Fliegen palaearkt. Reg. 64g:
1313. JAPAN.
Holotype ♂: Nukabira Hokkaido 2.3.VII.1949
S.Takano/ 379/ Blepharomyia ciliata Mesnil L.P.
Mesnil det., 1970/ CNC Holotype Blepharomyia
foliacea Mesnil Label affixed 1995/ TYPE/ EX
L.-P. MESNIL COLLECTION 1985.
- Bogosia bequaerti* Villeneuve, 1913, Revue zool. afr. 3: 45.
ZAIRE.
Holotype ♀: Kibombo 2-XI-1910/ Bogosia Bequaerti
Villen. Dr Villeneuve det. Typ./ TYPE/ EX L.-P.
MESNIL COLLECTION 1985.
- Bogosia latifascia* Villeneuve, 1932a, Bull. Annls Soc. ent. Belg.
71 (1931): 244. TAIWAN.
Lectotype ♂: Formosa Sauter/ Kosempo 1911. VII./
Bogosia latifascia n.sp./ LECTO-TYPE/ Bogosia
latifascia Vill. LECTOTYPE ♂ designated by
R.W.Crosskey 1974/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.
Remarks: For lectotype designation see Crosskey
(1976: 265).
- Bogosia similis* Villeneuve, 1926a, Revue zool. afr. 14: 64. ZAIRE.
Syntypes: 2♂♂.
♂: Urwald-Beni Okt.'10.Grauer/ Bogosia similis
♂ n.sp. Villen/ TYPE/ CNC Syntype Bogosia
similis Villeneuve Label affixed 1994/ EX L.-P.
MESNIL COLLECTION 1985.
♂: MUSÉE DU CONGO Albertville ♂ XII-1918
R. Mayné/ CNC Syntype Bogosia similis
Villeneuve Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1985.
- Bombyliomyia purpurea* Thompson, 1963a, Can. J. Zool. 41: 425.
TRINIDAD.
Holotype ♀: HOLOTYPE Bombyliomyia purpurea
Thomp. CNC No. 9779/ Trinidad,BWI. DEC.
1959/ 1125/ Type ♀/ Bombyliomyia purpurea nsp.
- Bonellimyia glauca* Brooks, 1944b, Can. Ent. 76: 200. CANADA.
Holotype ♂: Holo TYPE ♂ Bonellimyia glauca No.
Brks 5398/ Covey Hill,Que. 29.VI 1927
W.J.Brown.
Allotype ♀: Lyn Ont. 10.VIII.1926 F.P.Ide/ AlloTYPE
♀ Bonellimyia glauca Brks No. 5398/ SLIDE No.
0346.
Paratypes: 3♀♀, Nova Scotia; 1♂, Quebec; 1♂, 1♀,
Ontario; 1♂, 1♀, British Columbia; 1♀, Ohio; 1♂,
Michigan; 1♂, Colorado; 1♂, Massachusetts; 1♂,
New Hampshire.

- Bonellimyia subpolita* Brooks, 1944b, Can. Ent. 76: 200.
CANADA.
Holotype ♂: Ottawa,Ont. 2. VI 1927 C.H.Curran/
HoloTYPE ♂ Bonellimyia subpolita No. Brks
5399.
Allotype ♀: MerBleue Ont. 8.VI.1927 G.S.Walley/
AlloTYPE ♀ Bonellimyia subpolita No. Brks
5399.
Paratypes: 4♂♂, Ontario; 1♀, New Brunswick.
- Bothria japonica* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 22.
JAPAN.
Holotype ♂: Sapporo S.Takano [underside of label] 1/V
'26/ 283/ Bothria japonica Mesn. L.P. Mesnil det.,
1966/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.
- Braziliomyia obscura* Thompson, 1968, Mem. ent. Soc. Can. 56:
98. TRINIDAD.
Holotype ♂: HOLOTYPE Braziliomyia obscura
Thomp. CNC No. 9863/ Forest 1 m.w. of Arima
XII.29 60 Trinidad,BWI.
Allotype ♀: ALLOTYPHOPE Braziliomyia obscura Thomp.
CNC No. 9863/ Brazil Trinidad IV.14.60/ 1111
IV.14.60.
- Bucentes albocincta* Villeneuve, 1942a, Bull. Soc. ent. Fr. 47: 55.
ZAIRE.
Holotype ♀: MUSÉE DU CONGO P.N.A.:gîte -1933
Nydragongo, 2300 M. Dr De Wulf/ Bucentes
albocincta Typ. Villen./ Siphona albocincta Villen.
L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
Remarks: Holotype with left mid and both hind legs
glued to mount strip.
- Bucentes nigricans* Villeneuve, 1930c, Bull. Annls Soc. ent. Belg.
70: 100. SWEDEN.
Holotype ♂: Gallivara 24.7.20/ Bucentes nigricans Typ.
Villen./ TYPE/ EX L.-P. MESNIL COLLECTION
1970/ Siphona nigricans Villen L.P. Mesnil det.,
1970/ Holotype ♂ Siphona nigricans Vill. =
hokkaidensis Mesnil = silvarum Herting det. Stig
Andersen 1984.
Remarks: Holotype terminalia are stored in glycerine in
a plastic tube pinned below specimen and
abdomen is glued to a piece of paper below
specimen.
- Cacozelus riederi* Reinhard, 1943a, Can. Ent. 75: 168. USA.
Holotype ♂: Breitenbush Lake Marion Co.Ore.
VI-23-1940 R.E.Rieder,Col./ HOLOTYPE
Cacozelus riederi Reinhard/ Cacozelus riederi
Rnh. det.H.J.Reinhard.
- Cadurcia borbonensis* Villeneuve, 1926b, Revue zool. afr. 14: 245.
RÉUNION.
Syntype: 1♂.
♂: [blank small square label]/ Lophyrom.
sp.aff.clausae Alp.maritum ♂♀/ borbonensis
Villen./ Cadurcia borbonensis Vill. L.P. Mesnil
det., 1969/ TYPE/ CNC Syntype Cadurcia
borbonensis Villeneuve Label affixed 1994/ EX
L.-P. MESNIL COLLECTION 1970.

Cadurcia depressa Villeneuve, 1926b, Revue zool. afr. 14: 244.
ZAIRE.
Syntype: 1♂.
♂: Kayombo 6-V-11/ Dr Villeneuve det./
Cadurcia depressa Typ. Villen./ *Cadurcia depressa*
Villen. L.P. Mesnil det., 1969/ TYPE/ CNC
Syntype *Cadurcia depressa* Villeneuve Label
affixed 1994/ EX L.-P. MESNIL COLLECTION
1970.

Cadurciella rufipalpis Villeneuve, 1927, Revue zool. afr. 15: 120.
ZIMBABWE.
Syntype: 1♀.
♀: 14/12/11 Salisbury 370./ 4 St./ *Cadurciella*
rufipalpis Typ. Villen./ TYPE/ CNC Syntype
Cadurciella rufipalpis Villeneuve Label affixed
1994/ EX L.-P. MESNIL COLLECTION 1970.

Calocarcelia aureocephala Thompson, 1964, Studia ent. 7: 136.
TRINIDAD.
Holotype ♀: HOLOTYPE *Calocarcelia aureocephala*
Thomp. CNC No. 9822/ Santa Cruz V.X.7.60
Trinidad, BWI/ 1299 IV.29.63/ HOLOTYPE
CNCNo.
Paratypes: 3♀♀, Trinidad.

Calocarcelia excisoides Mesnil, 1957, Mém. Soc. r. ent. Belg. 28:
3. JAPAN.
Holotype ♀: Obihiro S.Takano [underside of label]
2/IX'52/ 327/ *Carcelia excisoides* Mesnil L.P.
Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Calocarcelia minima Thompson, 1964, Studia ent. 7: 146.
TRINIDAD.
Holotype ♀: HOLOTYPE *Calocarcelia minima* Thomp.
CNC No. 9825/ Forest 1 m.W.of Arima XII.29.60
Trinidad,BWI/ 1309 VI.14.63/ HOLOTYPE
CNCNo./ *Calocarcelia minima* n.sp.
Paratype: 1♀, Trinidad.

Calocarcelia takanoi Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 1.
JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label]
3/VIII'52/ 165/ *Calocarcelia takanoi* Mesnil L.P.
Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Calocarcelia trinitatis Thompson, 1964, Studia ent. 7: 141.
TRINIDAD.
Holotype ♀: Brazil II.13. 61 Trinidad,BWI/ 1302 V.30
63/ HOLOTYPE *Calocarcelia trinitatis* Thomp.
CNC No.9824.
Allotype ♂: Trinidad,BWI. Brazil IV 2. 60 St.
Augustine./ 1301 V.29. 63/ ALLOTYPE
Calocarcelia trinitatis Thomp. CNC No.9824/
Calocarcelia trinitatis nsp.
Paratypes: 1♂, 2♀♀, Trinidad.
Remarks: Allotype is missing terminalia. Date of
collection of allotype was published as 2 February
1960.

Calodexia callani Thompson, 1968, Mem. ent. Soc. Can. 56: 109.
TRINIDAD.

Holotype ♂: HOLOTYPE *Calodexia callani* Thomp.
CNC No. 9865/ St. Augustine, Trinidad, BWI. 21
Dec. 1943/ E. McC. Callan Collector/ I.C.T.A.
11483/ *Calodexia callani* n.sp.
Paratype: 1♂, Trinidad.

Calodexia nigripes Thompson, 1968, Mem. ent. Soc. Can. 56: 115.
TRINIDAD.
Holotype ♀: HOLOTYPE *Calodexia nigripes* Thomp.
CNC No. 9867/ MARACAS V. Trinidad,BWI.
Nov.11.1952/ Collector F.J.Simmonds/ 1434
VII.8.64/ *Calodexia nigripes* n.sp.
Allotype ♂: ALLOTYPE *Calodexia nigripes* Thomp.
CNC No. 9867/ 1360 1.23.64.
Remarks: The femora and tibia of two legs of holotype
are glued to the locality label. Year of collection
of allotype was published as 1965.

Calodexia rubripes Thompson, 1968, Mem. ent. Soc. Can. 56: 114.
TRINIDAD.
Holotype ♀: HOLOTYPE *Calodexia rubripes* Thomp.
CNC No. 9866/ ST. JOSEPH. Trinidad,BWI.
Nov.12.1953/ Collector F.J.Simmonds/ *Calodexia*
rubripes n.sp./ 14/ 1453 V.3.65.

Calolydella summatis Reinhard, 1974, Can. Ent. 106: 1158.
MEXICO.
Holotype ♂: MEX.Dgo. 14 mi.SW. El Salto,8000' 26
June 1964 W.R.M.Mason/ HOLOTYPE
Calolydella summatis Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 5♂♂, 3♀♀, Mexico.

Calyptromyia barbata Villeneuve, 1915b, Annls hist.-nat. Mus.
natn. hung. 13: 92. TAIWAN.
Holotype ♂: Formosa, Sauter/ Kosempo, 908. V/
Calyptromyia regina, n.g., n.sp.
Remarks: Although the label reads *regina*, the name
was published as *barbata*.

Campogaster vicina Mesnil, 1970, Mushi 44: 119. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label]
9/VII'54/ 336/ *Microsoma vicina* Mesnil L.P.
Mesnil det., 1975/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.
Paratypes: 2♀♀, Japan.

Campylochaeta (Campylochaeta) umbrinervis Mesnil, 1974,
Fliegen palaearkt. Reg. 64g: 1255. JAPAN.
Holotype ♂: 29/IV 1949 Takaosan N.FUKUHARA/
339/ *Campylochaeta umbrinervis* Mesn. L.P.
Mesnil det., 1972/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.
Remarks: Date of collection of holotype was published
as 29 April 1942.

Canelomyia fumator Reinhard, 1958d, J. Kans. ent. Soc. 31: 282.
USA.
Holotype ♂: Canelo, ARIZ. Oct. 11, 1955 G. D. Butler/
HOLOTYPE *Canelomyia fumator* Reinhard/
Canelomyia fumator R Rnh.
Paratype: 1♂, Arizona.

Carcelia abrelicta Mesnil, 1950a, Revue Zool. Bot. afr. 43: 16.
SOUTH AFRICA.

Syntype: 1♀.

♀: Cape Town 1915/ Carcelia abrelicta Mesnil
L.P. Mesnil det., 1964/ Co - type/ CNC Syntype
Carcelia abrelicta Mesnil Label affixed 1994/ EX
L.-P. MESNIL COLLECTION 1970.

Carcelia aequalis Villeneuve, 1939b, Bull. Mus. r. Hist. nat. Belg.
15(48): 1. SOUTH AFRICA.

Syntype: 1♂.

♂: 195/ [locality label illegible]/ Carcelia aequalis
Villen./ Carcelia aequalis Villen. L.P. Mesnil det.,
1969/ Co - type/ CNC Syntype Carcelia aequalis
Villeneuve Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

Carcelia angulicornis Villeneuve, 1916c, Ann. S. Afr. Mus. 15:
481. NIGERIA.

Syntypes: 1♂, 1♀.

♂: Oshogbo, S.Nigeria. Dr.T.F.G.Mayer.
20-30IX.1910/ Dr Villeneuve det./ Carcelia
angulicornis Villen. Typ./ TYPE/ CNC Syntype
Carcelia angulicornis Villeneuve Label affixed
1994/ EX L.-P. MESNIL COLLECTION 1970.
♀: Oshogbo, S.Nigeria. Dr.T.F.G.Mayer. 1-13.
Octobor [sic] 1910/ Carcelia angulicornis Vill.
L.P. Mesnil det., 1964/ Co - type/ CNC Syntype
Carcelia angulicornis Villeneuve Label affixed
1994/ EX L.-P. MESNIL COLLECTION 1970.

Carcelia dilaticornis Mesnil, 1950a, Revue Zool. Bot. afr. 43: 14.
MOROCCO.

Syntypes: 2♂♂, 2♀♀.

♂: 3095 Maroc Goundafa exP.14.2.47 Ch. Rungs/
ex Phragmatobia sp?/ Parexorista dilaticornis n.sp
L.P. Mesnil det., 1964/ TYPE/ Carcelia
dilaticornis Mes. L.P. Mesnil det., 1969/ CNC
Syntype Carcelia dilaticornis Mesnil Label affixed
1994/ EX L.-P. MESNIL COLLECTION 1970.
♂: Same data as above except without TYPE
label.
2♀♀: 3095 Maroc Goundafa exP.17.2.47 Ch.
Rungs/ ex Phragmatobia sp?/ Parexorista
dilaticornis Mesn. L.P. Mesnil det., 1967/ Carcelia
dilaticornis Mesn. L.P. Mesnil det., 1969/ CNC
Syntype Carcelia dilaticornis Mesnil Label affixed
1994.

Remarks: Second male syntype is missing terminalia.

Carcelia inusta Mesnil, 1950a, Revue Zool. Bot. afr. 43: 11.
SOUTH AFRICA.

Syntype: 1♂.

♂: 467/ Marley Stella B 4 -1915/ in shady places
low bush/ Carcelia inusta Mesn. L.Mesnil det.,
1949/ TYPE/ CNC Syntype Carcelia inusta Mesnil
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

Carcelia lucidula Villeneuve, 1941b, Bull. Soc. ent. Fr. 46: 125.
ZAIRE.

Syntype: 1♀.

♀: D.2. MUSÉE DU CONGO Uele: Dembia
10-XI-1933 J. V. Leroy/ lucidula Villen./ Carcelia

lucidula Vill. L.P. Mesnil det., 1969/ Co - type/
CNC Syntype Carcelia lucidula Villeneuve Label
affixed 1994/ EX L.-P. MESNIL COLLECTION
1970.

Carcelia noumeensis Mesnil, 1969, Entomophaga 13: 203. NEW
CALEDONIA.

Holotype ♂: ex Noctuidae s/Avoine - Nouméa. X.
1964/ Carcelia noumeensis Mesnil L.P. Mesnil
det., 1967/ TYPE/ HOLOTYPE ♂ of Carcelia
noumeensis Mesnil Examined 1970
R.W.Crosskey.

Carcelia oblectanea Mesnil, 1950a, Revue Zool. Bot. afr. 43: 15.
ZAIRE.

Syntype: 1♀.

♀: R 2000m 12.IV.14/ Carcelia oblectanea Mesn.
L.P. Mesnil det., 1964/ Co - type/ CNC Syntype
Carcelia oblectanea Mesnil Label affixed 1994/
EX L.-P. MESNIL COLLECTION 1970.

Carcelia oblimata Mesnil, 1950a, Revue Zool. Bot. afr. 43: 14.
SOUTH AFRICA.

Syntype: 1♀.

♀: Cape Town/ Carcelia oblimata Mesnil L.P.
Mesnil det., 1969/ Co - type/ CNC Syntype
Carcelia oblimata Mesnil Label affixed 1994/ EX
L.-P. MESNIL COLLECTION 1970.

Carcelia oblitterata Mesnil, 1950a, Revue Zool. Bot. afr. 43: 13.
SOUTH AFRICA.

Holotype ♀: Kransp. 30.I.07. 6197 with Chrys./
Carcelia oblitterata Mesn. L.P. Mesnil det., 1969/
TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Carcelia peraequalis Mesnil, 1950a, Revue Zool. Bot. afr. 43: 18.
ZIMBABWE.

Syntype: 1♂.

♂: Mfongosi Zulu L. W E Jones/ Carcelia
peraequalis Mesn. L.P. Mesnil det., 1969/ Co -
type/ CNC Syntype Carcelia peraequalis Mesnil
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

Remarks: The type locality was cited as Salisbury (now
Harare), so this specimen is possibly not part of
the original syntype series.

Carcelia pollinosa Mesnil, 1941b, Bull. Soc. ent. Fr. 46: 98.
CZECH REPUBLIC.

Lectotype ♂: F. Kowarz Franzsbd./ Extrema W. of
Bohemia C S R/ Eucarcelia pollinosa Mesnil L.P.
Mesnil det., 1969/ TYPE/ CNC Syntype Carcelia
pollinosa Mesnil Label affixed 1994/ Lectotype
fixation by Mesnil, 1963: 3. Label affixed 1995/
EX L.-P. MESNIL COLLECTION 1970.

Remarks: We regard Mesnil's (1963a: 3) reference to a
holotype as a lectotype fixation for this species.

Carcelia (Carcelina) nigrapex Mesnil, 1944a, Fliegen palaearkt.
Reg. 64g: 29, in key (1949a: 53, description).
CHINA.

Lectotype ♀: Kou-ling 22-8-18/ Carcelina nigrapex
Mesn. L. Mesnil, det/ Co - type/ LECTO-TYPE/
Carcelia nigrapex Mesnil LECTOTYPE ♀
designated by R.W.Crosskey 1974.

Remarks: For lectotype designation see Crosskey (1976: 265).

Carcelia (Parexorista) delicatula Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 173. INDIA.
Holotype ♂: Dehra Dun, U.P. J.C.M. Gardner, 22.I. 1940/ 911/ Parexorista delicatula n.sp. L.P. Mesnil det., 1967/ Parasitic on ? Laelia cocoon under stone/ TYPE/ Carcelia delicatula Mesn. L.P. Mesnil det., 1969/ HOLOTYPE ♂ of Carcelia (Parexorista) delicatula Mesn Examined 1970 R.W.Crosskey.
Paratype: 1♀, India.

Carmodymyia aencylostomiae Thompson, 1968, Mem. ent. Soc. Can. 56: 130. TRINIDAD.
Holotype ♂: HOLOTYPE Carmodymyia aencylostomiae Thomp. CNC No. 9871/ PARADISE MT. TRINIDAD W.I. MARCH. 1959./ Ex-POD PIGEON PEA/ F.D.Bennett Collector.
Allotype ♀: ALLOTYPE Carmodymyia aencylostomiae Thomp. CNC No. 9871/ AROUCA MT. TRINIDAD,BWI. JULY. 1953./ REARED FROM ANCYLOSTOMIA STERCOREAIN PIGEON PEA/ Collector F.D. Bennett/ 76.
Remarks: Holotype puparium is glued to a paper strip below specimen.

Catagonia subdistincta Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 484. SOUTH AFRICA.
Syntype: 1♂.
♂: frequents a sand creeper where it hides away beneath the leaves very sharp & active fly/ Point Durhan apl. 1915 Marley/ Catagonia subdistincta ♂ Villen./ Sisyropa subdistincta Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Catagonia subdistincta Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Catalinovoria discalis Brooks, 1945b, Can. Ent. 77: 81. CANADA.
Holotype ♂: Fton, [Fredericton] N. B. 7.VIII 1913 J. D. Tothill/ Holo TYPE ♂ Catalinovoria discalis No. Brks 5518.
Allotype ♀: Fort à la Corne Sask. 14. 7.1925 Kenneth M.King/ Allo TYPE ♀ Catalinovoria discalis No. Brks 5518.
Paratypes: 1♀, Saskatchewan; 1♀, Alberta; 1♀, Ohio.

Cataphoriniopsis fumipennis Thompson, 1968, Mem. ent. Soc. Can. 56: 134. TRINIDAD.
Holotype ♀: 1371 1.29.64/ HOLOTYPE Cataphoriniopsis fumipennis Thomp. CNC No. 9872.
Remarks: Holotype is missing terminalia and right wing.

Cataphoriniopsis galbae Thompson, 1968, Mem. ent. Soc. Can. 56: 137. TRINIDAD.
Holotype ♂: HOLOTYPE Cataphoriniopsis galbae Thomp. CNC No. 9873/ Trinidad,BWI. 30.IX.56/ Ex 'Galba' seeds (C.antillanum)/ 16278 I.C.T.A./ M.H.Breese Collector/ B165/ COM. INST ENT.

COLL NO 14680/ Cataphorinia? sp not in B.M. van Emden det. 1956.

Allotype ♀: ALLOTYPE Cataphoriniopsis galbae Thomp. CNC No. 9873/ St. Augustine, TRINIDAD,BWI. 1.30.60.

Remarks: Holotype head and terminalia are glued to a paper strip below specimen.

Catemophrys sororcula Reinhard, 1974, Can. Ent. 106: 1159. MEXICO.

Holotype ♀: Techuan Puebla,Mex. VI-23-1951/ On fls. of Eysenhardtia polystachya(Ort.)/ P.D. Hurd Collector/ HOLOTYPE Catemophrys sororcula Rnh. CNC No.

Cavalieria genibarbis Villeneuve, 1908b, Feuille jeun. Nat. 38: 114, 116. FRANCE.

Lectotype ♂: Cavalière (Var) VU-06/ Cavalieria genibarbis type ♂/ Cavalieria genibarbis Vill. L.P. Mesnil det., 1969/ CNC Syntype Cavalieria genibarbis Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1954: 320. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: We regard Mesnil's (1954a: 320) reference to a holotype as a lectotype fixation for this species.

Celatoria maracasi Thompson, 1968, Mem. ent. Soc. Can. 56: 172. TRINIDAD.

Holotype ♀: HOLOTYPE Celatoria maracasi Thomp. CNC No. 9880/ MARACAS V. TRINIDAD,BWI DEC.2.1952/ Collector F.J.Simmonds/ 53/ Celatoria maracasi n.sp.

Celotrophus soporis Reinhard, 1958d, J. Kans. ent. Soc. 31: 281. USA.

Holotype ♂: Amherst, Ohio V-10-1924/ H J Reinhard Collector/ HOLOTYPE Celotrophus soporis Reinhard/ Celotrophus soporis R Rnh./ Ictericophyto catskillensis (West) Reinhard Coll'n. Allotype ♀: Same data as holotype except with allotype label.
Paratype: 1♂, Ohio.

Cephaloplagia nubecula Reinhard, 1964a, J. Kans. ent. Soc. 37: 44. USA.

Holotype ♂: 1 mi.S.E Murphy Ida. Owyhee Co. V.27 1958/ W. F. Barr Collector/ HOLOTYPE Cephaloplagia nubecula Reinhard/ Cephaloplagia nubecula [sic] R (Rnh.).

Remarks: Holotype terminalia are glued to locality label.

Ceranthia japonica Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 33. JAPAN.

Holotype ♂: Kyushu K.KAMIJO [underside of label with "IX. 1958"]/ Ceranthia japonica n.sp. L.P. Mesnil det., 1963/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Ceranthia sulfurea Mesnil, 1971a, Entomophaga 16: 72. JAPAN.
Holotype ♀: Nakabira Hokkaido 27. VII. 1959 S.Takano/ 386/ Ceranthia sulfurea Mesnil L.P.

Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.

Ceratochaeta clausa Villeneuve, 1937b, Bull. Mus. r. Hist. nat. Belg. 13(13): 1. "LAPLAND".
Holotype ♂: Ceratochaeta clausa typ. Villen./ TYPE/ P. clausa Vill. L.P. Mesnil det., 1969/ EX L.-P. MESNIL COLLECTION 1970.

Ceratochaeta (Gymnophryxe) nudigena Villeneuve, 1922a, Bull. Mus. natn. Hist. nat., Paris 28: 293. ALGERIA.
Lectotype ♀: Parasite de l'Orgyia dubia var. des environs de Laghouat. Chenill fin Oct. 1911. eclos. parasit 28.2.1912/ Gymnophryxe nudigena Typ ♀/ TYPE/ Archiclops nudigena Vill. L.P. Mesnil det., 1969/ CNC Syntype Ceratochaeta (Gymnophryxe) nudigena Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1956: 514. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: We regard Mesnil's (1956b: 514) reference to a holotype as a lectotype fixation for this species.

Ceratochaeta (Lasiopales) pachychaeta Villeneuve, 1922a, Bull. Mus. natn. Hist. nat., Paris 28: 292. ALGERIA.
Lectotype ♂: Parasite de L. Staudingeri Lambèse Eclos. 3.4.1914/ Lasiopales pachychaeta Typ ♂/ Lasiopales pachychaeta Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Ceratochaeta (Lasiopales) pachychaeta Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1956: 514. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Lectotype right mid tibia and tarsus are glued to a piece of paper pinned below specimen. We regard Mesnil's (1956b: 514) reference to a holotype as a lectotype fixation for this species.

Ceromasia borealis Brooks, 1945b, Can. Ent. 77: 89. CANADA.
Holotype ♀: HoloTYPE ♀ Ceromasia borealis No. Brks 5537/ Cameron Bay, Great Bear Lake, N W T 27.VII-1937 T.N.Freeman.
Allotype ♂: Banff, Alta 3VIII.1922 C.B.D.Garrett 8000ft/ AlloTYPE ♂ Ceromasia borealis No. Brks 5537.
Paratypes: 2♀♀, Northwest Territories; 1♀, Alberta.
Remarks: Date of collection of holotype was published as 27 August 1937.

Ceromya balli O'Hara, 1994, Can. Ent. 126: 786. CANADA.
Holotype ♂: Masham Twp.QUE. Gatineau Co. 27-31.V.1974 D. M. Wood/ HOLOTYPE Ceromya balli O'Hara CNC No. 21512.
Allotype ♀: Same data as holotype except with allotype label and collected "3-7.VI.1974".
Paratypes: 2♂♂, 1♀, New Brunswick; 72♂♂, 7♀♀, Quebec; 54♂♂, 11♀♀, Ontario; 4♂♂, Manitoba; 3♂♂, Alberta; 3♂♂, 1♀, Maryland.

Ceromya flava O'Hara, 1994, Can. Ent. 126: 790. CANADA.
Holotype ♂: Masham Twp.Que Gatineau Co. 4-7.VI 1975 D.M.Wood/ A67/ HOLOTYPE Ceromya flava O'Hara CNC No. 21513.

Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.

Ceromya occidentalis O'Hara, 1994, Can. Ent. 126: 793. CANADA.
Holotype ♂: Lakelse L. bog nr. Terrace,B.C. 14.VI 1960 C.H. Mann/ HOLOTYPE Ceromya occidentalis O'Hara CNC No. 21515.
Allotype ♀: Lumby, B.C. June 22,1973 H.J. Teskey/ ALLOTYPE Ceromya occidentalis O'Hara CNC No. 21515.
Paratypes: 5♂♂, 4♀♀, British Columbia; 1♂, 1♀, California.

Ceromya oriens O'Hara, 1994, Can. Ent. 126: 791. CANADA.
Holotype ♂: Masham Twp.QUE. Gatineau Co. 15. V .1977 D.M.Wood/ HOLOTYPE Ceromya oriens O'Hara CNC No. 21514.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 17♂♂, 3♀♀, New Brunswick; 3♂♂, 2♀♀, Nova Scotia; 45♂♂, 30♀♀, Quebec; 31♂♂, 11♀♀, Ontario; 1♂, 2♀♀, North Carolina; 2♂♂, New York; 1♂, 1♀, Michigan; 1♀, Massachusetts.

Ceromyia (Proceromyia) macronychia Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 35. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label with "10/V'49"]/68/ Proceromyia macronychia Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Cestonionerva latigena Villeneuve, 1939a, Bull. Annls Soc. ent. Belg. 79: 353. CHINA.
Holotype ♀: Tchounsin 2.VI.18 [underside of label with "Ordos Licent"]/ Cestonionerva latigena Type Villen/ Cestonionerva latigena Villeneuve L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Cestonoptera mesnili Villeneuve, 1939a, Bull Annls Soc. ent. Belg. 79: 348. TUNISIA.
Holotype ♂: Tunis. V. 52779./ Cestonoptera mesnili Type Villen./ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Chaetexorista klapperichi Mesnil, 1960b, Fliegen palaearkt. Reg. 64g: 645. CHINA.
Holotype ♂: Kuatun(2300m)27,40n.Br. 117,406.L.J.Klapperich 25.5.1938(Fukien)/ Chaetexorista klapperichi Mesn L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Chaetogena (Spoggosia) soror Mesnil, 1971a, Entomophaga 16: 69. TURKMENISTAN.
Holotype ♂: Tr. Casp. Gr. Balchan/ TYPE/ Chaetogena soror Mesnil L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1985.

Chaetona pictilis Reinhard, 1974, Can. Ent. 106: 1159. MEXICO.
Holotype ♀: Chilpancingo, Guer., Mex. 7-24-61 R.&K. Dreisbach/ HOLOTYPE Chaetona pictilis Rnh. CNC No./ HOMOTYPE Chaetona congrua Wulp Compared by Wood '79.

Chaetona tuchucheensis Thompson, 1968, Mem. ent. Soc. Can. 56: 87. TRINIDAD.

Holotype ♀: HOLOTYPE *Chaetona tuchucheensis* Thomp. CNC No. 9860/ El Tucuche, 3000ft. Trinidad,BWI. 9 Sept. 1941/ E. McC. Callan Collector/ I.C.T.A. 9817/ *Chaetona tuchucheensis* [sic] n.sp.

Chaetophlepsis atriceps Reinhard, 1952a, J. Kans. ent. Soc. 25: 21. USA.

Holotype ♂: Amherst, Ohio VI-26 1924/ H J Reinhard Collector/ HOLOTYPE *Chaetophlepsis atriceps* Reinhard.

Chaetophlepsis polita Brooks, 1945b, Can. Ent. 77: 83. USA.

Holotype ♂: Agr. Col. Miss. IV-23 1921 H. W. Allen/ Holotype ♂ *Chaetophlepsis polita* No. Brks 5521.

Allotype ♀: Aylmer, Que. 18.VII 1924 C.H.Curran/ AlloTYPE ♀ *Chaetophlepsis polita* No. Brks 5521.

Chaetophlepsis semiothisae Brooks, 1945b, Can. Ent. 77: 82. CANADA.

Holotype ♂: Constance Bay Ont. 6-VII -'35 T.N.Freeman/ Holotype ♂ *Chaetophlepsis semiothisae* No. Brks 5520.

Allotype ♀: Same data as holotype except with allotype label an dated "V-VII -'35".

Paratypes: 1♀, Nova Scotia; 1♂, 1♀, New Brunswick; 1♂, 13♀♀, Quebec; 2♂♂, 6♀♀, Ontario; 5♂♂, 1♀, British Columbia.

Remarks: Holotype puparium is pinned below specimen.

Chaetophlepsis teliosis Reinhard, 1952a, J. Kans. ent. Soc. 25: 18. USA.

Holotype ♂: Raleigh, NC VIII-17-26/ HOLOTYPE *Chaetophlepsis teliosis* Reinhard.

Chaetoria micronyx Mesnil, 1971a, Entomophaga 16: 71. NEW CALEDONIA.

Holotype ♂: Noumea NEW CALEDONIA 12 II 1957 CharlesP.Hoyt/ *Chaetoria micronyx* Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Chaetovoria melancholica Mesnil, 1953b, Bull. Annls Soc. ent. Belg. 89: 171. "PALESTINE".

Holotype ♀: PALESTINE 1946 J.Palmoni/ 6620/ Uclesia melancholica Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Charasoma pammelas Reinhard, 1952b, Bull. Brooklyn ent. Soc. 47: 12. USA.

Holotype ♂: Forest Grove Ore V-20-18/ FRCole Coll/ ♂/ HOLOTYPE *Charasoma pammelas* Reinhard/ *Charasoma pammelan* [sic] R. Rnh./ *Charasoma pammelas* R. Rnh.
Paratype: 1♂, Oregon.

Charasoma residis Reinhard, 1952b, Bull. Brooklyn ent. Soc. 47: 12. USA.

Holotype ♀: Amherst, Ohio 7-12 1934/ H J Reinhard Collector/ HOLOTYPE *Charasoma residis* Reinhard/ *Charasoma residis* R. Rnh.
Paratype: 1♀, Michigan.

Charasoma subolis Reinhard, 1952b, Bull. Brooklyn ent. Soc. 47: 11. USA.

Holotype ♂: College Station Apr. 11, 1944 Tex/ H J Reinhard Collector/ HOLOTYPE *Charasoma subolis* Reinhard/ *Charasoma subolis* R. Rnh./ HOMOTYPE *Erycia seticauda* Rnh Compared by Wood'80.

Allotype ♀: Same data as holotype except with allotype label and collected "Apr.16,1946".

Paratype: 1♀, New York.

Charitella nigrescens Mesnil, 1977, Bull. Soc. ent. Suisse. 50: 325. MALAWI.

Holotype ♀: Mt. Mlanje, Nyasaland, 10.IX. 1913. S. A. Neave./ *Charitella nigrescens* Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Paratypes: 1♂, 1♀, Madagascar.

Chesippus notialis Reinhard, 1967, J. Kans. ent. Soc. 40: 107. USA.

Holotype ♂: Sta Rita MtsAriz/ Jun DKDuncan/ HOLOTYPE *Chesippus notialis* Reinhard.

Allotype ♀: Santa Rita Mts Ariz 7-19-38/ ALLOTYPE *Chesippus notialis* Reinhard.

Paratype: 1♀, Arizona.

Chlorolydella schistacea Mesnil, 1955a, Annls Mus. r. Congo belge Sér. 8vo (Zool.) 40: 365. RWANDA.

Holotype ♂: Congo belge: Ruanda Kibga (Volc. Bishoke) 2400 m. 11-ii-1935 G. F. De Witte: 1105/ *Chlorolydella schistacea* Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Chloropales luteifacies Mesnil, 1950b, Fliegen palaearkt. Reg. 64g: 109. PAPUA NEW GUINEA.

Syntype: 1♀.

♀: NeuPommern Kinigunang C.Ribbe/ coll.Lichtwardt/ *Chloropales luteifacies* Mesn. L.P. Mesnil det., 1966/ CNC Syntype *Chloropales luteifacies* Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Cistogaster atrota Reinhard, 1935c, Ann. ent. Soc. Am. 28: 173. USA.

Holotype ♀: Amherst, Ohio June 1934. A.J.Barckert Collector/ HOLOTYPE *Cistogaster atrota* Reinhard/ *Cistogaster atrota* det.H.J.Rein Rein./ *Gymnoclytia atrota* R Rnh.

Clairvillia amicta Reinhard, 1962a, Ent. News 73: 169. USA.

Holotype ♂: Canelo,Ariz. VII-20-1958 M. Adachi/ HOLOTYPE *Clairvillia amicta* Reinhard/ *Clairvillia amicta* R Rnh.

Paratype: 1♂, Arizona.

Clairvillia curialis Reinhard, 1958e, Ent. News 69: 235. MEXICO.

- Holotype ♂: Cuernavaca, Mex.9-28-57
R.&K.Dreisbach/ HOLOTYPE Clairvillia curialis
Reinhard/ Clairvillia curialis R Rnh.
Allotype ♀: Cuernavaca, Morelos,Mex. 10-29-57
R.&K.Dreisbach/ ALLOTYPE Clairvillia curialis
Reinhard.
Paratype: 1♀, Mexico.
- Clastoneuriopsis meralis* Reinhard, 1939, Bull. Brooklyn ent. Soc. 34: 69. USA.
Holotype ♂: Vantage, Wn. [Washington] IV-1- 1933/
CHAS.H. MARTIN/ HOLOTYPE
Clastoneuriopsis meralis Reinhard.
Paratypes: 3♂♂, Washington.
- Clemelis gymnops* Herting, 1975, Stuttg. Beitr. Naturk. (A) 287: 3.
TUNISIA.
Holotype ♂: Parasite Phycita ricinivora Dumont Nefta
28.IV.27/ MUSEUM PARIS TUNISIE NEFTA C.
DUMONT 1927/ *Clemelis apicalis* Villen. L.P.
Mesnil det., 1969/ *Clemelis gymnops* B. Herting
det. n.sp. Type/ EX L.-P. MESNIL COLLECTION
1970.
- Clemelis majuscula* Mesnil, 1954a, Fliegen palaearkt. Reg. 64g:
349. FRANCE.
Holotype ♂: Digne/ *Clemelis majuscula* Mesn. L.P.
Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
- Clemelis pullata obliquata* Mesnil, 1954a, Fliegen palaearkt. Reg.
64g: 352. FRANCE.
Holotype ♂: La Grave 56827/ *Clemelis oblectanea*
Mesnil L.P. Mesnil det., 1969./ TYPE/ *Clemelis*
pullata Meig. B. Herting det.
Remarks: Holotype was labeled as *Clemelis oblectanea*
(an unpublished name) by Mesnil, in error.
- Clistomorpha alberta* Curran, 1927b, Can. Ent. 59: 298. CANADA.
Holotype ♂: *Clistomorpha* TYPE alberta ♂ Curran
No.2463/ Slave Lake, Alta 14.VI.1924. O. Bryant.
- Clistomorpha deceptor* Curran, 1927b, Can. Ent. 59: 298.
CANADA.
Holotype ♂: *Clistomorpha* TYPE deceptor Curran
No.2457./ Hedley,B.C. 21VII 1923 C.B.Garrett.
- Cloacina filialis* Reinhard, 1945, Can. Ent. 77: 35. USA.
Holotype ♂: Hartley Co VI-14-34 Tx/ HOLOTYPE
Cloacina filialis Reinhard/ *Cloacina filialis* R Rnh.
Paratype: 1♂, Texas.
- Collatia ornata* Reinhard, 1937, Bull. Brooklyn ent. Soc. 32: 71.
USA.
Holotype ♂: Colorado Aug. 1933/ Holotype *Collatia*
ornata Reinhard/ *Collatia ornata* Rein.
det.H.J.Reinhard/ Meigenia submissa (A&W)
Reinhard Coll'n/ Meigenia submissa (Ald.+Web.).
- Colorus downsi* Reinhard, 1953b, J. Kans. ent. Soc. 26: 98.
MEXICO.
- Holotype ♂: HOLOTYPE *Colorus downsi* Reinhard/
AMANALCO MEX 5-V-50 coll.WGDowns.
Compsilurooides flavipalpis Mesnil, 1957, Mém. Soc. r. ent. Belg.
28: 22. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label with
"9/VII'50 h + s"]/ 93/ *Compsilurooides flavipalpis*
Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P.
MESNIL COLLECTION 1970.
- Compsoptesis phoenix* Villeneuve, 1915b, Annls hist.-nat. Mus.
natn. hung. 13: 91. TAIWAN.
Syntype: 1♂.
♂: Formosa, Sauter/ Sokotsu, 1912 V./
Compsoptesis phoenix, Type Villen.
- Compsoptesis rufula* Villeneuve, 1915b, Annls hist.-nat. Mus. natn.
hung. 13: 91. TAIWAN.
Holotype ♂: Formosa, Sauter/ Tainan, 1911 III./
Compsoptesis rufula, Type Villen.
- Cowania wheeleri* Reinhard, 1952b, Bull. Brooklyn ent. Soc. 47: 3.
MEXICO.
Holotype ♂: Fresnillo, Zac. Aug. 25, 1947/ F. A.
Cowan M. R. Wheeler Collectors/ HOLOTYPE
Cowania wheeleri Reinhard/ *Cowania wheeleri* R.
Rnh.
Allotype ♀: Same data as holotype except with allotype
label and missing last label.
Paratypes: 3♂♂, Mexico.
- Crocinosoma cornualis* Reinhard, 1947, J. Kans. ent. Soc. 20: 21.
USA.
Holotype ♀: College Station Jun.29,1944 Tex/ H J
Reinhard Collector/ HOLOTYPE *Crocinosoma*
cornualis Reinhard.
Allotype ♂: Austin Tex 27.IX.22/ ALLOTYPE
Crocinosoma cornualis Reinhard.
- Crocuta antennalis* Mesnil, 1952a, Bull. Inst. r. Sci. nat. Belg.
28(23): 9. ZIMBABWE.
Holotype ♂: Salisbury S. Rhodesia, 5. V. 1935 A.
Cuthbertson/ *Siphona antennalis* Mesn. L.P.
Mesnil det., 1970/ TYPE.
- Crossocosmia jacobsoni angustifrons* Mesnil, 1967, Mushi 41: 41.
JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label]
16/IX'50/ 211/ *Crossocosmia angustifrons* Mesn.
L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970/ *Blepharipa angustifrons*
(Mesn) det. H. Shima, 1992.
Remarks: Holotype terminalia are glued to a paper
mount below specimen.
- Crossocosmia (Blepharipa) carbonata* Mesnil, 1970, Mushi 44: 92.
JAPAN.
Holotype ♂: Mt. Moiwa Sapporo 27.VII.1965
S.Takano/ 398/ *Crossocosmia carbonata* Mesn.
L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970/ *Blepharipa carbonata*
(Mesn) det. H. Shima, 1992.
Remarks: Holotype terminalia are glued to a paper
mount below specimen.

Crossocosmia (Blepharipa) latigena Mesnil, 1970, Mushi 44: 92.
JAPAN.

Holotype ♂: Miyazaki K.Shimizu [underside of label]
3/V '55/ 166/ Crossocosmia latigena Mesn. L.P.
Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970/ Blepharipa latigena (Mesn.)
det. H. Shima, 1992.

Crossocosmia (Blepharipa) nigrina Mesnil, 1970, Mushi 44: 94.
CHINA.

Holotype ♂: Mandschurei Charbin, 28.6.53 V. Alin leg.,
vend. Eing.Nr.25,1953/ Crossocosmia nigrina
Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P.
MESNIL COLLECTION 1970.

Cryptomeigenia brimleyi Reinhard, 1947, J. Kans. ent. Soc. 20: 17.
USA.

Holotype ♂: Raleigh, NC CSBrimley 8-VI-1926/
HOLOTYPE Cryptomeigenia brimleyi Reinhard/
Cryptomeigenia brimleyi R Rnh.
Allotype ♀: Raleigh, N.C 1-VII-31 C.S.Brimley/ B1E/
ALLOTYPE Cryptomeigenia brimleyi Reinhard.

Cryptomeigenia crassipalpis Reinhard, 1947, J. Kans. ent. Soc. 20:
18. USA.

Holotype ♀: Babylon, L.I., N.Y. V-21 1933 F. S.
Blanton/ HOLOTYPE Cryptomeigenia
crassipalpis Reinhard/ Cryptomeigenia
crassipalpis R Rnh.
Paratype: 1♀, New York.

Cryptomeigenia dubia Curran, 1926a, Trans. R. Soc. Can., Ser. 3,
20: 164. CANADA.

Holotype ♀: C. HoloTYPE ♀ dubia Curran No. 1949/
Pt. Pelee, Ont. 23-VI 1925 G.S.Walley.

Cryptomeigenia muscoidea Curran, 1926a, Trans. R. Soc. Can., Ser.
3, 20: 157. CANADA.

Holotype ♂: Cryptomeigenia TYPE muscoidea Curran
No. 1489/ Aweme, Man. 16 V 1916 N. Criddle/
Host Phyllophaga rugosa/ 8078.

Allotype ♀: Same data as holotype except with allotype
label and collected "24 V 1916".

Remarks: Holotype abdomen is glued to a paper strip
below specimen, and terminalia are stored in
glycerine in a glass microvial pinned below
specimen. Allotype abdomen is glued to locality
label.

Cryptomeigenia nigripes Curran, 1926a, Trans. R. Soc. Can., Ser.
3, 20: 162. CANADA.

Holotype ♂: C. HoloTYPE ♂ nigripes Curran No. 1950/
Bothwell Ont. 23.V. 1925 G.S.Walley.

Allotype ♀: Same data as holotype except with allotype
label.

Cryptomeigenia nigripilosa Curran, 1926a, Trans. R. Soc. Can.,
Ser. 3, 20: 161. CANADA.

Holotype ♂: Cryptomeigenia TYPE nigripilosa Curran.
No. 1490/ Lillooet, BC. 17.VI.1917 J.D. Tothill.

Cryptomeigenia ochreigaster Curran, 1926a, Trans. R. Soc. Can.,
Ser. 3, 20: 165. CANADA.

Holotype ♂: TYPE Cryptomeigenia ochreigaster No.
Curran 1456/ Aweme, Man. R.M.White 31. V.
1923.

Cryptomeigenia ontario Curran, 1926a, Trans. R. Soc. Can., Ser. 3,
20: 159. CANADA.

Holotype ♂: Cryptomeigenia HoloTYPE ♂ ontario
Curran No. 1491/ O 13.5.01.
Allotype ♀: BELLEVILLE ONT., CAN./
Cryptomeigenia AlloTYPE ♀ ontario Curran No.
1491.

Paratype: 1♀, Quebec.

Remarks: Date of collection of holotype was published
as 13 May 1910.

Cryptomeigenia simplex Curran, 1926a, Trans. R. Soc. Can., Ser. 3,
20: 163. CANADA.

Holotype ♂: C. HoloTYPE ♂ simplex Curran No. 1947/
Hull, Que 28.V.1924 C.H.Curran.
Allotype ♀: Same data as holotype except with allotype
label and collected "21 V 1924".

Cryptomeigenia triangularis Curran, 1926a, Trans. R. Soc. Can.,
Ser. 3, 20: 160. CANADA.

Holotype ♂: C. HoloTYPE ♂ triangularis Curran No.
1948/ Queens Park, Aylmer 14.V.1924 Que. C B.
Hutchings.

Allotype ♀: Same data as holotype except with allotype
label and collected "15.6 1925".
Paratypes: 1♂, Quebec; 1♀, Manitoba.

Cryptomeigenioidea longipes Thompson, 1968, Mem. ent. Soc.
Can. 56: 40. TRINIDAD.

Holotype ♂: HOLOTYPE Cryptomeigenioidea longipes
Thomp. CNC No. 9848/ Simla Trinidad/ 1473
V.16.66.

Allotype ♀: ALLOTYPE Cryptomeigenioidea longipes
Thomp. CNC No. 9848/ Simla Trinidad.

Paratype: 1♀, Trinidad.
Remarks: Holotype is missing terminalia. Year of
collection of holotype was published as 1965.

Ctenophorinia grisea Mesnil, 1967, Mushi 41: 43. JAPAN.

Holotype ♂: Maruyama Sapporo 15.VI.1966 S. Takano/
408/ Ctenophorinia grisea Mesn. L.P. Mesnil det.,
1967/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.

Ctenophorocera angustifrons Mesnil, 1963a, Bull. Inst. r. Sci. nat.
Belg. 39(24): 6. JAPAN.

Holotype ♂: Sapporo Hokkaido, 2.VIII T.Kumata 1957/
Host Bombyx mori L. (Lep.: Bombycidae)/
Ctenophorocera angustifrons n.sp. L.P. Mesnil
det., 1962/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970/ Blepharipa angustifrons
(Mesn) det. H. Shima 1992.

Remarks: Holotype puparium is pinned below
specimen.

Ctenophorocera (Ctenophorocera) aethiopica Mesnil, 1950b,
Fliegen palaearkt. Reg. 64g: 124. TANZANIA.

Holotype ♂: Urw.hint.d.Randbg. d.N.W.Tanganika-S.
18-2200 m. Grauer/ Pales aethiopica Mesn. L.P.
Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

- Ctenophorocera (Ctenophorocera) coerulea* var. *coeruleonigra*
Mesnil, 1950b, Fliegen palaearkt. Reg. 64g: 126.
ZIMBABWE.
Holotype ♂: Vumba mts. Umtali Dist. Dept. Agric S.
Rhodesia Mar 1935/ A.Cuthbertson Collector/
Pales coeruleo-nigra Mesn. L.P. Mesnil det., 1969/
TYPE/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: Holotype right hind leg is glued to locality
label.
- Ctenophorocera (Ctenophorocera) coerulea maculisquama* Mesnil,
1950b, Fliegen palaearkt. Reg. 64g: 126.
ZIMBABWE.
Holotype ♂: Salisbury Dist. Dept. Agric. S. Rhodesia
16.VIII.1938/ at flowers of Mango/ A.
Cuthbertson Collector/ Pales maculisquama Mesn.
L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
- Ctenophorocera (Ctenophorocera) macrocephala* Mesnil, 1950b,
Fliegen palaearkt. Reg. 64g: 123. MALAWI.
Holotype ♂: 146 [underside of round label with
"N.NYASA. NYIKA PLATEAU MSWENTI
STRM 28.8.09. J.n.D."/] Pales macrocephala
Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P.
MESNIL COLLECTION 1970.
Remarks: Holotype right front leg is glued to round
paper mount.
- Ctenophorocera (Ctenophorocera) pauciseta* Mesnil, 1950b,
Fliegen palaearkt. Reg. 64g: 125. ZAIRE.
Holotype ♂: NW. Eala 10-XI-1917 R. Mayné/ Pales
pauciseta Mesnil L.P. Mesnil det., 1969/ TYPE/
EX L.-P. MESNIL COLLECTION 1970.
Remarks: The type locality was published as Kisantu.
- Ctenophorocera (Ctenophorocera) tessellans* Mesnil, 1950b,
Fliegen palaearkt. Reg. 64g: 123. SOUTH
AFRICA.
Holotype ♂: Xocavtree 26/9/13/ NATAL FRERE
D.C.M.DIBBEN/ Pales tessellans Mesn. L.P.
Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
- Ctenophorocera (Parapales) pallidula* Mesnil, 1950b, Fliegen
palaearkt. Reg. 64g: 123. MADAGASCAR.
Holotype ♂: Madagascar Bekily- III.30 A.Seyrig/
Parapales pallidula Mesnil L.P. Mesnil det., 1969/
TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Cubaemyiopsis trinitatis* Thompson, 1963b, Studia ent. 6: 386.
TRINIDAD.
Holotype ♀: HOLOTYPE Cubaemyiopsis trinitatis
Thomp. CNC No. 9797/ Santa Cruz V. 3. 5.60
Trinidad,BWI/ 1078 III.5.60/ Cubaemyiopsis
trinitatis nsp.
- Cuphocera trinitatis* Thompson, 1963a, Can. J. Zool. 41: 396.
TRINIDAD.
Syntype: 1♂.
♂: HOLOTYPE Cuphocera trinitatis Thomp.
CNC No. 9775/ 2.17.60 Brazil Trinidad,BWI/
Type ♂/ CNC Syntype Cuphocera trinitatis
Thompson Label affixed 1994.

- Cylindromyia aurora* Herting, 1983, Fliegen palaearkt. Reg. 64c:
54. TUNISIA.
Holotype ♂: Muséum Paris, Tunisie, Maknassy, L.
DUMONT 1929, août/ Exogaster aurora Mesn./
Cylindromyia aurora n.sp., Type, B.Herting det.
- Cylindromyia compressa* Aldrich, 1926, Proc. U.S. natn. Mus. 68:
10. CANADA.
Holotype ♀: Banff, Alta. 23 IX. 1922 C.B.D.Garrett/
Type/ TYPE Cylindromyia compressa Aldrich No.
641.
- Cylindromyia sternalis* Reinhard, 1955d, Ent. News 66: 237.
MEXICO.
Holotype ♂: Ahuacatlan, Nay., Mex. VII-18-22-51/ On
fls. of Donnellsmithia Hintonii M & C/ P. D. Hurd
Collector/ HOLOTYPE Cylindromyia sternalis
Reinhard.
Paratype: 1♂, Mexico.
Remarks: Holotype terminalia are glued to locality
label.
- Cylindromyia (Apinocyptera) thompsoni* Guimarães, 1976, Archos
Zool. S Paulo 27: 28. TRINIDAD.
Holotype ♂: ST.JOSEPH Trinidad,BWI.
NOV.19.1952/ Collector F.J.Simmonds/
HOLOTYPE CYLINDROMYIA THOMPSONI
Guim.
Paratypes: 27♂♂, 10♀♀, Trinidad.
Remarks: Date of collection of holotype was published
as 15 July 1953.
- Cyosoprocta auriceps* Reinhard, 1952b, Bull. Brooklyn ent. Soc.
47: 6. MEXICO.
Holotype ♂: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan
M. R. Wheeler Collectors/ HOLOTYPE
Cyosoprocta auriceps Reinhard.
- Cyosoprocta funebris* Reinhard, 1952b, Bull. Brooklyn ent. Soc.
47: 6. MEXICO.
Holotype ♂: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan
M. R. Wheeler Collectors/ HOLOTYPE
Cyosoprocta funebris Reinhard/ Cyosoprocta
funebris R Rnh.
Allotype ♀: Same data as holotype except with allotype
label and lacking last label.
Paratypes: 4♂♂, 1♀, Mexico.
- Cyrtophloeba nitida* Curran, 1930a, Jl N.Y. ent. Soc. 38: 74.
CANADA.
Holotype ♂: TYPE ♂ Cyrtophloeba nitida 3214 No.
Curran TYPE ♀ Cyrtophloeba nitida 3214 No.
Curran/Cyrtophloeba nitida Det. Curran
C.H.Curran Types 4 para-types./ Aweme, Man. N.
Cridle 21 V 1921.
Allotype ♀: Mounted on same pin as holotype.
Paratypes: 2♀♀, Quebec; 2♀♀, Ontario.
- Descampsina sesamiae* Mesnil, 1956c, Entomophaga 1: 76.
CAMEROON.
Syntype: 1♂.
♂: ex Sesamia Garoua XII. 53 Descamps/
Descampsina sesamiae Mesnil L.P. Mesnil det.,
1969/ COTYPE/ CNC Syntype Descampsina

- Dexodes sesamiae* Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Deuteroclara regalis* Villeneuve, 1915a, Revue zool. afr. 4: 208. MADAGASCAR.
Holotype ♂: Madagascar Tananarive Lamberton 1912/ Deuteroclara regalis Typ. ♂ Villen./ regalis/ EX L.-P. MESNIL COLLECTION 1986.
- Dexia monticola* Villeneuve, 1935b, Revue Zool. Bot. afr. 27: 137. UGANDA.
Holotype ♂: R. [Ruwenzori] 1900m 22.V.14/ Sumatro-Dexia monticola Typ. Villen./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Dexia pollinosa* Villeneuve, 1943b, Revue Zool. Bot. afr. 37: 94. NIGERIA.
Syntype: 1♂.
♂: N. Nigeria. Abinsi R.Benue 1912-1913.
Dr.J.M.Daziel/ Dexia pollinosa Typ. n.sp./ TYPE/ CNC Syntype Dexia pollinosa Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.
- Dexia (Eomyocera) seticincta* Mesnil, 1980, Fliegen palaearkt. Reg. 64f: 46. JAPAN.
Holotype ♂: 20-V-1951 Tanashi-nojyo (Tokyo) I.HATTORI/ 199/ Dexia seticincta Mesn. L.P. Mesnil det., 1979/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Dexiosoma aristatum* Mesnil, 1970, Mushi 44: 118. INDIA.
Holotype ♀: INDE RÉGION HIMALAYENNE KURSÉONG (1500 m ALT.)/ Dexiosoma aristatum Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Dexiotrix pubiseta* Mesnil, 1967, Mushi 41: 54. JAPAN.
Holotype ♀: Hokkaido S.Takano [underside of label] 18/VII'63/ 365/ Dexiotrix pubiseta Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Dexiotrix rufiventris* Mesnil, 1967, Mushi 41: 52. CHINA.
Holotype ♂: Kina S. Kansu/ Sven Hedins Exp. Ctr. Asien Dr Hummel/ Dexiotrix rufiventris Mesn. L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Dexodes conica* Reinhard, 1934b, Bull. Brooklyn ent. Soc. 29: 189. USA.
Holotype ♂: Madison Wis. 6-12 1930/ HOLOTYPE Dexodes conica Reinhard.
- Dexodes sartura* Reinhard, 1959a, Ent. News 70: 234. USA.
Holotype ♂: Babylon, L.I., N.Y. V-26 1935 Blanton & Borders/ HOLOTYPE Dexodes sartura Reinhard.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 1♂, New York; 1♀, Ohio.
- Dexodes valida* Curran, 1927c, Can. Ent. 59: 21. CANADA.
Holotype ♂: Dexodes TYPE valida ♂ Curran No. 2244/ SUDBURY, ONT, CAN.
- Dexodes xylota* Curran, 1927c, Can. Ent. 59: 22. CANADA.
Holotype ♂: Dexodes TYPE xylota Curran. No. 2239/ Orillia, Ont. 7.VI.1925 C.H.Curran.
Allotype ♀: Dexodes AlloTYPE ♀ xylota Curran No 2239./ Orillia,Ont. 11-VI-1925 C.H.Curran.
- Dexodimyia discalis* Thompson, 1968, Mem. ent. Soc. Can. 56: 181. TRINIDAD.
Holotype ♀: HOLOTYPE Dexodimyia discalis Thomp. CNC No. 9883/ N. NEW FARM TRINIDAD 16-9-1964.
- Dexodioida aurea* Thompson, 1968, Mem. ent. Soc. Can. 56: 81. TRINIDAD.
Holotype ♂: TUNAPUNA TRINIDAD 17-8-1964/ HOLOTYPE CNCNo/ Dexodiodea aurea n.g. n.sp./ 1492 X.18.66.
Remarks: Holotype is missing terminalia, and right hind leg is glued to locality label.
- Dexodiopsis aurea* Thompson, 1968, Mem. ent. Soc. Can. 56: 202. TRINIDAD.
Holotype ♂: HOLOTYPE Dexodiopsis aurea Thomp. CNC No. 9886/ CUREPE TRINIDAD 28-9-1964.
Allotype ♀: Same data as holotype except with allotype label.
Remarks: Holotype terminalia are missing.
- Diaprochaeta (Diaprochaeta) illustris* Mesnil, 1970, Mushi 44: 105. ZIMBABWE.
Holotype ♂: Sankishia 3-IX-1911/ Diaprochaeta illustris n.sp. Villen. Diaprochaeta illustris Villen. L.P. Mesnil det., 1969/ TYPE.
- Diaprochaeta (Kaiseriola) aperta* Mesnil, 1970, Mushi 44: 105. SOUTH AFRICA.
Holotype ♂: Durban Marley 4.16/ Diaprochaeta aperta Mesnil L.P. Mesnil det., 1969/ TYPE.
- Dichocera robusta* Brooks, 1945b, Can. Ent. 77: 80. CANADA.
Holotype ♂: Holotype ♂ Dichocera robusta No. Brks 5517/ TABER Alta JUNE 30 1923 H.E.Gray.
Allotype ♀: Orion Alta. SEP 9 1924 H.L.Seamans/ AlloTYPE ♀ Dichocera robusta No. Brks 5517/ SLIDE No. 0439.
Paratypes: 2♂♂, Saskatchewan; 1♂, 1♀, Alberta.
- Diotrepes formosus* Reinhard, 1964b, Acta zool. mex. 7: 8. MEXICO.
Holotype ♂: Cuernavaca, Mex. March 22 1934 S. E. Jones, Coll/ HOLOTYPE Diotrepes formosus Reinhard.
- Dolichocodia erratilis* Reinhard, 1958d, J. Kans. ent. Soc. 31: 278. MEXICO.
Holotype ♂: Teotihuacan, VII-15-36Mex/ HOLOTYPE Dolichocodia erratilis Reinhard/ Dolichocodia erratilis R Rhn.
- Dolichocodia furacis* Reinhard, 1958d, J. Kans. ent. Soc. 31: 277. USA.
Holotype ♂: Hidalgo County 8-19 1931 Tex/ H J Reinhard Collector/ HOLOTYPE Dolichocodia furacis Reinhard.
Paratypes: 2♂♂, Texas.

- Dolichocolon africanum* Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 176. ZAIRE.
Holotype ♂: Congo belge : P.N.A. Rwindi 1000m. 26-xi-1934 G.F.de Witte : 818/ Dolichocolon africanum Mesnil L.P. Mesnil det., 1967/ TYPE/ HOLOTYPE ♂ of Dolichocolon africanum Mesnil Examined 1970 R.W.Crosskey.
- Dolichocolon klapperichi* Mesnil, 1967, Mushi 41: 43. CHINA.
Holotype ♂: Kwangtseh-Fukien J.Klapperich O 9.10 1937/ Dolichocolon klapperichi Mesn. L.P. Mesnil det., 1967/ TYPE/ HOLOTYPE ♂ of Dolichocolon klapperichi Mesnil Examined 1970 R.W.Crosskey.
- Dolichocolon vicinum* Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 176. VIETNAM.
Holotype ♂: INDOCHINE FRANÇAISE Nhatranget Saigon A. KREMP 1913/ Dolichocolon vicinum Mesnil L.P. Mesnil det., 1967/ TYPE/ HOLOTYPE ♂ of Dolichocolon vicinum Mesn. Examined 1970 R.W.Crosskey.
Paratype: 1♀, Japan.
- Dolichopodomintho takanoi* Mesnil, 1973a, Fliegen palaearkt. Reg. 64g: 1161. JAPAN.
Holotype ♂: Utatsuyama Kanazawa JPN. 5.X.'64 S.Takano/ 416/ Dolichopodomintho takanoi Mesnil L.P. Mesnil det., 1971/ TYPE/ EX. L.-P. MESNIL COLLECTION 1985.
- Dolichotarsus griseus* Brooks, 1945b, Can. Ent. 77: 95. CANADA.
Holotype ♀: Trinity Valley 16.VI.37 B.C. Hugh B. Leech/ Holotype ♀ Dolichotarsus griseus No. Brks 5531.
- Dolichotarsus kingi* Brooks, 1945b, Can. Ent. 77: 95. CANADA.
Holotype ♂: 16446 114AEI/ Saskatoon, Sask. June 8 1927 Kenneth M. King/ [pink blank label]/ HoloTYPE ♂ Dolichotarsus kingi No. Brks. 5530.
Allotype ♀: Cut Knife, Sask. 7-VI-1940 A.R.Brooks/ AlloTYPE ♀ Dolichotarsus kingi Brks. No. 5530.
Paratypes: 1♂, 1♀, Saskatchewan.
Remarks: Holotype puparium is glued to a paper point below specimen.
- Dolichotarsus livescens* Reinhard, 1958b, J. Kans. ent. Soc. 31: 226. USA.
Holotype ♀: Milagro, N.M. VI-23-41/ HOLOTYPE Dolichotarsis [sic] livescens Reinhard/ Dolichotarsis [sic] livescens R Rnh.
- Dolichotarsus stipatus* Reinhard, 1958b, J. Kans. ent. Soc. 31: 225. MEXICO.
Holotype ♀: 9600ft. W.slope Popocatepetl, Mex. VII-5-51/ P. D. Hurd/ HOLOTYPE Dolichotarsis [sic] stipatus Reinhard/ Dolichotarsis [sic] stipatus R Rnh.
Paratype: 1♀, Mexico.
- Doliolomyia alactaga* Reinhard, 1974, Can. Ent. 106: 1160. MEXICO.
Holotype ♂: Jalapa, Ver. 9/28-X/3/61 R&K Dreisbach/ HOLOTYPE Doliolomyia alactaga Reinhard.

- Doliolomyia thessa* Reinhard, 1974, Can. Ent. 106: 1160. MEXICO.
Holotype ♂: Jalapa, Ver. 9/28-X/3/61 R&K Dreisbach/ HOLOTYPE Doliolomyia thessa Reinhard.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 9♂♂, Mexico.
- Doryphorophaga australis* Reinhard, 1935a, Jl N.Y. ent. Soc. 43: 389. USA.
Lectotype ♂: College Station 6-14-19 Tex/ R.J.Reinhard Collector/ HOLOTYPE Doryphorophaga australis Reinhard/ CNC Syntype Doryphorophaga australis Reinhard Label affixed 1994/ Lectotype fixation by Wood, 1985: 63. Label affixed 1995.
Paratypes: 33♂♂, 4♀♀, Texas.
Remarks: We regard Wood's (1985: 63) reference to a holotype as a lectotype fixation for this species.
- Doryphorophaga macella* Reinhard, 1935a, Jl N.Y. ent. Soc. 43: 390. USA.
Holotype ♂: College Station Oct.27,1934 Tex/ H J Reinhard Collector/ HOLOTYPE Doryphorophaga macella Reinhard/ Doryphorophaga macella det.H.J.Reinhard Rein.
Paratypes: 1♂, 1♀, Ohio; 2♂♂, 14♀♀, Texas.
- Doryphorophaga patrita* Reinhard, 1935a, Jl N.Y. ent. Soc. 43: 394. USA.
Syntypes: 9♀♀.
9♀♀: Amherst, Ohio [various dates July-August 1933-35]/ H J Reinhard Collector/ PARATYPE Doryphorophaga patrita Reinhard/ CNC Syntype Doryphorophaga patrita Reinhard Label affixed 1994.
Remarks: Though Wood (1985: 66) referred to a female holotype, no holotype was designated or labeled by Reinhard.
- Doryphorophaga sedula* Reinhard, 1935a, Jl N.Y. ent. Soc. 43: 392. USA.
Holotype ♂: Amherst, Ohio 7-17 1934/ H J Reinhard Collector/ HOLOTYPE Doryphorophaga sedula Reinhard/ Doryphorophaga sedula det.H.J.Reinhard Rein.
Paratypes: 9♂♂, 1♀, Ohio; 2♂♂, Texas.
- Drino aureocauda* Thompson, 1966, Studia ent. 8 (1965): 398. TRINIDAD.
Holotype ♂: HOLOTYPE Drino aureocauda Thomp. CNC No. 9833/ SAN JUAN Trinidad,BWI. OCT.13.1952./ Collector F.J.Simmonds/ Drino aureocauda n.sp.
- Drino curepei* Thompson, 1966, Studia ent. 8 (1965): 408. TRINIDAD.
Holotype ♀: HOLOTYPE Drino curepei Thomp. CNC No.9836/ PARADISE MT. Trinidad,BWI. FEB.19.1953./ Collector F.J.Simmonds/ Analcombs m. [word illegible] large/ Drino curepei n.sp/ 925/ 88.
Allotype ♂: ALLOTYPE Drino curepei Thomp. CNC No.9836/ Cumuto TRINIDAD 21-7-1964/ 1440 XI.17.64 [underside of label] Trinidad, WI.

Remarks: Date of collection of holotype was published as 14 February 1953. Allotype terminalia are missing.

Drino nigricauda Thompson, 1966, Studia ent. 8 (1965): 394.
TRINIDAD.

Holotype ♂: HOLOTYPE Drino nigricauda Thomp. CNC No.9832/ Trinidad,BWI. Bamboo Gr. 2 X.26.60.

Allotype ♀: Arima V. TRINIDAD W.I. III. 9. 1961 W.R.Thompson/ ALLOTYPE Drino nigricauda Thomp. CNC No. 9832.

Drino nigripalpis Thompson, 1966, Studia ent. 8 (1965): 404.
TRINIDAD.

Holotype ♀: HOLOTYPE Drino nigripalpis Thomp. CNC No. 9835/ St. Augustine, Trinidad,BWI. I. 16. 60/ 1038 A-7 17. I. 60 [underside of label] St. Augustine, Trinidad,BWI.

Paratypes: 2♀, Trinidad.

Drino ruficauda Thompson, 1966, Studia ent. 8 (1965): 401.
TRINIDAD.

Holotype ♂: HOLOTYPE Drino ruficauda Thomp. CNC No.9834/ Trinidad,BWI. Bamboo Grove X.26.60/2 Drino ruficauda n.sp.

Allotype ♀: ALLOTYPE Drino ruficauda Thomp. CNC No.9834/ St. Augustine, Trinidad,BWI. I 19. 60/ ♀: p black abd.n. [word illegible].

Drino trinidadensis Thompson, 1966, Studia ent. 8 (1965): 411.
TRINIDAD.

Holotype ♂: HOLOTYPE Drino trinidadensis Thomp. CNC No. 9837/ CUREPE Trinidad,BWI. OCT.6.1952./ Collector F.D. Bennett/ Drino trinidadensis nsp.

Allotype ♀: ALLOTYPE Drino trinidadensis Thomp. CNC No. 9837/ 1058 A2 I. 17. 60/ St. Augustine, Trinidad,BWI. I. 16. 60/ Larvae with anal combs/ Anal combs present M. [word illegible].

Drino (Isosturmia) chatterjeeana japonica Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 13. JAPAN.

Holotype ♂: Obihiro S.Takano [underside of label] 23/VI'54/ 10/ Isosturmia trisetosoides Bar. L.P. Mesnil det., 1967/ subsp. japonica Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ Isosturmia japonica (Mesn.) det. H. Shima, 1986.

Remarks: Date of collection of holotype was published as 6 July 1952.

Drino (Prosturmia) amicula Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 30. MOZAMBIQUE.

Syntype: 1♂.

♂: MAI/ ZAMBEZE AMONT DE TAMBARA ALFIATE (GARÉ) P. LESNE 1929/ Drino amicula Mesnil L.P. Mesnil det., 1969/ COTYPE/ CNC Syntype Drino (Prosturmia) amicula Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Drino (Prosturmia) angustifacies Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 23. JAPAN.

Holotype ♂: Prosturmia TYPE angustifacies Mesn. No. 5826/ 18146H37 300 a 15.VII. 38/ Host Neodiprion sertifer/ Kyushu Japan/ 3684/ Prosturmia angustifacies Mesn. L. Mesnil det., 1949/ TYPE.

Paratypes: 3♂, 5♀, Japan.

Drino (Prosturmia) aurora Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 15. JAPAN.

Holotype ♂: Prosturmia TYPE gilva s.sp. aurora Mesn No. 5827/ Japan/ 18165H35 300 a 28.IX.36/ Host Diprion nipponica/ 3709/ Prosturmia aurora Mesn. L. Mesnil det., 1949/ TYPE.

Remarks: There may be six paratypes among the CNC material but they are not labeled as such.

Drino (Prosturmia) bohemica Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 23. SWEDEN.

Holotype ♂: 18111 H38 300 a-2 4.VIII.39/ Host G. hercynia Htg/ 4759/ TYPE Prosturmia bohemica Mesn No. 5824/ Sturmia sp. ♂/ TYPE/ Prosturmia bohemica Mesn. L. Mesnil det 1949/ Vittangi Sweden.

Remarks: Holotype puparium is stored in a gelatin capsule that is pinned with the specimen. Host pupal case is also pinned with the specimen. Mesnil (1949b: 23) erroneously cited the type locality as "Nova Bistrice, en Bohême" (see explanation by Crosskey 1966: 135).

Drino (Prosturmia) gilpiniae Mesnil, 1971a, Entomophaga 16: 67. PAKISTAN.

Holotype ♂: Neelam Valley 7. V. 70/ Ex. Gilpinia sp cocoon/ 801/ Drino gilpiniae Mesnil L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1985/ TYPE.

Remarks: Holotype terminalia are glued to a paper strip below specimen.

Drino (Prosturmia) inconspicua var. *inusta* Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 25. POLAND.

Holotype ♂: Prosturmia TYPE inconspicua v. inusta Mesn. No.5825/ Czecho-slovakia/ 18111 H37 300 www4 20.VIII.38/ Host Gilpinia hercyniae/ 3696/ Prosturmia inconspicua Mg. var. inusta Mesn. L. Mesnil det., 1949/ TYPE.

Remarks: Holotype puparium is glued to a paper strip below specimen. Country of origin was published as Poland, though the holotype bears a "Czechoslovakia" label.

Drino (Prosturmia) iterata Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 31. UGANDA.

Syntype: 1♂.

♂: Uganda/ Drino iterata Mesnil L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Drino (Prosturmia) iterata Mesnil Label affixed 1994.

Drino (Prosturmia) latigena maroccana Mesnil, 1951, Fliegen palaearkt. Reg. 64g: 194. MOROCCO.

Holotype ♂: 1246 Maroc 5a lé exl 7.XII.34 Ch. Rungs/ ExTaragama repandum Hbn. (Lariocamp)/ TYPE/ Drino maroccana Mesnil L.P. Mesnil det., 1969.

Drino (Prosturmia) melancholica Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 16. ZIMBABWE.

Syntypes: 1♂, 1♀.

♂: Salisbury S. Rhodesia. Dept. Agric. 21 March, 1931./ on cotton plants./ Drino melancholica Mesn. L.P. Mesnil det., 1969/ SYN-TYPE/ TYPE/ Syntype/ CNC Syntype
Drino (Prosturmia) melancholica Mesnil Label affixed 1994.
♀: Salisbury S. Rhodesia. Dept. Agric. 21 March 1931/ Drino melancholica Mesn L.P. Mesnil det., 1963/ CNC Syntype Drino (Prosturmia) melancholica Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Drino (Prosturmia) nova Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 27. MADAGASCAR.

Syntype: 1♂.

♂: MADAGASCAR BEKILY RÉG SUD DE L'ILE/ Drino nova Mesnil L.P. Mesnil det., 1969/ CNC Syntype Drino (Prosturmia) nova Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970/ COTYPE.

Drino (Prosturmia) patruelis Mesnil, 1949b, Bull. Inst. r. Sci. nat. Belg. 25(42): 17. ZIMBABWE.

Syntype: 1♂.

♂: C.1083 Salisbury S. Rhodesia. Dept. Agric. 30.IV. 193.8/ Coll. 8591/ A.Cuthbertson Collector./ Drino patruelis Mesn. L.P. Mesnil det., 19/ TYPE/ SYN-TYPE/ CNC Syntype Drino (Prosturmia) patruelis Mesnil Label affixed 1994.

Drino (Prosturmina) aureola Mesnil, 1970, Mushi 44: 110. SIERRA LEONE.

Holotype ♂: BAFODEA Sierra Leone, 14.IV. 1912 Jas. J. Simpson/ Drino aureola Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Dufouria nova Mesnil, in Mesnil & Pschorr-Walcher, 1968, Mushi 41: 174. JAPAN.

Holotype ♂: Mt. Teine N. Sapporo Japan 10. VII. 1961/ Dufouria nova Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Echinomyia genurufa Villeneuve, 1936e, Ark. Zool. 27A: 4. CHINA.

Holotype ♀: Kina S. Kansu/ Sven Hedins Exp. Ctr. Aslen Dr Hummel/ 23|8/ Echinomyia genurufa ♀ n.sp./ Tachina genurufa Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ Tachina corsicana Villen. L.P. Mesnil det., 1970.

Echinomyia grandigena Villeneuve, 1936d, Bull. Mus. r. Hist. nat. Belg. 12(42): 6. BHUTAN.

Holotype ♀: British Bootan Padong. L. Durel 1913./ Echinomyia grandigena Villen. ♀ Typ./ Mikia grandigena Villen. L.P. Mesnil det., 1970/ TYPE/ Mikia tepens Walker L.P. Mesnil det., 1970/ HOLOTYPE ♀ of Echinomyia grandigena Villen. Examined 1970 R.W.Crosskey.

Echinomyia hedini Villeneuve, 1936e, Ark. Zool. 27A: 3. CHINA.

Lectotype ♂: Kina S. Kansu/ 27/6/ Sven Hedins Exp. Ctr. Asien Dr Hummel/ Eudoromyia Hedinsi [sic] Typ. Villen./ TYPE/ Nowickia polita Zimin L.P. Mesnil det., 1970/ Nowickia hedini Villen. L.P. Mesnil det., 1970/ Eudoromyia hedini Villeneuve LECTOTYPE ♂ designated by R.W.Crosskey 1970/ LECTOTYPE ♂ Eudoromyia hedini Villeneuve Designated by R.W.Crosskey, 1974.

Remarks: For lectotype designation see Crosskey (1976: 269). Date of collection of Lectotype was published as 27 August. There may be a paralectotype among the CNC material but it was not labeled as such.

Echinomyia magnicornis var. *corsicana* Villeneuve, 1931, Konowia 10: 48. ALGERIA.

Syntype: 1♂.

♂: Oasis Gafsa Tunis B.v.Bodemeyer/ Echinomyia corsicana Villen. =prompta Macq./ Tachina corsicana Vill. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype Echinomyia magnicornis var. corsicana Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Echinomyia splendidula Villeneuve, 1936d, Bull. Mus. r. Hist. nat. Belg. 12(42): 6. VIETNAM.

Holotype ♂: Hanoi 4376/ Echinomyia splendidula n.sp. Typ. Villen./ Mikia splendidula Villen L.P. Mesnil det., 1970/ TYPE/ Mikia tepens Walker L.P. Mesnil det., 1970/ HOLOTYPE ♂ of Echinomyia splendidula Villen. Examined 1970 R.W.Crosskey.

Echinomyia trigonata Villeneuve, 1936e, Ark. Zool. 27A: 3. CHINA.

Holotype ♂: Kina S. Kansu/ Sven Hedins Exp. Ctr. Asien Dr. Hummel/ 30.10.1930/ Eudoromyia trigonata Typ. ♂ Villen./ TYPE/ Tachina trigonata Vill. L.P. Mesnil det., 1970/ HOLOTYPE ♂ of Eudoromyia trigonata Villen. Examined 1970 R.W.Crosskey.

Echinomyia (Larvaevora) rubrapex Villeneuve, 1932b, Bull. Soc. ent. Fr. 37: 268. TAIWAN.

Lectotype ♀: LECTO-TYPE/ Formosa Sauter/ Polisha 908.XII./ Echinomyia rubrapex Typ. Villen./ Mikia aureocephala Towns L.P. Mesnil det., 1970/ TYPE/ Mikia rubrapex Villen. L.P. Mesnil det., 1970/ Echinomyia rubrapex Vill. LECTOTYPE ♀ designated by R.W.Crosskey 1971/ LECTOTYPE ♀ Echinomyia rubrapex Villeneuve Designated by R.W. Crosskey, 1974.

Remarks: For lectotype designation see Crosskey (1976: 267).

Echinopyrrhosia arrogans Reinhard, 1974, Can. Ent. 106: 1160. COLOMBIA.

Holotype ♂: PRMO GUERRERQ CUNDINAMARCA COLOMBIA ALT. 3200 NI.K.67/ [word illegible] OR H. OSORNO FECHA 25-VII-48/ HOLOTYPE Echinopyrrhosia arrogans Reinhard.

- Echinopyrrhosia pellacis* Reinhard, 1974, Can. Ent. 106: 1161.
COLOMBIA.
Holotype ♀: PRMO GUERRERQ CUNDINAMARCA COLOMBIA ALT. 3200 NL.K.67/ HOLOTYPE Echinopyrrhosia pellacis Reinhard/
Echinopyrrhosia pellacis R Rnh.
- Eipogona americana* Curran, 1927a, Bull. Brooklyn ent. Soc. 22: 148. USA.
Holotype ♂: SiouxCity Ia Aug '20/ CNAinalie Collector/ Reared from Papilio asterias/ SiouxCity No 2063/ Eipogona TYPE americana ♂ Curran No. 2237.
- Elassomyia defecta* Reinhard, 1967, J. Kans. ent. Soc. 40: 103. USA.
Holotype ♂: Calif/ HOLOTYPE Elassomyia defecta Reinhard/ Elassomyia defecta R Rnh.
- Eleodiphaga martini* Reinhard, 1937, Bull. Brooklyn ent. Soc. 32: 69. USA.
Holotype ♂: Parma,Ida. IX.4.1934/ Chas.H.Martin Col./ HOLOTYPE Eleodiphaga martini Reinhard/
Eleodiphaga martini det.H.J.Reinhard Rein.
- Elfia viridis* Reinhard, 1967, J. Kans. ent. Soc. 40: 104.
COLOMBIA.
Holotype ♂: Pasto Nar 8-V-58/ A. UNIGARRO PAPA/ HOLOTYPE Elfia viridis Reinhard.
Allotype ♀: Pasto Nar. 8-V-58/ A. UNIGARRO PAPA./ ALLOTYPHOE Elfia viridis Reinhard/ Elfia viridis R Rnh.
Paratypes: 10♀, Colombia.
- Elfriedella amoena* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 69. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label] 2-3/VII'49/ 94/ Elfriedella amoena Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Eloceria delecta grandis* Mesnil, 1973b, Fliegen palaearkt. Reg. 64g: 1226. SWITZERLAND.
Holotype ♂: Delmont 26. VII. 1972/ Eloceria grandis Mesn. L.P. Mesnil det., 1972/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
Paratypes: 1♂, 5♀, Switzerland.
- Elodimyia tricincta* Mesnil, 1952b, Fliegen palaearkt. Reg. 64g: 243. INDONESIA.
Lectotype ♂: Sunda Exp. Rensch Nlawangan Lombok 3. IV. 1927/ Elodimyia tricincta Mesnil L.P. Mesnil det., 1967/ TYPE/ HOLOTYPE ♂ of Elodimyia tricincta Mesnil Examined 1970 R.W.Crosskey/ LECTOTYPE ♂ Elodimyia tricincta Mesnil Designated by R.W.Crosskey, 1974.
Remarks: For lectotype designation see Crosskey (1976: 268).
- Enchomyia shewelli* Cortés, 1975, Boln tēc Estac. exp. agron. Fac. Agronomía Univ. Chile 40: 5. CHILE.
Holotype ♂: ♂/ HOLOTIPO/ Banos El Toro Coquimbo,CHILE 3300-4000 m. 6-12.I.66,Pena/

- Enchomyia n.sp. Det.D.M.Wood 1969/
Enchomyia ♂ shewelli n.sp. R.Cortés det. 1974.
- Eomeigenielloides segestris* Reinhard, 1974, Can. Ent. 106: 1162. MEXICO.
Holotype ♀: Cuernavaca, Mor., Mex. 7-29-61 7000' R & K Dreisbach/ HOLOTYPE Eomeigenielloides segestris Reinhard.
- Epidexia mimela* Reinhard, 1953b, J. Kans. ent. Soc. 26: 96. USA.
Holotype ♂: College Station Jun.16.1947 Tex/ H J Reinhard Collector/ HOLOTYPE Epidexia mimela Reinhard.
Paratypes: 2♂♂, Texas; 6♂♂, Mexico.
- Epineura minor* Villeneuve, 1913, Revue zool. afr. 3: 45. ZAIRE.
Syntype: 1♂.
♂: Mt. Mlanje, Nyasaland, 10.X.1913. S. A. Neave./ ♀/ Epineura minor ♂ Villen. Dr VILLENEUVE det. 1899/ TYPE/ CNC Syntype Epineura minor Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.
Remarks: Syntype is missing abdomen.
- Epineura rubens* Villeneuve, 1923b, Revue zool. afr. 11: 78. NIGERIA.
Syntype: 1♂.
♂: Bende. S.Nigeria. J.J.Simpson. 22.V.1910/ ♂/ Bogosia (Epineura) rubens V./ TYPE/ CNC Syntype Epineura rubens Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.
- Ergolabus bonasus* Reinhard, 1964b, Acta zool. mex. 7: 16. MEXICO.
Holotype ♀: Cuernavaca, Mex March 22 1934 S. E. Jones, Coll/ HOLOTYPE Ergolabus bonasus Reinhard/ HOMOTYPE Heliae mirabilis Cn. Compared by Wood 73/ HOMOTYPE Plagia setifrons Wlp Compared by Wood 89.
- Erigone cristata* Villeneuve, 1920b, Annls Soc. ent. Belg. 60: 116. FRANCE.
Syntype: 1♂.
♂: Montloruis 22. VI/ Eurithia cristata Villen. L.P. Mesnil det., 1970/ Paratype/ CNC Syntype Erigone cristata Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Eriothrix umbrinervis* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 68. JAPAN.
Holotype ♀: Obihiro S.Takano [underside of label] 16/IX'50/ 28/ Eriothrix umbrinervis Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Ernestia nigropalpis* Tothill, 1921, Can. Ent. 53: 247. CANADA.
Holotype ♂: TYPE No.483/ Stickeen R.Can. B. C. Wickham./ TypeNo. 24354 U.S.N.M./ Ernestia nigropalpis Tothill.
Paratypes: 5♂♂, British Columbia.
- Ernestia (Faustia) fasciata* Curran, 1924a, Ent. News 35: 246. CANADA.

Holotype ♂: HoloTYPE Ernestia fasciata Curr No. 796/ Cranbrook,B.C. 8.V 1922 C.B.D.Garrett.
 Allotype ♀: TYPE E. fasciata Curran No. 796/ Ok.Falls B.C. 24. IV. 1919 E.R. Buckell.
 Paratypes: 1♂, 1♀, British Columbia.
 Remarks: Holotype terminalia are glued to locality label. Date of collection of holotype was published as 9 May 1922. CNC type number was published as 79.

Erycia aldrichi Curran, 1927c, Can. Ent. 59: 17. CANADA.
 Holotype ♂: Erycia TYPE ♂ aldrichi Curran No. 2240/ Hedley, B.C. 29.VIII.1923 C.B.Garrett/ ♂.
 Allotype ♀: Same data as holotype except with allotype label.
 Paratypes: 5♂♂, 1♀, British Columbia.

Erycia delecta Curran, 1927c, Can. Ent. 59: 16. CANADA.
 Holotype ♂: Erycia TYPE delecta Curran. No. 2243./ Winnipeg,Man. A.V.Mitchener July 6 1925.

Erycia levata Reinhard, 1934b, Bull. Brooklyn ent. Soc. 29: 193. USA.
 Holotype ♂: California May 19, 1927./ HOLOTYPE Erycia levata Reinhard/ Erycia levata det.H.J.Reinhard Rein/ Erycia occidentalis det.H.J.Reinhard Coq.
 Paratypes: 2♂♂, Washington; 1♂, Oregon.

Erycia sectilis Reinhard, 1953c, Pan-Pacif. Ent. 29: 59. USA.
 Holotype ♂: Teton Pass, Mont Aug. 2, 1947/ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE Erycia sectilis Reinhard/ Sisyrosturmia sectilis R (Rnh.).
 Paratype: 1♂, Wyoming.
 Remarks: The locality label of holotype and paratype indicates Teton Pass in the state of Montana, whereas the original description gives it in Wyoming.

Erycia varifrons Curran, 1927c, Can. Ent. 59: 18. CANADA.
 Holotype ♂: Erycia TYPE ♂ varifrons Curran ♀ No. 2241./ Lillooet,B.C. 27-V 1925 E.R.Buckell.
 Allotype ♀: Mounted on the same pin as holotype.
 Paratype: 1♂, British Columbia.

Erycilla amoena Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 20. JAPAN.
 Holotype ♀: Obihiro S.Takano [underside of label] 5/VIII '50/ 58/ Erycilla amoena Mesnil L.P. Mesnil det., 1969/ TYPE.

Erynnia condecor Reinhard, 1968b, J. Kans. ent. Soc. 41: 507. CANADA.
 Holotype ♂: Summit Lake,B.C. Mi392 Alaska Hwy 17-19. VII. 1959 4500' R.E. Leech/ HOLOTYPE Erynnia condecor Reinhard.
 Paratypes: 2♂♂, California.

Erynnia coracina Reinhard, 1968b, J. Kans. ent. Soc. 41: 504. CANADA.
 Holotype ♂: Whitehorse, Y.T. 24.VIII.1959/ HOLOTYPE Erynnia coracina Reinhard/ Reinhard Collection.
 Paratypes: 1♂, 1♀, British Columbia.

Erynnia tricincta Reinhard, 1968b, J. Kans. ent. Soc. 41: 506. MEXICO.

Holotype ♂: TLAHUALILO DGO MEX 8-3-'36/ 1650/ C. S. RUDE COLL./ HOLOTYPE Erynnia tricincta Reinhard.
 Allotype ♀: Same data as holotype except with allotype label.

Eubrachymera florentis Reinhard, 1964a, J. Kans. ent. Soc. 37: 40. USA.

Holotype ♂: HOLOTYPE Eubrachymera florentis Reinhard/ Glacier Nat. Pk. July 27, 1947.
 Allotype ♀: Same data as holotype except with allotype label.

Eucarcelia nudicauda Mesnil, 1967, Mushi 41: 37. JAPAN.

Holotype ♀: Mt.Horaiji Aichi Pref. 7.VII.1960 S.Takano/ 412/ Eucarcelia nudicauda Mesn L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Eucarcelia villeneuvei Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 3. HUNGARY.

Holotype ♂: Ungarn 53345./ Eucarcelia villeneuvei Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Eucelatoria teutonia Sabrosky, 1981, Tech. Bull. U.S. Dep. Agric. 1635: 14. BRAZIL.

Holotype ♂: Nova Teutonia 27°11'S 52°23'W Brazil, 300-500m. II-1965 Fritz Plaumann/ HOLOTYPE Eucelatoria ♂ teutonia C.W.Sabrosky/ Eucelatoria teutonia det SABR. Sabrosky.
 Paratypes: 35♂♂, Brazil.

Eucelatorioida nigripalpis Thompson, 1968, Mem. ent. Soc. Can. 56: 177. TRINIDAD.

Holotype ♀: HOLOTYPE Eucelatorioida nigripalpis Thomp. CNC No. 9881/ Trinidad,BWI./ Eucelatorioida nigripalpis.

Eucelatorioida physonotae Thompson, 1968, Mem. ent. Soc. Can. 56: 179. TRINIDAD.

Lectotype ♀: Caroni Trinidad,BWI. Nov. 1946/ Collector F.J.Simmonds/ LECTOTYPE ♀ of Eucelatorioida physonotae Thompson designated 1984 D.M.Wood.
 Paralectotypes: 3♂♂, 3♀♀, Trinidad.
 Remarks: For lectotype designation see Wood (1985: 98). Lectotype puparium is pinned below specimen.

Euceromasia neptis Reinhard, 1947, J. Kans. ent. Soc. 20: 23. USA.

Holotype ♀: College Station Jun.23,1944 Tex/ H J Reinhard Collector/ HOLOTYPE Euceromasia neptis Reinhard/ Euceromasia neptis R. Rnh.

Paratypes: 1♂, 31♀♀, Texas.
 Remarks: The male labeled as a paratype may actually be the allotype.

Euceromasia sobrina Reinhard, 1974, Can. Ent. 106: 1162.

CANADA.
 Holotype ♂: Black Sturgeon Lake, Ontario Em. 10-VII-63 Insectary/ Club top4 Picea mariana R'd/ HOLOTYPE Euceromasia sobrina Reinhard.

Allotype ♀: Black Sturgeon Lake, Ontario Em.
26-VII-62 Insectary/ Club top7 Picea mariana R'rd
ALLOTYPE Euceromasia sobrina Reinhard.
Paratypes: 14♂♂, 4♀♀, Ontario.

Euceromasia solata Reinhard, 1947, J. Kans. ent. Soc. 20: 24.
USA.

Holotype ♂: College Station May 2,1944 Tex/ H J
Reinhard Collector/ HOLOTYPE Euceromasia
solata Reinhard.
Allotype ♀: Same data as holotype except with allotype
label and collected "Jun 27,1944".
Paratypes: 2♂♂, 2♀♀, Texas.

Eufrontina ethniae Brooks, 1945b, Can. Ent. 77: 91. CANADA.
Holotype ♂: C.E.Farm Ottawa, Ont. emerged in lab.

4.II.1933 J.McDunnough/ Lep.bred on
Lithospermum officinale/ Host. Ethnia
longimaculella Clem./ HoloTYPE ♂ Eufrontina
ethniae No. Brks 5526.

Allotype ♀: Same data as holotype except with allotype
label and emergence date "Feb.27,1933", and one
additional label "etl".

Paratypes: 7♂♂, 2♀♀, Ontario.

Remarks: Two puparia on pin below holotype, and one
puparium on pin below allotype. There are 7♂♂
and 2♀♀ paratypes (not 5♂♂ and 4♀♀ as
published), see O'Hara (1993: 26).

Eugaediopsis rubentis Reinhard, 1961, J. Kans. ent. Soc. 34: 209.
USA.

Holotype ♂: Chiricahua Mts. July30,1957 ARIZ. G. D.
Butler/ HOLOTYPE Eugaediopsis rubentis
Reinhard/ Eugaediopsis rubentis R Rnh.

Allotype ♀: Same data as holotype except with allotype
label.

Paratypes: 2♂♂, 1♀, Arizona; 2♂♂, Mexico.

Eugaediopsis simmondsi Thompson, 1963b, Studia ent. 6: 283.
TRINIDAD.

Syntype: 1♂.

♂: St. Augustine, Trinidad,BWI. II.7.60/ ♂ Type/
HOLOTYPE Eugaediopsis simmondsi Thomp.
CNC No. 9786/ CNC Syntype Eugaediopsis
simmondsi Thompson Label affixed 1994.

Eugaediopsis vinnula Reinhard, 1961, J. Kans. ent. Soc. 34: 210.
USA.

Holotype ♂: Sunnyside Canyon Huachuca Mts. Ariz.
VII.9.40 R. H. Beamer/ HOLOTYPE Eugaediopsis
vinnula Reinhard.

Euhystricia cussiliris Reinhard, 1953a, Bull. Brooklyn ent. Soc. 48:
91. MEXICO.

Holotype ♂: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan
M. R. Wheeler Collectors/ HOLOTYPE
Euhystricia cussiliris Reinhard/ Euhystricia
cussiliris R. Rnh.

Eulasiona nigra Curran, 1924c, Can. Ent. 56: 194. CANADA.

Holotype ♂: HOLOTYPE Eulasiona nigra Curr No.
705/ Victoria,B.C. 5.VII.1923 K.F.Auden/ Host.
Tmetocera and Cacoecia.

Allotype ♀: TYPE E. nigra Curr No. 705./
Victoria,B.C. 5.VII.1923 K.F.Auden/ 10575/ Host.
Tmetocera and Cacoecia.
Paratypes: 1♂, 1♀, British Columbia.

Eulasiona tibialis Curran, 1927a, Bull. Brooklyn ent. Soc. 22: 150.
CANADA.

Holotype ♀: Eulasiona TYPE tibialis Curran No. 2057./
Aylmer,Que 24.V. 1924 C.H.Curran/ Comp. w.
type of E. spinosa Coq. and seems to me to be a
different sp. GES. (see notes) '53.

Eumacrohoughia minor Thompson, 1963b, Studia ent. 6: 333.
TRINIDAD.

Syntypes: 1♂, 1♀.

♂: HOLOTYPE Eumacrohoughia minor Thomp.
CNC No. 9791/ Brazil II.18.61. Trinidad,BWI/
CNC Syntype Eumacrohoughia minor Thompson
Label affixed 1994.2

♀: ALLOTYPE Eumacrohoughia minor Thomp.
CNC No. 9791/ Arima V. TRINIDAD W.I. III. 9.
1961 W.R.Thompson/ 19.01.III 2911/ CNC
Syntype Eumacrohoughia minor Thompson Label
affixed 1994.

Euphorocera cumutoensis Thompson, 1968, Mem. ent. Soc. Can.
56: 16. TRINIDAD.

Holotype ♂: HOLOTYPE Euphorocera cumutoensis
Thomp. CNC No. 9841/ II 20 64 Trinidad, WI.

Allotype ♀: ALLOTYPE Euphorocera cumutoensis
Thomp. CNC No. 9841/ TRINIDAD 28-7-1964
CUMUTO.

Paratype: 1♂, Trinidad.

Eupododexia diaphana Villeneuve, 1915a, Revue zool. afr. 4: 202.
MADAGASCAR.

Holotype ♂: Madagascar Tananarive Lambert 1912/
Eupododexia diaphana Typ. Villen. ♂/ TYPE/ EX
L.-P. MESNIL COLLECTION 1985.

Eupododexia festiva Villeneuve, 1915a, Revue zool. afr. 4: 201.
MADAGASCAR.

Syntype: 1♂.

♂: Mdkg. Sikora Andrangoloaka/ Eupododexia
festiva Typ. Villen. ♂/ TYPE/ CNC Syntype
Eupododexia festiva Villeneuve Label affixed
1994/ EX L.-P. MESNIL COLLECTION 1985.

Remarks: Syntype right wing is glued to locality label.

Euptilopareia vicinalis Reinhard, 1956c, Ent. News 67: 125. USA.
Holotype ♂: SEQUOIA N.P. 30-VII-39 CAL./

HOLOTYPE Euptilopareia vicinalis Reinhard.

Allotype ♀: Same data as holotype except with allotype
label.

Paratype: 1♂, California.

Eurylochus dissitus Reinhard, 1967, J. Kans. ent. Soc. 40: 109.
MEXICO.

Holotype ♂: Jalapa, Ver. 9/28-E/3/61 R & K Dreisbach/
HOLOTYPE Eurylochus dissitus Reinhard.

Allotype ♀: Same data as holotype except with allotype
label.

Paratypes: 2♂♂, Mexico.

Eurysthaea (Discochaeta) veniseta Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 181. PAKISTAN.

Holotype ♀: Ex.larva feedin/ g leaves of *Salix* sp./ [word illegible] 4.VII.63/ C.I.B.C FPO.7163-93/ 1333/ *Eurysthaea veniseta* Mesnil L.P. Mesnil det., 1967/ TYPE.

Remarks: Holotype puparium is pinned below specimen.

Eurythia emdeni Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 58. JAPAN.

Holotype ♂: Obihiro S.Takano [underside of label] 14/V'50/ 111/ *Eurithia emdeni* Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Eurythia fucosa Mesnil, 1975c, Fliegen palaearkt. Reg. 64g: 1400. ITALY.

Holotype ♂: Bormio 15 - 19 VII/ *Eurithia fucosa* Mesn. L.P. Mesnil det., 1975/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Paratype: 1♀, France.

Eurythia monticola Mesnil, 1967, Mushi 41: 48. FRANCE.

Holotype ♂: Montlouis Po. VII/ *Eurythia monticola* Mesn. L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: There may be paratypes among the CNC material, but they are not labeled as such.

Eusisyropa blandooides Thompson, 1963b, Studia ent. 6: 297. TRINIDAD.

Holotype ♀: HOLOTYPE *Eusisyropa blandooides* Thomp. CNC No. 9789/ Santa Cruz V. 3. 5.60 Trinidad,BWI./ HOLOTYPE CNCNo./ *Eusisyropa blandooides* n.sp.

Eutachina atricans Villeneuve, 1938b, Bull. Mus. r. Hist. nat. Belg. 14(4): 3. MALAWI.

Syntypes: 1♂, 3♀♀.

♂: Mt. Mlanje, Nyasaland, 16. IX. 1913. S. A. Neave./ *Tachina atricans* n.sp./ TYPE/ Exorista atricans Villen. L.P. Mesnil det., 1970/ Syntype/ CNC Syntype *Eutachina atricans* Villeneuve Label affixed 1994/ EX L.-P. MESNIL

COLLECTION 1970.

♀: Mt. Mlanje, Nyasaland, 26.VIII 1913 S. A. Neave./ Exorista atricans Villen. L.-P. Mesnil det., 1970/ CNC Syntype *Eutachina atricans* Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

♀: Mt. Mlanje, Nyasaland, 27.VIII 1913 S. A. Neave./ Exorista atricans Villen. L.P. Mesnil det., 1970/ CNC Syntype *Eutachina atricans* Label affixed 1994/ EX L.-P. MESNIL

COLLECTION 1970.

Eutelothyria trinitatis Thompson, 1963a, Can. J. Zool. 41: 484.

TRINIDAD.

Syntype: 1♂.

♂: MARACAS V. Trinidad,BWI. MARCH.5.1953./ Collector F.J.Simmonds/ *Eutelothyria trinitatis* ns/ HOLOTYPE *Eutelothyria trinitatis* Thomp. CNC No. 9783/ CNC Syntype *Eutelothyria trinitatis* Thompson Label affixed 1994.

Euthera setifacies Brooks, 1945b, Can. Ent. 77: 79. CANADA.

Holotype ♂: Trinity Valley B.C. July 25 1937 K. Graham/ HoloTYPE ♂ *Euthera setifacies* No. Brks 5514.

Allotype ♀: Same data as holotype except with allotype label and collected "July 26 1937".

Paratype: 1♂, British Columbia.

Eutheresia trivittata Curran, 1929, Can. Ent. 61: 33. CANADA.

Holotype ♀: *Eutheresia* TYPE ♀ trivittata Curran No 2754/Rondeau, Ont. 21-VI 1922 A.W.Baker.

Paratype: 1♀, Ontario.

Eutrapelus atlxicoensis Reinhard, 1974, Can. Ent. 106: 1164.

MEXICO.

Holotype ♂: Atlixco, Pueb. Sept. 8, 1947/ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE *Eutrapelus atlxicoensis* Rnh. CNC No.

Allotype ♀: Atlixco 8-IX-47/ F. A. Cowan M. R. Wheeler collectors/ IDEMA illustration/ ALLOTYPE CNC No.

Paratypes: 2♂♂, Mexico.

Eutricogena polita Brooks, 1945b, Can. Ent. 77: 92. CANADA.

Holotype ♂: Holo TYPE ♂ *Eutricogena polita* Brks No. 5533/ Agassiz,B.C. 8.V 1926 R.Glendenning.

Allotype ♀: Cultus L., B.C. 20.X-1938 J.K.Jacob/ Allo TYPE ♀ *Eutricogena polita* Brks No. 5533.

Paratype: 1♀, British Columbia.

Eutrixa laxifrons Reinhard, 1962b, Pan-Pacif. Ent. 38: 216. USA.

Holotype ♀: nr.Flagstaff,ARIZ.Ft. ValleyExForHq. VI-4- 1959.LACarruth/ HOLOTYPE *Eutrixa laxifrons* Reinhard/ *Eutrixa laxifrons* R Rnh.

Paratype: 1♀, California.

Evanalia medicinensis Strickland, 1941, Can. Ent. 73: 64.

CANADA.

Holotype ♀: HOLOTYPE *Evanalia medicinensis* 5160 No.Strickland/ Medicine Hat,Alta. 9 VIII 1939 E. H. Strickland/ Type *Evanalia medicinen-sis* E.H.S.

Remarks: The CNC type number was published as 5161.

Evidomyia infida Reinhard, 1958b, J. Kans. ent. Soc. 31: 226. USA.

Holotype ♂: 12mi.SE.Ivanpah S.Berna.Co,Calif V-1-1956/ M. Wasbauer Collector/ HOLOTYPE *Evidomyia infida* Reinhard.

Allotype ♀: Same data as holotype except with allotype label.

Paratype: 1♂, California.

Remarks: Date of collection of holotype was published as 1 November 1956. The collectors name was published as Vasbauer.

- Exorista aberrans* var. *syriaca* Mesnil, 1941a, Bull. Soc. ent. Fr. 46: 21. ISRAEL.
 Lectotype ♂: Route de Jerusalem Hebron 5. 3. 1935/ *Exorista syriaca* Mesnil L.P. Mesnil det., 1970/ TYPE/ CNC Syntype *Exorista aberrans* var. *syriaca* Mesnil Label affixed 1994/ Lectotype fixation by Mesnil, 1960: 585. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.
 Paralectotype: 1♂, Israel.
 Remarks: We regard Mesnil's (1960a: 585) reference to a holotype as a lectotype fixation for this species (see O'Hara in press).
- Exorista aplomyoides* Villeneuve, 1936e, Ark. Zool. 27A: 4. CHINA.
 Holotype ♀: Kina N.O. Szechuan/ Sven Hedin's Exp. Ctr. Aslen Dr Hummel/ 25/5/ *Exorista aplomyoides* n.sp./ Thelyconychia *aplomyoides* Vill. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Exorista caesar* Aldrich, 1916, Can. Ent. 48: 20. CANADA.
 Holotype ♂: TYPE No. 488/ *Exorista caesar* Ald. Type./ Simcoe, Ont. L.Caesar. 1914/ Archips cerasivorana Parasite.
 Paratypes: 1♂, 2♀♀, Ontario.
- Exorista loxostegeae* Reinhard, 1922, Ann. ent. Soc. Am. 14 (1921): 331. USA.
 Syntypes: 6♂♂, 2♀♀.
 3♂♂, 1♀: College Station 6-14-20 Texas/ H.J.Reinhard Collector/ Bria/ PARATYPE *Exorista loxostegeae* Reinhard/ CNC Syntype *Exorista loxostegeae* Reinhard Label affixed 1994.
 ♂: College Station 6-15-20 Texas/ H.J.Reinhard Collector/ br/ PARATYPE *Exorista loxostegeae* Reinhard/ Pseudoperichaeta R erecta (Coq.)/ CNC Syntype *Exorista loxostegeae* Reinhard Label affixed 1994.
 2♂♂: College Station 6-15-20 Texas/ H.J.Reinhard Collector/ br/ PARATYPE *Exorista loxostegeae* Reinhard/ CNC Syntype *Exorista loxostegeae* Reinhard Label affixed 1994.
 ♀: College Station 6-16-20 Texas/ H.J.Reinhard Collector/ br/ PARATYPE *Exorista loxostegeae* Reinhard/ CNC Syntype *Exorista loxostegeae* Reinhard Label affixed 1994.
- Exorista perdives* Villeneuve, 1926d, Bull. Annls Soc. ent. Belg. 66: 198. SPAIN.
 Holotype ♂: Andalusia Algeciras 1.-10.V.'25.Zerny/ *Exorista (Zenillia) dives* Villen/ Eumeella perdives Villen. L.P. Mesnil det., 1969/ TYPE.
- Exorista (Alsomyia) triseriella* Villeneuve, 1929a, Bull. Annls Soc. ent. Belg. 69: 185. EGYPT.
 Holotype ♂: WADI-HOFF 21.3.22/ *Alsomyia triseriella* n.sp./ Nealsomyia *triseriella* Vill. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Exorista (Guerinia) lacteipennis* Mesnil, 1970, Mushi 44: 115. TUNISIA.

Holotype ♂: Tassili S. Tunisia 4. 1953/ *Guerinia lacteipennis* Mesn. L.P. Mesnil det., 1970/ TYPE.
 Remarks: Holotype is missing abdomen, and terminalia are glued to a card mount below specimen.

- Exorista (Guerinia) paligera* Mesnil, 1970, Mushi 44: 113. FRANCE.

Holotype ♂: EAUX CH [Chaudes] 23.VIII.18 B. [Basses] P. [Pyrénées]/ *Guerinia paligera* Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Holotype terminalia are glued to a card mount below specimen. There may be a paratype among the CNC material, but it is not labeled as such.

- Exorista (Podotachina) rutilans* Mesnil, 1970, Mushi 44: 113. EGYPT.

Holotype ♂: Sinai IV Thal Israel 27.III.1969 leg. Kugler/ T 260/ *Exorista rutilans* Mesnil L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970/ TYPE.
 Paratype: 1♀, Egypt.

- Exorista (Pokornia) kugleri* Mesnil, 1960a, Fliegen palaearkt. Reg. 64g: 592. ISRAEL.

Holotype ♂: 1-V 1957. Eingedi Israel Coll Kugler I/ *Exorista kugleri* Mesnil L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970/ TYPE.

- Exorista (Pokornia) aberrans* var. *numidica* Mesnil, 1946, Encycl. ent. (B) II Dipt. 10: 54. MOROCCO.

Lectotype ♂: MAROC SIDI TAIBI PRÉS KENITRA R BENOIST 1928/ 13 MAI 28/ *Exorista numidica* Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ CNC Syntype *Exorista (Pokornia) aberrans* var. *numidica* Mesnil Label affixed 1994/ Lectotype fixation by Mesnil, 1960: 585. Label affixed 1995.

Paralectotype: 1♀, Morocco.
 Remarks: We regard Mesnil's (1960a: 585) reference to a holotype as a lectotype fixation for this species (see O'Hara in press).

- Exorista (Pokornia) ghanii* Mesnil, 1971a, Entomophaga 16: 68. PAKISTAN.

Holotype ♂: Karis 27.VII.66/ Pakistan/ Ex.larva feeding on leaves of Halogoton flaconeri/ *Exorista ghanii* Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Remarks: Holotype puparium is pinned below specimen, and terminalia are glued to a card mount. Date of collection of holotype was published as 8 August 1966.

- Exorista (Prosalia) thomasi* Mesnil, 1960a, Fliegen palaearkt. Reg. 64g: 563. INDONESIA.

Holotype ♂: Ned. Niauw Guinea Kota Nica Res. Hollandia 8-VI 1959 leg. R.T. Simon Thomas/ alt.: 75 m. Host. Psychidae loc.: cups./ TYPE/ *Exorista (s.g. Prosalia) thomasi* Mesnil L.P. Mesnil det., 1960/ HOLOTYPE ♂ of *Exorista thomasi* Mesn. Examined 1970 R.W.Crosskey.

Exorista (Scotiella) cantans Mesnil, 1960a, Fliegen palaearkt. Reg. 64g: 574. JAPAN.

Holotype ♂: VIII-14 1949 Hanno N.FUKUHARA/ 175/ Exorista cantans Mesnil L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970/ TYPE.

Exorista (Scotiella) patelliforceps Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 20. JAPAN.

Holotype ♂: Hanase Kyoto, Japan July 14, 1957 Coll. T. Horio/ 348/ Exorista patelliforceps Mesnil L.P. Mesnil det., 1960/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Exoristoides harrisi Reinhard, 1935c, Ann. ent. Soc. Am. 28: 160. USA.

Holotype ♂: Ames, Iowa May 25 1932 H. M. Harris/ Reared from Centipede Geophilus sp.? over [underside of label] Pupated May 12 Emerged May 25 HMH/ HOLOTYPE Exoristoides harrisi Reinhard/ Exoristoides harrisi det.H.J.Reinhard Rein.

Remarks: Holotype puparium is pinned below specimen.

Exoristoides verticalis Reinhard, 1935c, Ann. ent. Soc. Am. 28: 161. MEXICO.

Holotype ♂: Mexico 28.VIII.23/ HOLOTYPE Exoristoides verticalis Reinhard/ Helioplagia verticalis Rnh./ Exoristoides verticalis det.H.J.Reinhard Rein.

Fabriciella actinosa Reinhard, 1938, Can. Ent. 70: 8. USA.

Holotype ♀: Alamosa, 8.2.34 Colo. 5466 Acc./ HOLOTYPE Fabriciella actinosa Reinhard/ Fabriciella actinosa Rnh. Reinhard Coll'n/ Fabriciella actinosa Rein det. H.J.Reinhard.

Paratype: 1♂, Colorado.

Fabriciella ampliforceps Rowe, 1931, Ann. ent. Soc. Am. 24: 673. CANADA.

Holotype ♂: HOLOTYPE Fabriciella ampliforceps Rowe No. 3315/ Mahone Bay 9.VII.87 C.A. Hamilton/ 15/ Holotype.

Fabriciella argentea Rowe, 1931, Ann. ent. Soc. Am. 24: 654. CANADA.

Holotype ♂: HOLOTYPE Fabriciella argentea Rowe No. 3317/ Banff, Alta. 9 VIII 1922 C.B.D.Garrett 7700 ft/ Holotype.

Fabriciella brevirostris Tothill, 1924b, Can. Ent. 56: 264. CANADA.

Holotype ♂: HoloTYPE F. brevirostris Tothill No. 809/ Keremeos, B.C. 16.VII.1923 C.B.Garrett.

Remarks: There is one specimen labeled as a paratype but it is from a locality not mentioned in original description.

Fabriciella canadensis Tothill, 1924b, Can. Ent. 56: 264. CANADA.

Holotype ♂: HoloTYPE F. canadensis Toth No. 803./ Godbout, Que. 20-VII-1918 E.M.Walker.

Paratypes: 2♂♂, Alberta.

Fabriciella cordiforceps Rowe, 1931, Ann. ent. Soc. Am. 24: 649. USA.

Holotype ♂: HOLOTYPE Fabriciella cordiforceps Rowe No. 3316/ Pine Val. Mt. in. Utah 7000 ft./ BROOKLYN MUSEUM COLL. 1929/ ♂/ Holotype.

Allotype ♀: ALLOTYPE Fabriciella cordiforceps Rowe No. 3316/ Oak Creek Canon Ariz. 6000 ft. Aug F H.Snow/ Allotype/ SLIDE No. 0263.

Paratypes: 2♂♂, Arizona.

Fabriciella egula Reinhard, 1938, Can. Ent. 70: 9. USA.

Holotype ♂: San Luis Val. VIII-10-'30/ HOLOTYPE Fabriciella egula Reinhard/ Fabriciella egula Rnh. Reinhard Coll'n.

Paratype: 1♂, Oregon.

Fabriciella eureka Reinhard, 1942c, Bull. Brooklyn ent. Soc. 37: 28. USA.

Holotype ♂: Eureka, Cal 6-V 1935/ E.W.Baker Collector/ HOLOTYPE Fabriciella eureka Reinhard/ Fabriciella eureka R. Rnh.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 4♂♂, 1♀, California; 1♀, British Columbia.

Fabriciella evanida Reinhard, 1953c, Pan-Pacif. Ent. 29: 53. USA.

Holotype ♂: Lake City Colo. 8700 Aug.8-15'36/ Chas.L.Fluke Collector/ HOLOTYPE Fabriciella evanida Reinhard/ Fabriciella evanida R. Rnh.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 3♂♂, 3♀♀, Colorado; 1♂, Utah.

Fabriciella hispida Tothill, 1924b, Can. Ent. 56: 265. CANADA.

Holotype ♂: TYPE F. hispida Toth. No. 805./ SUDBURY, ONT. CAN. 1890.

Allotype ♀: ALLOTYPE F. hispida Toth. No. 805/ Halifax,N.S. 21 VIII 1915 J.Perrin.

Paratypes: 3♂♂, Ontario.

Remarks: The status of the specimen labeled as allotype is uncertain because, its locality is not the same as that published for the allotype. There is one additional specimen labeled as a paratype but it is from a locality not mentioned in original description.

Fabriciella latianulum Tothill, 1924b, Can. Ent. 56: 266. CANADA.

Holotype ♂: Holo TYPE F. latianulum Toth. No. 813./ Royal Oak, B.C. 30-VI-17. B.H.T.

Allotype ♀: Allo TYPE F. latianulum Toth. No. 813./ Agassiz,B.C. 22.VIII.1921 R.Glendenning.

Paratypes: 16♂♂, British Columbia; 1♂, Ontario.

Fabriciella latiforceps Tothill, 1924b, Can. Ent. 56: 266.

CANADA.

Holotype ♂: Godbout, Que. 25-VII-1918 E.M.Walker/ TYPE F. latiforceps [sic] Toth. No. 804.

Paratype: 1♂, British Columbia.

Fabriciella latifrons Tothill, 1924b, Can. Ent. 56: 260, 269.

CANADA.

Holotype ♂: N. W. T. CAN./ HoloTYPE F. latifrons Tothill No. 811.

Paratypes: 2♂♂, 2♀♀, Manitoba; 1♀, Northwest Territories; 1♀, Colorado.

Fabriciella latigena Tothill, 1924b, Can. Ent. 56: 262. CANADA.
Holotype ♂: Lillooet, B.C. 3000 ft 26-VII-17. J.D.T./
Holo. TYPE F. latigena Toth No. 806.
Allotype ♀: TYPE F. latigena Tothill No. 806.
Paratypes: 3♂♂, British Columbia.

Fabriciella longiunguis Tothill, 1924b, Can. Ent. 56: 265.
CANADA.
Holotype ♂: Holo. TYPE F. longiunguis Toth. No. 807/
Vernon, B.C. 19 M.H. Ruhmann.
Remarks: Holotype abdomen is stored in gelatin
capsule below specimen.

Fabriciella nigella Reinhard, 1938, Can. Ent. 70: 10. USA.
Holotype ♂: Dillon Mont. July 16, 1934/ Dorothy
Martin Coll./ HOLOTYPE *Fabriciella nigella*
Reinhard/ *Fabriciella nigella* Rnh. Reinhard
Coll'n/ *Fabriciella nigella* Rein. det. H.J. Reinhard.

Fabriciella orbitalis Reinhard, 1942c, Bull. Brooklyn ent. Soc. 37:
26. USA.
Holotype ♂: ASA Cr Kane Co. Utah/ C.Lynn Hayward
collector/ July/ HOLOTYPE *Fabriciella orbitalis*
Reinhard/ *Fabriciella orbitalis* Rein
det. H.J. Reinhard/ *Fabriciella orbitalis* Rnh.
Reinhard Coll'n.
Paratype: 1♀, Colorado.

Fabriciella pilosa Tothill, 1924b, Can. Ent. 56: 263. CANADA.
Holotype ♂: Holo. TYPE F. pilosa Toth. No. 801/
Hedley, B.C. 29.VII.1923 C.B. Garrett.
Paratypes: 3♂♂, Alberta; 1♂, British Columbia.

Fabriciella rostrata Tothill, 1924b, Can. Ent. 56: 267. CANADA.
Holotype ♂: Holo. TYPE F. rostratum Toth No. 814/
Penticton B.C. 8.VII.1913 E.M. Anderson.
Allotype ♀: Allo. TYPE F. rostratum Toth No. 814/
Hedley, B.C. 29.VIII.1923 C.B. Garrett.
Paratypes: 5♂♂, Alberta; 3♂♂, British Columbia.
Remarks: Date of collection of holotype was published
as 8 August 1913.

Fabriciella spineiventer Tothill, 1924b, Can. Ent. 56: 266.
CANADA.
Holotype ♂: Holo. TYPE F. spineiventer Toth. No. 812./
Mt. McLean, B.C. 8000-8500 25-VII-17 J.D.T.
Paratype: 1♂, British Columbia.

Fabriciella spinosa Tothill, 1924b, Can. Ent. 56: 263. CANADA.
Holotype ♂: Holo. TYPE F. spinosa Tothill No. 800./
Peachland, B.C. 22.VII. 12 J. B. Wallis.
Paratype: 1♂, California.

Fausta inusta Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 57.
JAPAN.

Holotype ♂: Obihiro S.Takano [underside of label]
30/VI'51/ 110/ *Fausta inusta* Mesnil L.P. Mesnil
det., 1967/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Floradalia major Thompson, 1963a, Can. J. Zool. 41: 486.
TRINIDAD.

Syntype: 1♀.
♀: HOLOTYPE *Floradalia major* Thomp. CNC
No. 9784/ MARACAS V. TRINIDAD, BWI NOV. 21.
1952./ Collector F.J.Simmonds/ 19/ *Floridalia* [sic]
major ng. n.sp./ CNC Syntype *Floradalia major*
Thompson Label affixed 1994.

Floradalia minor Thompson, 1963a, Can. J. Zool. 41: 488.
TRINIDAD.

Syntype: 1♀.
♀: HOLOTYPE *Floradalia minor* Thomp. CNC
No. 9785/ St. Augustine, TRINIDAD, BWI OCT. 30.
1953./ Collector F.J.Simmonds/ V V-2/ *Floradalia*
minor TT./ CNC Syntype *Floradalia minor*
Thompson Label affixed 1994.

Frontina malacosomae Curran, 1925b, Can. Ent. 57: 155.
CANADA.

Holotype ♂: Holo. TYPE ♂ *Frontina malacosomae* Curr.
No. 1306/ Truro, N.S. 18 July 14/ No 246/ Parasite
of American Tent caterpillar/ *Lespesia frenchii*
(Williston) Det. D.F. Beneway 1960.
Allotype ♀: Same data as holotype except with allotype
label and without tent caterpillar label.

Frontina spectabilis Aldrich, 1916, Can. Ent. 48: 21. CANADA.
Holotype ♀: TYPE *Frontina spectabilis* Ald. No. 490/
Waubamic, Parry Sd., Ont./ VIII-5-15.

Frontiniella incarcerata O'Hara, 1993, Can. Ent. 125: 19. USA.

Holotype ♂: USA AZ Graham Co. 2.4km w. on
Hwy.366 from Hwy.666 1160m 27-28.v.1991
desert malaise J.E. O'Hara/ HOLOTYPE
Frontiniella incarcerata O'Hara CNC No. 21508.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 68♂♂, 4♀♀, Arizona.

Frontiniella jorgensenii O'Hara, 1993, Can. Ent. 125: 24. USA.

Holotype ♂: NEW MEXICO, Chaves Co. 20 mi NW
Kenna 13 May 1988 Malaise T. N. Jorgensen,
Coll./ Retained from Det Lot 91-195/ HOLOTYPE
Frontiniella jorgensenii O'Hara CNC No. 21510.
Allotype ♀: Same data as holotype except with allotype
label and collected 15 miles northwest of Kenna,
New Mexico.
Paratypes: 1♂, 1♀, Arizona.
Remarks: Holotype terminalia are stored in glycerine in
a plastic microvial pinned below specimen.

Gaedea hispanica Mesnil, 1953a, Fliegen palaearkt. Reg. 64g: 273.
SPAIN.

Holotype ♂: El Ventorrillo Espanye 14 Sept. 1935/
Gaedea hispanica Mesn. L.P. Mesnil det., 1969/
TYPE/ EX L.-P. MESNIL COLLECTION 1970.
Paratype: 1♂, Spain.

Gaediophana monnula Reinhard, 1951, Bull. Brooklyn ent. Soc.
46: 5. MEXICO.

Holotype ♂: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan
M. R. Wheeler Collectors/ HOLOTYPE

- Gaediophana monnula* Reinhard/ *Gaediophana monnula* R. Rnh.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 14♂♂, 10♀♀, Mexico.
- Gaedioxenix haematoxides* Villeneuve, 1937a, Revue Zool. Bot. afr. 29: 207. SOUTH AFRICA.
Holotype ♂: ♂/ Somerset W. STand Dr. Brauns.
Kapland/ *Gaedioxenix haematoxides* Typ. Villen./
Gaedioxenix haematoxides Vill. L.P. Mesnil det.,
1969/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.
- Gaedioxenix setifrons* Villeneuve, 1937a, Revue Zool. Bot. afr. 29:
207. SOUTH AFRICA.
Holotype ♀: [pink blank label]/ Capland Stellenbosch
Nov.10 1925 Dr. H. Brauns./ *Gaedioxenix*
setifrons Typ. Villen./ *Gaedioxenix setifrons* Vill.
L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
- Gastropitlops ater* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 78.
JAPAN.
Holotype ♀: 5-V-1951 Nokogiriyama (Okutama)
S.KATO/ *Gastropitlops ater* Mesnil L.P. Mesnil
det., 1976/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
- Germaria (Atractochaeta) barbara* Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 37. ALGERIA.
Holotype ♂: La Calle (algérie)/ *Germaria barbara* Mesn.
L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
Remarks: Holotype with only head and portion of thorax remaining.
- Germaria (Germaria) hispanica* Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 38. SPAIN.
Holotype ♂: Villaverde Arias Encobet [underside of label] 14.vi.1906/ *Germaria hispanica* Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
Remarks: There may be two paratypes among the CNC material but they are not labeled as such.
- Gibsonomyia nigricosta* Curran, 1925a, Can. Ent. 57: 282.
CANADA.
Holotype ♂: *Gibsonomyia* TYPE *nigricosta* Curran
No.1427/ Banff, Alta. 27 VII.1922 C.B.D.Garrett.
Paratype: 1♂, British Columbia.
- Gigamyiopsis funebris* Reinhard, 1964b, Acta zool. mex. 7: 6.
MEXICO.
Holotype ♂: Acatlipa,Morelos W.G.Downs,Coll./
HOLOTYPE *Gigamyiopsis funebris* Reinhard/
Gigamyiopsis funebris R Rnh.
Allotype ♀: Acatlipa Morelos July 29, 1950 Coll.
WGDowns/ ALLOTYPE *Gigamyiopsis funebris*
Reinhard.
- Ginglymia bicolor* Curran, 1923, Can. Ent. 55: 246. CANADA.
Holotype ♀: TYPE *Ginglymia* [sic] *bicolor* No. Curr
553/ Saanich B.C. 13. VIII. 1918 W. Downes.
Paratypes: 2♀♀, British Columbia.
- Ginglymia fracida* Reinhard, 1962b, Pan-Pacif. Ent. 38: 220.
MEXICO.
Holotype ♂: Colima,Mex. Mar. 27,1934/ HOLOTYPE
Ginglymia fracida Reinhard/ *Ginglymia fracida* R
Rnh.
Allotype ♀: Colima,Mex. Mar. 27,1934/ ALLOTYPE
Ginglymia fracida Reinhard.
- Gonia albagenae* Morrison, 1940, Can. J. Res. 18: 349. CANADA.
Holotype ♂: Penticton, B.C. 12.IV.1927 E.R.Buckell/
Gonia TYPE *albagena* Morr No. 5249.
Paratypes: 4♂♂, British Columbia.
Remarks: The CNC type number was published as
5034.
- Gonia aldrichi* Tothill, 1924a, Can. Ent. 56: 209. CANADA.
Holotype ♂: Holotype G. *aldrichi* Tothill No. 786./
Coaldale, Alta. April 26 1915 E. H. Strickland/
/12/X/ Host. *Euxoa ochrogaster*/ SLIDE F.O.M.
No. 38 (S).
Paratypes: 5♂♂, Alberta.
Remarks: Holotype terminalia are mounted on a
microscope slide.
- Gonia aturgida* Brooks, 1944a, Can. Ent. 75 (1943): 233. USA.
Holotype ♂: Idaho April 28, 1919/ E.H.Quayle
Collector/ HoloTYPE ♂ *Gonia aturgida* No. Brks
5456.
Allotype ♀: Idaho April 28, 1919/ E H. Quayle
Collector/ 118/ *Gonia turgida* [sic] Det.
C.H.Curran Coq./ AlloTYPE ♀ *Gonia aturgida* No.
Brks 5456.
Paratypes: 2♂♂, Saskatchewan; 1♂, Washington.
- Gonia breviforceps* Tothill, 1924a, Can. Ent. 56: 210. CANADA.
Holotype ♂: Holotype G. *breviforceps* [sic] Tothill
No. 788/ Lethbridge Alta. B April 15 1915 E. H.
Strickland //12/Y/ SLIDE F.O.M. No. 35 (S).
Paratypes: 3♂♂, Alberta; 1♂, British Columbia; 1♂,
Montana.
Remarks: Holotype terminalia are mounted on a
microscope slide.
- Gonia brevipulvilli* Tothill, 1924a, Can. Ent. 56: 211. CANADA.
Holotype ♂: Holotype G. *brevipulvilli* Tothill No.
790./ F'ton [Fredericton] 3.V.12/ JDTothill
Collector/ SLIDE F.O.M. No. 31. (S).
Paratypes: 1♂, New Brunswick; 1♂, Massachusetts.
Remarks: Holotype terminalia are mounted on a
microscope slide.
- Gonia contumax* Brooks, 1944a, Can. Ent. 75 (1943): 233. USA.
Holotype ♂: Westport, Wn. [Washington] IV-8-1933/
S.E.Crumb, Coll./ HoloTYPE ♂ *Gonia contumax*
No. Brks/ *Gonia contumax* Det. Brks A.R.Brooks/
Reinhard Collection.
Allotype ♀: Westport, Wash. IV-9 33 J. Wilcox, Col./
AlloTYPE ♀ *Gonia contumax* No. Brks/ Reinhard
Collection.
Paratype: 1♂, Washington.
- Gonia discalis* Morrison, 1940, Can. J. Res. 18: 347. CANADA.
Holotype ♂: Nicola, B.C. 28 IV 1922 P.N.Vroom/
Gonia TYPE *discalis* Morr No. 5248.

Paratypes: 12♂♂, British Columbia.

Remarks: Holotype is missing head. The CNC type number was published as 5003.

Gonia fissiforceps Tothill, 1924a, Can. Ent. 56: 207. CANADA.

Holotype ♂: Holotype G. fissiforceps [sic] Tothill No. 782/ Lillooet B.C 24.VI 1917 det. Tothill.

Paratype: 1♂, Alberta.

Remarks: Holotype terminalia are stored in glycerine in a glass microvial pinned below specimen.

Gonia frontosa Say, 1829, J. Acad. nat. Sci. Philad. 6: 175. CANADA.

Neotype ♂: NeoTYPE G. frontosa Tothill Say. No. 785/ Aweme, Man. 21.V. 1921 P. Vroom/ SLIDE broken 34(S) No. F.O.M.

Remarks: For neotype designation see Tothill (1924a: 211). Neotype terminalia are mounted on a microscope slide.

Gonia grandipulvilli Morrison, 1940, Can. J. Res. 18: 345. CANADA.

Holotype ♂: Edmonton 25/4/23 at Salix/ G. grandipulvilli Teste Morrison/ Coll. F.S. Carr. (Authority of E.H. Strickland [Strickland] 1940.)/ XXVII/ Gonia TYPE grandipulvilli Morr No. 5251.

Paratypes: 3♂♂, Quebec; 13♂♂, Ontario; 14♂♂, Saskatchewan; 5♂♂, Alberta; 1♂, British Columbia; 1♂, Montana.

Remarks: Holotype terminalia are stored in glycerine in a glass microvial pinned below specimen. The CNC type number was published as 5032.

Gonia longiforceps Tothill, 1924a, Can. Ent. 56: 208. CANADA.

Holotype ♂: TYPE G. longiforceps [sic] Tothill No. 784/ L'brid, [Lethbridge] Alta. B April 21 1915 E.H. Strickland/ Host Porosagrotis orthogonia/ /12/Y./ SLIDE F.O.M. No 32 (S).

Paratypes: 6♂♂, British Columbia; 3♂♂, Alberta.

Remarks: Holotype terminalia are mounted on a microscope slide.

Gonia longipulvilli Tothill, 1924a, Can. Ent. 56: 211. CANADA.

Holotype ♂: Holotype G. longipulvilli Tothill No. 789/ Royal Oak B.C 30-VI-17 B.H.T.

Paratypes: 1♂, Alberta; 1♂, British Columbia.

Remarks: Holotype terminalia are stored in glycerine in a glass microvial pinned below specimen.

Gonia nana Becker, 1908, Mitt. zool. Mus. Berl. 4: 197.

MADEIRA.

Syntype: 1♂.

♂: Madeira 51039. IV/ Typus/ Gonia nana Beck./ Gonia nana Becker L.P. Mesnil det, 1969/ TYPE/ CNC Syntype Gonia nana Becker Label affixed 1994.

Gonia occidentalis Brooks, 1944a, Can. Ent. 75 (1943): 234. CANADA.

Holotype ♂: Hedley BC 15/V 1920 W.B. Anderson/ Holotype ♂ Gonia occidentalis No. Brks 5454.

Allotype ♀: Penticton, B.C. 4.V 1927 E.R. Buckell/ AlloTYPE ♀ Gonia occidentalis No. Brks 5454.

Paratypes: 1♂, 2♀♀, British Columbia; 4♀♀, Colorado; 1♂, Montana; 1♂, Washington.

Gonia pilosa Brooks, 1944a, Can. Ent. 75 (1943): 231. CANADA.

Holotype ♀: Treesbank, Man. 20.VII 10 J. B. Wallis/ b13y/ Holotype ♀ Gonia pilosa No. Brks 5453.

Gonia reinhardi Brooks, 1944a, Can. Ent. 75 (1943): 232. USA.

Holotype ♂: Pingree Park Colo. VIII-17-32/ Holotype ♂ Gonia reinhardi No. Brks/ Gonia reinhardi Det. Brks A.R.Brooks/ Reinhard Collection.

Allotype ♀: Yakima, Wn. [Washington]/ A. L. Lovett Coll. 7-26-22/ Gonia longipulvilli det. Aldrich Toth/ AlloTYPE ♀ Gonia reinhardi No. Brks/ Reinhard Collection.

Paratype: 1♂, Wyoming.

Gonia robusta Brooks, 1944a, Can. Ent. 75 (1943): 234. CANADA.

Holotype ♂: Wabamum Alta. May 11 1924 E. H. Strickland/ Holotype ♂ Gonia robusta No. Brks 5457.

Paratypes: 5♂♂, Alberta.

Gonia sagax Townsend, 1892b, Can. Ent. 24: 65. USA.

Neotype ♂: NeoTYPE G. sagax Tns. No. 406./ MiddlxCo IV-6 NJ/ SLIDE F.O.M No. 30. (S.).

Remarks: Neotype terminalia are mounted on a microscope slide. For neotype designation see Tothill (1924a: 199).

Gonia smithi Brooks, 1944a, Can. Ent. 75 (1943): 235. CANADA.

Holotype ♂: Holotype ♂ Gonia smithi Brks No. 5455/ onto, [Toronto] Ont. 13-VI-05/ Paratype G. discalis Morr No. 5248.

Allotype ♀: 14-4-98/ AlloTYPE ♀ Gonia smithi No. Brks 5455.

Paratypes: 1♀, Prince Edward Island; 3♂♂, Ontario; 1♂, Manitoba.

Gonia tenuiforceps Morrison, 1940, Can. J. Res. 18: 356. CANADA.

Holotype ♂: Hopedale, Labr. 21-VII 1926 W.W. Perrett/ SLIDE Vial No. 110/ Gonia TYPE tenuiforceps Morr No. 5250.

Paratypes: 4♂♂, 3♀♀, Newfoundland.

Remarks: The CNC type number was published as 5035. Holotype abdomen and terminalia are stored in glycerine in a plastic microvial pinned below specimen.

Gonia yukonensis Tothill, 1924a, Can. Ent. 56: 210. CANADA.

Holotype ♂: TYPE G. yukonensis Tothill. No. 787/ Yukon Terr D.D. CAIRNER 1916/ LAT 62°30'-63°06' Long 137°30'-139°30'/ Lab. No. Yukon. Terr. 1916 D.D. Cairnes (Not to me. G.E.S.) Appears the same as G. fissiforceps Toth. Loc. F.O. Morrison Dom. Ent. Br. Ottawa F.O.M.

Remarks: Holotype terminalia are stored in glycerine in a glass microvial pinned below specimen.

Gonioclea apicalis Villeneuve, 1929b, Bull. Annls Soc. ent. Belg.

69: 100. TURKMENIA.

Holotype ♀: Tr. Casp. Gr. Balchan/ Gonioclea apicalis
Typ. Villen./ TYPE/ Eremogonia apicalis Rohd.
L.P. Mesnil det., 1969.

Gonioclea atricans Villeneuve, 1929b, Bull. Annls Soc. ent. Belg. 69: 100. TURKMENIA.
Holotype ♀: Askhabad/ Gonioclea atricans Typ.
Villen./ TYPE/ Eremogonia desertorum Rohd.
L.P. Mesnil det, 1969.

Goniophthalmus simonyi Villeneuve in Becker, 1910, Denkschr. Akad. Wiss., Wien 71: 145. SOCOTRA IS.
Syntype: 1♀.
♀: Ras Shoab, Sokotra; I. '99. O.Simony/
Goniophthalmus Simonyi ♀ Type Villen./ TYPE/
Goniophthalmus simonyi Villen. L.P. Mesnil det.,
1969/ CNC Syntype Goniophthalmus simonyi
Villeneuve Label affixed 1994.

Gopaulia nigrifrons Thompson, 1968, Mem. ent. Soc. Can. 56: 71. TRINIDAD.
Holotype ♂: HOLOTYPE Gopaulia nigrifrons Thomp.
CNC No. 9855/ NURSERY TRINIDAD
30-9-1964/ 1452a ♂ IV.20.65/ Gopaulia nigrifrons
n.sp.
Allotype ♀: ALLOTYPE Gopaulia nigrifrons Thomp.
CNC No. 9855/ NURSERY TRINIDAD
30-9-1964/ 1452B IV-20 65.
Remarks: Holotype terminalia are missing.

Graphogaster (Parahyria) buccata hirtipleura Mesnil, 1973b,
Fliegen palaearkt. Reg. 64g: 1202. FRANCE.
Holotype ♂: Col du Lautaret 2.7.03 (Htes Alpes)/
Graphogaster hirtipleura Mesn. L.P. Mesnil det.,
1975/ TYPE/ EX L.-P. MESNIL COLLECTION
1985.

Gremlinotrophus derisus Reinhard, 1943a, Can. Ent. 75: 163. USA.
Holotype ♀: Amherst, Ohio 7-9 1934/ H J Reinhard
Collector/ HOLOTYPE Gremlinotrophus derisus
Reinhard/ Gremlinotrophus derisus Rnh. det.
H.J.Reinhard.
Paratypes: 1♀, Montana; 2♀♀, Alberta.

Grisdalemyia bigelowi Curran, 1926b, Can. Ent. 58: 134.
CANADA.
Holotype ♀: Grisdalemyia TYPE bigelowi Curran. No.
1953/ LowBush Ont. Lake Abitibi VI-30-1925
N.K.Bigelow/ Grisdalemyia bigelowi Det. Curran
C.H.Curran.

Guerinia trudis Reinhard, 1951, Bull. Brooklyn ent. Soc. 46: 8.
USA.
Holotype ♂: Panguitch, Ut. June, 15 1948 G. F.
Knowlton S. L. Wood/ HOLOTYPE Guerinia
trudis Reinhard/ Guerinia trudis Rnh.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 6♂♂, 7♀♀, Utah; 1♂, Idaho.

Gueriniopsis plausilis Reinhard, 1943a, Can. Ent. 75: 167.
CANADA.

Holotype ♂: Aweme, Man. N. Criddle 29 V 1925/
Holotype G. plausilis Reinhard/ HOLOTYPE
Gueriniopsis plausilis Rnh No. 5402.

Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 2♂♂, 1♀, Manitoba; 1♀, Ohio; 1♂, Iowa;
5♂♂, Wisconsin.

Gymnocarcelia tenuiforceps Reinhard, 1964a, J. Kans. ent. Soc. 37:
49. USA.

Holotype ♂: Amherst, Ohio VII-5 1931/ H J Reinhard
Collector/ HOLOTYPE Gymnocarcelia
tenuiforceps Reinhard/ Gymnocarcelia
tenuiforceps R Rnh.

Gymnocheta frontalis Brooks, 1945b, Can. Ent. 77: 88. CANADA.
Holotype ♂: Mt. Lolo Kamloops, B.C. 2.VI-1938
J.K.Jacob/ HOLOTYPE ♂ Gymnocheta frontalis No.
Brks 5529.

Gymnocheta rufipalpis Brooks, 1945b, Can. Ent. 77: 87.
CANADA.

Holotype ♂: Vancouver 19.5.16 B.C./ R. S. Sherman/
60/ Gymnosoma vivida Will det.Tothill/
HOLOTYPE ♂ Gymnocheta rufipalpis No. Brks
5528.

Gymnodexia bigelowi Curran, 1926d, Can. Ent. 58: 88. USA.

Holotype ♂: Gymnodexia HOLOTYPE ♂ bigelowi
Curran No. 1793./ Robinson Delaware Co Iowa,
VII-24-24 NKBIGelow.

Allotype ♀: Same data as holotype except with allotype
label.

Gymnosoma occidentale Curran, 1927a, Bull. Brooklyn ent. Soc.
22: 144. CANADA.

Holotype ♂: VII 28 ,1920 Vernon ,B.C. M. H.
Ruhmann/ TYPE Gymnosoma occidentale Curr
No. 2636.
Paratypes: 2♂♂, British Columbia.

Hegesinus griphus Reinhard, 1964b, Acta zool. mex. 7: 11.
MEXICO.

Holotype ♂: Teotzlan, Morelos,Mex. 9-26-57
R.&K.Dreisbach/ HOLOTYPE Hegesinus griphus
Reinhard/ Hegesinus griphus R Rnh.

Heliodexodes argenteus Thompson, 1968, Mem. ent. Soc. Can. 56:
197. TRINIDAD.

Holotype ♂: HOLOTYPE Heliodexodes argenteus
Thomp. CNC No. 9885/ Am. Rd. Morvan Hill
XII.30.60 Trinidad,BWI. / [blank red label].

Allotype ♀: ALLOTYPE Heliodexodes argenteus
Thomp. CNC No. 9885/ Champs Fleurs
TRINIDAD, W.I. IX.28.60 W.R.Thompson/
[Blank red label].

Heliosprosopa electilis Reinhard, 1964c, Pan-Pacif. Ent. 40: 120.
MEXICO.

Holotype ♂: Colima,Mex. Mar.27.1934/ HOLOTYPE
Heliosprosopa electilis Reinhard/ Heliosprosopa
electilis R Rnh.

Allotype ♀: Same data as holotype except with allotype
label.

Heliosoprosopa liciata Reinhard, 1964c, Pan-Pacif. Ent. 40: 123.
MEXICO.

Holotype ♂: 7mi NE Jacala Hdgo. Mex. VII.27.54
5000ft/ HOLOTYPE *Heliosoprosopa liciata*
Reinhard/ HOMOTYPE Trichophora
convexincruis Wlp Compared by Wood 89/
Heliosoprosopa liciata R Rnh.
Allotype ♀: Fusa(Cund.) 10-IX 43 H. Osorno 1900/
ALLOTYPE *Heliosoprosopa liciata* Reinhard.

Heliosoprosopa macrocera Reinhard, 1964c, Pan-Pacif. Ent. 40: 120.
COLOMBIA.

Holotype ♀: 11302/ HOLOTYPE *Heliosoprosopa*
macrocera Reinhard/ *Heliosoprosopa macrocera* R
Rnh.

Heliosoprosopa veleda Reinhard, 1964c, Pan-Pacif. Ent. 40: 121.
COLOMBIA.

Holotype ♀: Fusa (Cund.) 10. IX. 43 H. Osorno 1900/
HOLOTYPE *Heliosoprosopa veleda* Reinhard/
Heliosoprosopa veleda R Rnh.

Helocera discolor Villeneuve, 1942a, Bull. Soc. ent. Fr. 47: 50.
JAPAN.

Holotype ♀: JAPON Chu-zenjii 16.8.1909 Edme
Gallois/ *Helocera discolor* Villen./ Eloceria
discolor Villen. L.P. Mesnil det., 1975/ TYPE/ EX
L.-P. MESNIL COLLECTION 1970.

Hemilinnaemyia decorata Villeneuve, 1932b, Bull. Soc. ent. Fr. 37:
269. TAIWAN.

Syntype: 1♀.
♀: Formosa Sauter/ Koshun 908. III./
Hemilinnaemyia decorata Typ. Villen./ TYPE/
Linnaemyia lateralis Towns. L.P. Mesnil det.,
1970/ CNC Syntype *Hemilinnaemyia decorata*
Villeneuve Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

Hemimasicera schnabli Villeneuve, 1908a, Wien. ent. Ztg 27: 286.
UKRAINE.

Lectotype ♀: [locality in Russian] 1907-VI-20/ det.
Dr Villeneuve/ TYPE/ *Hemimasicera Schnabli* ♀
Mohrzecki Type/ T. nidicola Townsend L.P.
Mesnil det., 1969/ CNC Syntype *Hemimasicera*
schnabli Villeneuve Label affixed 1994/ Lectotype
fixation by Mesnil, 1950: 116. Label affixed 1995/
EX L.-P. MESNIL COLLECTION 1970.

Remarks: We regard Mesnil's (1950b: 116) reference to
a holotype as a lectotype fixation for this species.

Hemiphania trispina Villeneuve, 1937a, Revue Zool. Bot. afr. 29:
205. ZAIRE.

Holotype ♂: TYPE/ R [Ruwenzori] 1800, 26.IV.14./
Hemiphania trispina, Type Villen.

Hemyda decumata Reinhard, 1958e, Ent. News 69: 235. MEXICO.

Holotype ♂: Rio Blanco, V.C., Mex., 11-13-57 R.&K.
Dreisbach/ HOLOTYPE *Hemyda decumata*
Reinhard/ *Hemyda decumata* det.
H.J.REINHARD.

Allotype ♀: Same data as holotype except with allotype
label.

Paratypes: 2♂♂, Mexico.

Hemyda latipennis Curran, 1924b, Can. Ent. 56: 252. CANADA.

Holotype ♂: TYPE *Hemyda latipennis* Curr No.745./
Aweme, Man. R.M.White 25.VIII.1923.

Heraultia albipennis Villeneuve, 1920b, Annls Soc. ent. Belg. 60:
119. FRANCE.

Lectotype ♂: Villeneuve leg May. 13.V./ Heraultia
albipennis Villen. L.P. Mesnil det., 1970/
Heraultia albipennis Villen. Typ./ TYPE/ CNC
Syntype Heraultia albipennis Villeneuve Label
affixed 1994/ Lectotype fixation by Mesnil, 1973:
1187. Label affixed 1995/ EX L.-P. MESNIL
COLLECTION 1970.

Remarks: The original description cites M.H. Sicard,
not Villeneuve, as collector. We regard Mesnil's
(1973b: 1187) reference to a holotype as a
lectotype fixation for this species.

Hermyia formosana Villeneuve, 1939a, Bull. Annls Soc. ent. Belg.
79: 353. TAIWAN.

Holotype ♂: Formosa, Sauter/ Kosempo, 1909 VII./
Hermyia formosana, Type Villen.

Hesperophasia aldrichi Curran, 1927b, Can. Ent. 59: 301.
CANADA.

Holotype ♂: *Hesperophasia* TYPE *aldrichi* ♂ Curran
No.2460/ Biggar, Sask. 12. VI. 1925 N.J. Atkinson.
Allotype ♀: *Hesperophasia* AlloTYPE *aldrichi* ♀
Curran No.2460/ Saskatoon, Sask. June 10 1925
Kenneth M. King.
Paratype: 1♀, Saskatchewan.

Histochaeta claripennis Reinhard, 1943b, J. Kans. ent. Soc. 16: 15.
USA.

Holotype ♂: Park City to Silver Lake 11Sep'17 Utah
R.C.Shannon/ HOLOTYPE *Histochaeta*
claripennis Reinhard/ *Histochaeta claripennis* det.
H.J.Reinhard Rein.

Remarks: *Histochaeta* was an unjustified emendation of
Istocheta.

Histochaeta pulchricornis Villeneuve, 1938b, Bull. Mus. r. Hist.
nat. Belg. 14(4): 3. UNKNOWN.

Holotype ♂: Erosba Pan Mar. 1923/ *Histochaeta*
pulchricornis Typ. Villen./ *Brachychaetoides*
pulchricornis Vill. L.P. Mesnil det., 1969/ TYPE/
EX L.-P. MESNIL COLLECTION 1970.

Remarks: Holotype thorax is badly damaged
surrounding the area where pin is inserted.
Histochaeta was an unjustified emendation of
Istocheta.

Homalactia brimleyi Curran, 1926d, Can. Ent. 58: 86. CANADA.

Holotype ♂: *Homalactia* TYPE *brimleyi* Curran No.
1868/ Emo, Ont VI-7-24 J. Brimley.

Houghia nigripalpis Reinhard, 1967, J. Kans. ent. Soc. 40: 108.
USA.

Holotype ♂: 6mi.S.E.Patagonia, ARIZ. Jul.26.'62
Werner, Bequaret, Nollar/ from Condalia

- lycioides and Baccharis glutinosa/ HOLOTYPE
Houghia nigripalpis Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 3♂♂, 2♀♀, Arizona.
- Hyalomyiopsis robusta* Brooks, 1945a, Scient. Agric. 25: 678.
USA.
Holotype ♂: Summit Prairie Grant Co.Ore. El.5500 ft.
VIII-7-41 M. & R.E.Rieder/ HoloTYPE ♂
Hyalomyiopsis robusta No. Brks/ Reinhard
Collection.
Paratypes: 1♂, 1♀, Oregon; 4♂♂, Colorado.
- Hyalurgus itoi* Mesnil, 1967, Mushi 41: 47. JAPAN.
Holotype ♂: R.63-79'B" Japan 1. IV. 64/
Pachynematus itoi/ 69.-5727/ 65-689/ *Hyalurgus*
itoi n.sp. L.P. Mesnil det., 1966/ TYPE.
Remarks: Holotype terminalia are stored in glycerine in
a plastic microvial pinned below specimen.
- Hyalurgus spathulatus* Mesnil, 1967, Mushi 41: 48. JAPAN.
Holotype ♂: Teine nr. Sapporo 10. VII. 1961/
Hyalurgus spathulatus n.sp. L.P. Mesnil det.,
1966/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.
- Hygiella nigripes* Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104:
182. INDIA.
Holotype ♂: PASHOK W. BENGAL 10.VIII.1965/
Resting on Toon/ CIBC-1S KPG.SS/ TYPE/
Hygiella nigripes n. sp. L.P. Mesnil det., 1966/
HOLOTYPE ♂ of *Hygiella nigripes* Mesnil
Examined 1970 R.W.Crosskey.
- Hyleorus hoyti* Mesnil, 1974, Fliegen palaearkt. Reg. 64g: 1259.
NEW CALEDONIA.
Holotype ♂: Noumea NEW CALEDONIA 12 II 1957
Charles P.Hoyt/ *Hyleorus hoyti* Mesnil L.P. Mesnil
det., 1976/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.
Paratype: 1♀, New Caledonia.
Remarks: Date of collection of holotype was published
as 12 February 1937.
- Hylotomomyia buckelli* Curran, 1926c, Can. Ent. 58: 216.
CANADA.
Holotype ♂: *Hylotomomyia* TYPE *buckelli* Curran No.
1952/ Osoyoos, B.C. 16.V.1924 E.R.Buckell.
- Hyperomyia rubidiapex* Reinhard, 1942b, Can. Ent. 74: 89. USA.
Holotype ♂: Utah 1927/ HOLOTYPE *Hyperomyia*
rubidiapex Reinhard/ *Hyperomyia rubidiapex*
det.H.J.Rein Rein/ Steveniopsis *rubidiapex* [sic] R
(Rnh.)/ Steveniopsis *rubidiapex* (Rnh.) Reinhard
Coll'n.
Allotype ♀: Wenatchee IX-15-28 Wash./ ALLOTYPE
Hyperomyia rubidiapex Reinhard/ Steveniopsis
rubidiapex (Rnh.) Reinhard Coll'n.
Paratypes: 1♂, California; 1♂, Utah; 1♂, Oregon; 1♀,
Washington.
- Hyperomyia subnitens* Reinhard, 1942b, Can. Ent. 74: 91. USA.
Holotype ♂: Garden City, Ut. July 31,1940 G. F.
Knowlton F. C. Harmston/ HOLOTYPE
- Hyperomyia subnitens* Reinhard/ *Hyperomyia*
subnitens det.H.J.Reinhard Rein/ Steveniopsis
subnitens R (Rnh.).
- Hyperecteina transcasica* Villeneuve, 1920c, Annls Soc. ent.
Belg. 60: 199. RUSSIA.
Lectotype ♀: Tr.Casp. Gr. Balchan/ *Hyperecteina*
transcasica n.sp./ *Hyperecteina transcasica* Vill.
L.P. Mesnil det., 1970/ TYPE/ CNC Syntype
Hyperecteina transcasica Villeneuve Label
affixed 1994/ Lectotype fixation by Mesnil,
1961b: 691. Label affixed 1995/ EX L.-P.
MESNIL COLLECTION 1970.
Paratypes: 4♀♀, Russia.
Remarks: We regard Mesnil's (1961b: 691) reference to
a holotype as a lectotype fixation for this species.
- Hypertophomma subita* Reinhard, 1958e, Ent. News 69: 240.
MEXICO.
Holotype ♀: Huajitlan, Morelos,Mex. 9-27-57
R.&K.Dreisbach/ HOLOTYPE *Hypertophomma*
subita Reinhard.
- Hypostena alberta* Curran, 1925b, Can. Ent. 57: 154. CANADA.
Holotype ♂: TYPE *Hypostena alberta* No Curran 4432/
Slave Lake, 2.VII.1924 Alta. O. Bryant.
Remarks: The CNC type number was published as
1303. This number appears on the type label but
has been crossed out and 4432 added.
- Hypostena columba* Curran, 1925b, Can. Ent. 57: 153. CANADA.
Holotype ♂: TYPE *Hypostena columba* 1310 No
Curran./ Eburne, B.C., 3.VI,1915. RS Sherman.
Paratype: 1♂, British Columbia.
- Hypostena imitator* Curran, 1925b, Can. Ent. 57: 154. CANADA.
Holotype ♂: HoloTYPE *Hypostena imitator* No Curran
1311/ Lillooet BC 15.VI.17 J.D.Tothill.
- Hypostena impatiens* Curran, 1925b, Can. Ent. 57: 154. CANADA.
Holotype ♂: TYPE *Hypostena impatiens* No. Curran
1312/ Teulon Man. 4.VIII.1922 A. J. Hunter.
- Hypostena (Arrhinomyia) ouelleti* Curran, 1925b, Can. Ent. 57:
150. CANADA.
Holotype ♂: TYPE *Hypostena oulettei* [sic] No. Curran
1307/ Joliette 9-VII Que.
Paratype: 1♂, Quebec.
- Hypostena (Arrhinomyia) tothilli* Curran, 1925b, Can. Ent. 57: 152.
CANADA.
Holotype ♂: TYPE *Hypostena tothilli* Curran No. 1309/
F'ton [Fredericton] 17.VII.12.
- Hypotachina fenestrata* Mesnil, 1971b, Fliegen palaearkt. Reg.
64g: 993. BURMA.
Holotype ♀: N. E. BURMA Kambaiti 7000m
28.IV.1934 R. MALAISÉ/ Dexiomima *fenestrata*
Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P.
MESNIL COLLECTION 1985.

Hypotachina raoi Mesnil, 1971b, Fliegen palaearkt. Reg. 64g: 992.
INDIA.

Holotype ♀: Shathla (H.P.) 13.7. 1963/ Kotgarh -S.S
C.I.B.C.-I.S./ Ex:Pupa of Gypsy moth(?) on oak/
Dexiomima raoi Mesnil L.P. Mesnil det., 1970/
TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Hypovoria hilaris Villeneuve, 1912b, Bull. Mus. natn. Hist. nat.,
Paris 18: 510. TUNISIA.
Holotype ♀: Tunisie Sfax 22-4-98/ Hypovoria hilaris
Villen. type/ Hypovoria hilaris Villen. L.P. Mesnil
det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Hystricia aldrichi Townsend, 1892a, Trans. Am. ent. Soc. 19: 91.
USA.
Syntype: 1♂.
♂: PARATYPE E. aldrichi Towns. No. 396/
Brookings SD/ JMA/ ParatypeNo 895 U.S.N.M./
CNC Syntype Hystricia aldrichi Townsend Label
affixed 1994.

Ignatomyia aurea Thompson, 1963d, Can. Ent. 95: 1308.
TRINIDAD.

Holotype ♂: HOLOTYPE Ignatomyia aurea Thomp.
CNC No. 9809/ St. Augustine, Trinidad,BWI.
NOV.10.1953/ Collector F.J.Simmonds/
HOLOTYPE CNCNo./ Ignatomyia aurea n.sp.
Allotype ♀: ALLOTYPE Ignatomyia aurea Thomp.
CNC No. 9809/ 2.17.60 Brazil Trinidad,BWI./
1062 II.18.60/ ALLOTYPE CNCNo.
Paratypes: 3♂♂, 1♀, Trinidad.

Imaguncula tabida Reinhard, 1958b, J. Kans. ent. Soc. 31: 230.
USA.
Holotype ♂: Tucson, Ar. VIII-16,1925/ HOLOTYPE
Imaguncula tabida Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 1♂, 1♀, Arizona; 2♀♀, Mexico.

Isidotus incanus Reinhard, 1962b, Pan-Pacif. Ent. 38: 216. USA.
Holotype ♂: BearCan.HH.Mi.11,Sta.
CatalinaMts.,ARIZ.VII 9-1959.B1 tr.FGWerner,
RWRadford&GASamuelson/ HOLOTYPE
Isidotus incanus Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 2♂♂, Arizona.

Istoglossa (Peristoglossa) aurantiaca Mesnil, 1973a, Fliegen
palaearkt. Reg. 64g: 1127. SENEGAL.
Holotype ♂: No 9174/ SENGAL Bambe 1940
v.Risbec/ Diptera des pommes de Acacia Sengal/
Peristoglossa aurantiaca Mesn L.P. Mesnil det.,
1971/ TYPE/ EX L.-P. MESNIL COLLECTION
1985.

Jalapamyia callitris Reinhard, 1964b, Acta zool. mex. 7: 13.
MEXICO.
Holotype ♀: Jalapa, Ver. 9/28-X/3/61 R & K
Dreisbach/ HOLOTYPE Jalapamyia callitris
Reinhard/ Jalapamyia callitris R Rnh.

Jurinia (Fabricia) hystricoides Williston, 1886, Trans. Am. ent.
Soc. 13: 300. USA.

Syntypes: 1♂, 1♀.
♂: D.C. Aug 10.83/ Type./ Jurinia PARATYPE
hystricoides Will. No.2030/ CNC Syntype Jurinia
(Fabricia) hystricoides Williston Label affixed
1994.
♀: Jurinia PARATYPE hystricoides Will.
No.2030/ Conn. Williston./ Type/ ?infuscatus
comp. to Ht ♀ Det.D.M.Wood 1989/ CNC
Syntype Jurinia (Fabricia) hystricoides Williston
Label affixed 1994.

Juriniopsis lampuris Reinhard, 1953a, Bull. Brooklyn ent. Soc. 48:
93. MEXICO.

Holotype ♂: Acatlpa, Morelos June 2, 1950
Coll.WGDowns/ HOLOTYPE Juriniopsis
lampuris Reinhard/ Juriniopsis lampuris R. Rnh.
Allotype ♀: Acatlpa Morelos July 29, 1950
Coll.WGDowns/ ALLOTYPE Juriniopsis lampuris
Reinhard.

Lasioneura dextella Reinhard, 1953a, Bull. Brooklyn ent. Soc. 48:
95. MEXICO.

Holotype ♂: alt. 9000' El Zarco,Mex IX/3/50 E.Bordas
Coll./ HOLOTYPE Lasioneura dextella Reinhard/
Lasioneura dextella R. Rnh.

Laufferiella elegans Villeneuve, 1929b, Bull. Annls Soc. ent. Belg.
69: 99. TURKMENIA.

Holotype ♂: Transasp Imam-baba/ Laufferiella
elegans Typ. Villen./ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Laximasicera sexualis Curran, 1927c, Can. Ent. 59: 14. CANADA.

Holotype ♂: Hemimasicera TYPE sexualis Curran. No.
2242/ Waterton Alta. July 10 1923 E.H.Strickland.

Leschenaultia halisidotae Brooks, 1947, Can. Ent. 78 (1946): 176.
CANADA.

Holotype ♂: HoloTYPE ♂ Leschenaultia halisidotae
No. Brks 5690/ Rec: 7101 F.I.Survey 1942
Em(Incubator) 6.III. 1943/ ex H.caryae Galt.Ont.
Allotype ♀: Same data as holotype except with allotype
label and emerged 9 March 1943.
Paratypes: 9♂♂, 5♀♀, Ontario; 2♂♂, 1♀, Quebec; 1♂,
1♀, Ohio; 1♀, Massachusetts; 1♀, New York.
Remarks: There are additional specimens labeled as
paratypes but they are from localities not
mentioned in the original description. Holotype
puparium is pinned below specimen.

Leschenaultia hospita Reinhard, 1952b, Bull. Brooklyn ent. Soc.
47: 7. MEXICO.

Holotype ♂: Zamora, Mich. Aug. 27, 1947/ F. A.
Cowan M. R. Wheeler Collectors/ Reinhard
Leschenaultia hospita Reinhard/ Leschenaultia
hospita R Rnh.

Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 6♀♀, Mexico.

- Leschenaultia nuda* Thompson, 1963b, Studia ent. 6: 288.
TRINIDAD.
- Syntype: 1♀.
♀: Brazil II. 18. 61 Trinidad,BWI/ HOLOTYPE
Leschenaultia nuda Thomp. CNC No. 9787/ Type
♀/ *Leschenaultia nuda* n.sp./ CNC Syntype
Leschenaultia nuda Thompson Label affixed 1994.
- Leskia miranda* Mesnil, 1973a, Fliegen palaearkt. Reg. 64g: 1129.
JAPAN.
Holotype ♀: Tsurugi Ishikawa p. JPN. 9.X.'64
S.Takano/ 406/ *Leskia miranda* Mesnil L.P.
Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.
Paratype: 1♂, Japan.
- Leskiopalpus sanctaecrucis* Thompson, 1963a, Can. J. Zool. 41:
348. TRINIDAD.
Syntypes: 2♀♀.
♀: HOLOTYPE *Leskiopalpus sanctaecrucis*
Thomp. CNC No. 9764/ 1095 III.22.60/ CNC
Syntype *Leskiopalpus sanctaecrucis* Thompson
Label affixed 1994.
♀: 1085 e III.9.60/ CNC Syntype *Leskiopalpus*
santaecrucis Thompson Label affixed 1994.
- Lespesia clavipalpis* Thompson, 1966, Studia ent. 8 (1965): 377.
TRINIDAD.
Holotype ♀: HOLOTYPE *Lespesia clavipalpis* Thomp.
CNC No. 9829/ Brazil 3. 11. 60 Trinidad,BWI./
1087 III. 11. 60/ *Lespesia clavipalpis* n.sp.
Paratype: 1♀, Trinidad.
- Lespesia pollinosa* Thompson, 1966, Studia ent. 8 (1965): 380.
TRINIDAD.
Holotype ♀: HOLOTYPE *Lespesia pollinosa* Thomp.
CNC No. 9830/ St. Augustine, Trinidad,BWI. XI
2. 60/ *Lespesia pollinosa* n.sp.
Allotype ♂: II.24.64 St. Augustine, Trinidad, WI./
ALLOTYPE *Lespesia pollinosa* Thomp. CNC No.
9830.
Paratypes: 2♀♀, Trinidad.
- Leucostoma africanum* Villeneuve, 1920a, Revue zool. afr. 8: 155.
SOUTH AFRICA.
Syntype: 1♂, 1♀.
♂: Willowmore, Capland, 15:12:11, Dr. Brauns/
XIV/ CNC Syntype *Leucostoma africanum*
Villeneuve Label affixed 1994.
♀: Willowmore, Capland, 15:12:11, Dr. Brauns/
XIV/ *Leucostoma africanum*, Types ♀ Villen./
CNC Syntype *Leucostoma africanum* Villeneuve
Label affixed 1994.
- Leucostoma effrenatum* Reinhard, 1956a, J. Kans. ent. Soc. 29: 165.
USA.
Holotype ♂: Logan Dry Canyon Ut.May 26 1949
Wilford J. Hanson/ HOLOTYPE *Leucostoma*
effrenatum Reinhard.
Paratype: 1♀, Utah.
Remarks: Holotype terminalia are glued to locality
label.

- Leucostoma fallax* Reinhard, 1974, Can. Ent. 106: 1164. MEXICO.
Holotype ♀: Cordoba Mex Vera Cruz VII-13-1960/
Dreisbach Coll./ HOLOTYPE *Leucostoma fallax*
Reinhard/ *Leucostoma fallax* R Rnh.
- Leucostoma peccator* Reinhard, 1956a, J. Kans. ent. Soc. 29: 167.
BRAZIL.
Holotype ♂: 25 9 1937 Brasilien Nova Teutonia
27°11'W. 52°23' L. Fritz Plaumann/ HOLOTYPE
Leucostoma peccator Reinhard/ *Leucostoma*
peccator R Rnh.
Remarks: Holotype abdomen is glued to mount strip.
- Leucostoma perrarum* Reinhard, 1956a, J. Kans. ent. Soc. 29: 164.
USA.
Holotype ♂: Duchesne, Utah Aug 23 1948 G. F.
Knowlton/ HOLOTYPE *Leucostoma perrarum*
Reinhard/ *Leucostoma perrarum* R Rnh.
Paratypes: 1♂, Utah; 2♂♂, Arizona; 1♂, 2♀♀, Oregon;
4♂♂, California; 1♂, Nevada.
Remarks: Date of collection of holotype was published
as 23 August 1928.
- Leucostoma politifrons* Reinhard, 1974, Can. Ent. 106: 1164. USA.
Holotype ♂: Southwestern Res.Sta. Cochise Co.,
Arizona August 10-12, 1966 H. J. Reinhard/
HOLOTYPE *Parapsalida politifrons* Reinhard/
Parapsalida politifrons R Rnh./ desc. in
Leucostoma.
Paratypes: 1♂, Arizona; 3♂♂, California.
- Leucostoma vapulare* Reinhard, 1956a, J. Kans. ent. Soc. 29: 165.
USA.
Holotype ♀: Sagehen nr. Hobart Mills Calif. VI-25-54/
E.I.Schlinger Collector/ HOLOTYPE *Leucostoma*
vapulare Reinhard/ *Leucostoma vapulare* R Rnh.
Paratype: 1♀, California.
- Ligeria latigena* Wood, 1985, Mem. ent. Soc. Can. 132: 51.
CANADA.
Holotype ♂: Y.T. 17 km WNW Burwash Flats
11.VII.80 1830m Wood & Lafontaine/
HOLOTYPE *Ligeria latigena* Wood CNC No.
18431.
Allotype ♀: Chesterfield N.W.T. 2.VIII.1960 J.G.
Chillcott/ IDEMA illustration/ ALLOTYPE
Ligeria latigena Wood CNC No. 18431.
Remarks: Holotype terminalia are glued to pin below
specimen.
- Lindigopalpus bogotensis* Reinhard, 1974, Can. Ent. 106: 1165.
COLOMBIA.
Holotype ♀: Bogata (Cund.) 14. VIII, 38/ HOLOTYPE
Lindigopalpus bogotensis Reinhard/ *Lindigopalpus*
bogotensis R Rnh.
- Linnaemyia breviseta* Villeneuve, 1941a, Bull. Soc. ent. Fr. 46:
109. ZAIRE.
Holotype ♀: Kabare VIII.14/ *Linnaemyia breviseta*
Typ. Villen./ *Linnaemyia breviseta* Vill. L.P.
Mesnil det., 1970/ EX L.-P. MESNIL
COLLECTION 1970/ TYPE.

- Linnaemyia consobrina* Villeneuve, 1941a, Bull. Soc. ent. Fr. 46: 108. ZIMBABWE.
Holotype ♀: Vumba Mts., S. Rhodesia, 14.I.1937 Major Drysdale/ Linnaemyia consobrina Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: Date of collection of holotype was published as 3 January 1937.
- Linnaemyia eburneola* Villeneuve, 1935b, Revue Zool. Bot. afr. 27: 141. ZAIRE.
Holotype ♀: R [Ruwenzori] 2300 19.IV.14/ Linnaemyia eburneola typ. n.sp./ Linnaemyia eburneola Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Linnaemyia nigrescens* Curran, 1925d, Ent. News 36: 15. CANADA.
Holotype ♂: Holo TYPE ♂ Linnaemyia nigrescens No. Cn. 1251/ Hedley, B.C. 23.VII.1923 C.B. Garrett.
- Linnaemyia sororcula* Villeneuve, 1941a, Bull. Soc. ent. Fr. 46: 107. ZAIRE.
Syntype: 1♀.
♀: Congo - belge Eala-V -1935 J. Ghesquière/ Linnaemyia sororcula Cotyp. Villen./ Linnaemyia sororcula Vill. L.P. Mesnil det., 1970/ CNC Syntype Linnaemyia sororcula Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Linnaemyia speciosissima* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 52. JAPAN.
Holotype ♀: Kyoto: Hanase 24.VII.1951 S.Ito/ 230/ Linnaemyia speciosissima Mesnil. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Linnaemyia takanoi* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 51. JAPAN.
Holotype ♀: Obihiro S.Takano [underside of label] 30/VI'51/ 245/ Linnaemyia takanoi Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Linnaemyia tessellata* Brooks, 1944b, Can. Ent. 76: 198. CANADA.
Holotype ♂: Holo TYPE ♂ Bonellimyia tessellata No. Brks 5397/ Petite Riviere, N.S. 18-vii.1935 J.McDunnough.
Allotype ♀: Aylmer, Que. 28.IX.1924 C.H. Curran/ AlloTYPE ♀ B. tessellata Brks No. 5397.
Paratypes: 2♂♂, 1♀, Nova Scotia; 5♂♂, 2♀♀, New Brunswick; 2♂♂, 1♀, Quebec; 3♂♂, 2♀♀, Ontario; 2♀♀, Saskatchewan; 4♂♂, 3♀♀, Alberta; 2♂♂, British Columbia; 2♂♂, 1♀, Northwest Territories; 5♂♂, Vermont; 1♂, Washington; 1♂, Idaho; 1♂, New Hampshire; 1♂, Colorado; 1♀, Massachusetts; 1♀, Tennessee.
- Linnaemyia varia* Curran, 1925d, Ent. News 36: 14. CANADA.
Holotype ♂: HoloTYPE ♂ Linnaemyia varia Cn. No. 1250/ Hopedale Labr. 12.6 1924.
Allotype ♀: Same data as holotype except with allotype label and collected "28.VII 1922".

- Linnaemyia (Gymnochaetopsis) dumonti* Mesnil, 1971b, Fliegen palaearkt. Reg. 64g: 1018. TUNISIA.
Holotype ♂: parasite chenille Cerocala scapulosa Nefta 5-2-27/ TUNISIE NEFTA C. DUMONT 1927/ Linnaemyia dumonti Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Linnaemyia (Micropalpus) formosensis* Villeneuve, 1932b, Bull. Soc. ent. Fr. 37: 269. TAIWAN.
Holotype ♂: Formosa Sauter/ Kosempo 908. 1.20./ Linnaemyia formosensis Typ. Villen./ TYPE/ Linnaemyia vulpinoides Bar. L.P. Mesnil det., 1970/ HOLOTYPE ♂ of Linnaemyia formosensis Vill. Examined 1970 R.W. Crosskey.
- Linnaemyia (Micropalpus) obscurior* Villeneuve, 1934a, Revue Zool. Bot. afr. 25: 409. SOUTH AFRICA.
Syntype: 1♂.
♂: Marley 5-16 Durban/ Micropalpus obscurior Cotyp. Villen./ Linnaemyia obscurior Vill. L.P. Mesnil det., 1970/ Linnaemyia agilis Curran L.P. Mesnil det., 1970/ CNC Syntype Linnaemyia (Micropalpus) obscurior Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Linnaemyia (Micropalpus) pulchella* Villeneuve, 1934a, Revue Zool. Bot. afr. 25: 410. NIGERIA.
Syntype: 1♂.
♂: ♂/ Oshogbo, S.Nigeria. Dr.T.F.G.Mayer. Nov 1910/ Micropalpus Dr. Villeneuve det. pulchellus Typ. Villen./ Linnaemyia pulchella Vill. L.P. Mesnil det., 19/ TYPE/ CNC Syntype Linnaemyia (Micropalpus) pulchella Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Lispideosoma flavipes* Reinhard, 1943a, Can. Ent. 75: 165. USA.
Holotype ♀: Dane Co Wis. IX-13-27/ HOLOTYPE Lispideosoma flavipes Reinhard/ Lispideosoma flavipes R. Rnh.
Allotype ♂: Wisconsin 7-VII-1934/ ALLOTYPE Lispideosoma flavipes Reinhard.
- Lispideosoma triste* Reinhard, 1961, J. Kans. ent. Soc. 34: 208. MEXICO.
Holotype ♂: W.Slope Cortez Pass. Mex. 10000' 8-11-54 R.R.Dreisbach/ HOLOTYPE Lispideosoma triste Reinhard/ Lispideosoma triste R Rnh.
- Lissoglossa bequaerti* Villeneuve, 1912b, Bull. Mus. natn. Hist. nat., Paris 18: 506. ALGERIA.
Lectotype ♂: CapFalcon Oran ♂ Dr. J. Bequaert/ Lissoglossa Bequaerti type Villen./ Lissoglossa bequaerti Villen. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype Lissoglossa bequaerti Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1973: 1217. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: We regard Mesnil's (1973b: 1217) reference to a holotype as a lectotype fixation for this species.

- Lissoglossa taeniata* Villeneuve, 1912b, Bull. Mus. natn. Hist. nat., Paris 18: 506. ALGERIA.

Lectotype ♀: Sa Cruz-Oran Algeria Dr. J. Bequaert/
Lissoglossa taeniata type Villen./ Lissoglossa
taeniata Vill. L.P. Mesnil det., 1970/ TYPE/ CNC
Syntype Lissoglossa taeniata Villeneuve Label
affixed 1994/ Lectotype fixation by Mesnil, 1973:
1217. Label affixed 1995/ EX L.-P. MESNIL
COLLECTION 1970.

Paralectotypes: 2♀♀, Algeria.

Remarks: We regard Mesnil's (1973b: 1217) reference
to a holotype as a lectotype fixation for this
species.

Litophasia albisquama Villeneuve, 1932a, Bull. Annls Soc. ent.
Belg. 71 (1931): 242. SPAIN.

Holotype ♂: Spanien, Pyrenäen, Farga de Moles bei
Seo de Urgel, 850-1300m. 4.-10.6.30/ F. Diehl u.
G.F.Meyer leg., Eing. Nr. 133, 1930./ *Litophasia*
albisquama Villeneuve.

Lixophaga fallax Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24):
32. JAPAN.

Holotype ♂: Toyama TAKAGI-S./ *Lixophaga fallax*
n.sp. L.P. Mesnil det., 1963/ TYPE/ EX L.-P.
MESNIL COLLECTION 1970/ *Oswaldia hirsuta* (Mesn.)
det. H. Shima 1992.

Lixophaga opaca Reinhard, 1945, Can. Ent. 77: 30. USA.

Holotype ♂: Blewett, Wn. [Washington] VI-18 1933 J.
Wilcox, Col./ HOLOTYPE *Lixophaga opaca*
Reinhard/ *Lixophaga opaca* Rnh.
det.H.J.Reinhard/ Aplomyopsis opaca (Rnh.).
Allotype ♀: Yakima Wn. VIII-19-1926/ ALLOTYPIC
Lixophaga opaca Reinhard.
Paratypes: 3♂♂, Alberta.

Lixophaga remora Reinhard, 1953b, J. Kans. ent. Soc. 26: 102.
MEXICO.

Holotype ♀: HOLOTYPE *Lixophaga remora* Reinhard/
Acatlipa, Morelos Sept. 1, 1950 Coll. WGDowns.
Paratype: 1♀, Mexico.

Lixophaga scintilla Reinhard, 1953b, J. Kans. ent. Soc. 26: 101.
MEXICO.

Holotype ♂: Acatlipa Morelos July 29, 1950 Coll.
WGDowns/ HOLOTYPE *Lixophaga scintilla*
Reinhard.

Loewia (Fortisia) nudigena Mesnil, 1973b, Fliegen palaearkt. Reg.
64g: 1208. SWITZERLAND.

Syntype: 1♂.
♂: Delemont Tente 23.VIII.1968/ *Loewia*
nudigena Mesn. L.P. Mesnil det., 1972/ TYPE/
CNC Syntype *Loewia (Fortisia) nudigena* Mesnil
Label affixed 1994/ EX. L.-P. MESNIL
COLLECTION 1985.

Loewia (Fortisia) piliceps Mesnil, 1973b, Fliegen palaearkt. Reg.
64g: 1209. FRANCE.

Syntype: 1♂.
♂: PAU 18.VI.10 B.PYR./ *Loewia piliceps*
Mesnil L.P. Mesnil det., 1972/ TYPE/ CNC
Syntype *Loewia (Fortisia) piliceps* Mesnil Label
affixed 1994/ EX. L.-P. MESNIL COLLECTION
1985.

Lomacantha parra hispanica Mesnil, 1962, Fliegen palaearkt. Reg.
64g: 773. SPAIN.

Holotype ♂: Ceredilla J. ARIAS/ *Lomacantha*
hispanica Mesn. L.P. Mesnil det., 1963/ TYPE/
EX L.-P. MESNIL COLLECTION 1970.

Lomatacantha apicalis Mesnil, 1957, Mém. Soc. r. ent. Belg. 28:
25. JAPAN.

Holotype ♂: Sapporo 5-VII. 1942 Y.Nishijima/ 116/
Oswaldia apicalis Mesn. L.P. Mesnil det., 1969/
TYPE/ EX L.-P. MESNIL COLLECTION 1970/
Oswaldia apicalis (Mesnil) det. H. Shima, 1992.

Lomatacantha cinctella Mesnil, 1957, Mém. Soc. r. ent. Belg. 28:
24. JAPAN.

Holotype ♀: Obihiro S. Takano [underside of label]
21/VII'55/ 201/ *Lixophaga cinctella* Mesnil L.P.
Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Lophosiosoma bicornis Mesnil, 1973b, Fliegen palaearkt. Reg. 64g:
1212. TAIWAN.

Holotype ♂: Formosa Sauter/ Mt.Hoozan 1910. VIII./
Lophosiosoma bicornis Mesnil L.P. Mesnil det.,
1972/ TYPE/ EX. L.-P. MESNIL COLLECTION
1985.

Paratype: 1♀, Taiwan.

Lubutana mayeri Mesnil, 1955a, Annls Mus. r. Congo belge Sér.
8vo (Zool.) 40: 363. NIGERIA.

Holotype ♀: Oshogbo, S.Nigeria. Dr.T.F.G.Mayer.
1-13. Oct 1910/ *Lubutana mayeri* Mesnil L.P.
Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Lydella connecta Curran, 1925a, Can. Ent. 57: 286. CANADA.

Holotype ♂: TYPE *Lydella connecta* Curran No. 1430/
BigCrk., B.R.V. Mt. 7-13-18.

Lydella hyphantriae Tothill, 1922, Bull. Can. Dep. Agric. (n. ser.)
3: 43. CANADA.

Holotype ♀: TYPE No. 487/ Ex *Hyphantriae Agassiz*
BC. 21.VII.1918 J.E.S./ *Lydella hyphantriae*
Tothill.

Lydella immissa Reinhard, 1959a, Ent. News 70: 232. MEXICO.

Lectotype ♂: Zamora, Mich. [Michoacan] Aug. 27,
1947/ F. A. Cowan M. R. Wheeler Collectors/
Lydella/ Reinhard Collection/ LECTOTYPE ♂ of
Lydella immissa Reinhard designated 1984
D.M.Wood.

Paralectotype: 1♂, Mexico.

Remarks: For lectotype designation see Wood (1985:
100).

Lydella laterale Curran, 1925a, Can. Ent. 57: 284. CANADA.

Holotype ♂: Holotype *Lydella lateralis* No. Curran
1453/ lillooet BC 23.VII.'17/ J.D. Tothill 1500'.

Lydella lathami Curran, 1925a, Can. Ent. 57: 284. USA.

Holotype ♂: Holotype ♂ *Paradexodes lathami* No
Curran 1429/ Greenpoint, L.I. 6-10-12 Roy
Latham.

Lydella minor Curran, 1925a, Can. Ent. 57: 283. CANADA.

Holotype ♂: TYPE Lydella minor Curran No./ Covey Hill, Que. 24.VI 1924 G.S.Walley.

Lydella obscura Curran, 1925a, Can. Ent. 57: 285. CANADA.
Holotype ♂: TYPE Lydella obscura No. Curran 1432/
Truro, N.S. 1 Aug. 17.

Lydellina anorbitalis Mesnil, 1970, Mushi 44: 99. BENIN.
Holotype ♂: Agouagon (Dahomey)/ Lydellina
anorbitalis Mesnil L.P. Mesnil det., 1969/ TYPE/
EX L.-P. MESNIL COLLECTION 1970.

Lydellina frontalis Mesnil, 1970, Mushi 44: 100. GHANA.
Holotype ♂: GOLD COAST ABURI 1912-13
W.H.PATTERSON Lydellina frontalis Mesnil
L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
Remarks: Holotype puparium is pinned below
specimen.

Lydellina umbripennis Mesnil, 1970, Mushi 44: 100. ZAIRE.
Holotype ♂: Congo-belge Eala-I -1936 J.Ghesquière/
Lydellina umbripennis Mesnil. L.P. Mesnil det.,
1969/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.

Lypha frontalis Brooks, 1945b, Can. Ent. 77: 85. CANADA.
Holotype ♂: Ottawa, Ont. 24.VI-1941 A. Brooks/ Holo
TYPE ♂ Lypha frontalis No. Brks 5524.
Allotype ♀: Ottawa, Ont. 2. VI 1927 C.H.Curran/ Allo
TYPE ♀ Lypha frontalis No. Brks 5524.

Lypha fumipennis Brooks, 1945b, Can. Ent. 77: 85. CANADA.
Holotype ♂: Lillooet V.1.17 J.D.T. / 2000'/ HoloTYPE
♂ Lypha fumipennis No. Brks 5523.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 3♂♂, 2♀♀, British Columbia.

Lypha intermedia Brooks, 1945b, Can. Ent. 77: 86. CANADA.
Holotype ♂: HoloTYPE ♂ Lypha intermedia No. Brks
5525/ Ex. S. duplex Gogama, Ont/ Rec.: 5873
F.I.Survey 1942 Em.(Incubator) 25.1 1943.
Allotype ♀: Same data as holotype except with allotype
label.
Paratype: 1♀, Ontario.
Remarks: Holotype puparium is pinned below
specimen.

Lypha parva Brooks, 1945b, Can. Ent. 77: 86. CANADA.
Holotype ♂: Mer Bleue Ont. 10.V -193 G.E.Shewell 8/
HoloTYPE ♂ Lypha parva No. Brks 5556.
Allotype ♀: Mer Bleue, Ont. 3.VI-1938 A.R.Brooks/
AlloTYPE ♀ Lypha parva No. Brks 5556.
Paratype: 1♀, Ontario.

Lypha (Lyphosia) barbata Mesnil, 1957, Mém. Soc. r. ent. Belg.
28: 56. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label]
28/V '49/ 29/ Lyphosia barbata Mesnil L.P. Mesnil
det., 1967/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Macquartia (Albiniola) nudigena Mesnil, 1972, Fliegen palaearkt.
Reg. 64g: 1100. FRANCE.

Holotype ♂: Rambouillet 10 avril/ Macquartia nudigena
Mesn. L.P. Mesnil det., 1971/ TYPE/ EX L.-P.
MESNIL COLLECTION 1985.

Paratype: 1♀, France.

Macquartia (Javetia) hystrix Mesnil, 1972, Fliegen palaearkt. Reg.
64g: 1101. FRANCE.

Holotype ♂: Lautaret (H. A.) VII./ Macquartia hystrix
Mesnil L.P. Mesnil det., 1975/ TYPE/ EX L.-P.
MESNIL COLLECTION 1985.

Macromya ciniscula Reinhard, 1968a, Can. Ent. 100: 1287.
BRAZIL.

Holotype ♂: Nova Teutonia 27°11'S, 52°23'W Brazil,
300-500m. 1. 1966 Fritz Plaumann/ HOLOTYPE
Macromya ciniscula Reinhard/ perfect for
anthemon but scutal hair pale Det.D.M.Wood
1989.

Paratype: 1♂, Brazil.

Macromya crocata Reinhard, 1968a, Can. Ent. 100: 1283. USA.

Holotype ♂: Big Bend Park Brewster Co. Texas July 8
1937 Rollin H. Baker/ HOLOTYPE Macromya
crocata Reinhard.

Allotype ♀: Same data as holotype except with allotype
label and dated "July 1, 1937".

Paratypes: 12♂♂, 20♀♀, Arizona; 2♂♂, Texas.

Mactomyia fracida Reinhard, 1958b, J. Kans. ent. Soc. 31: 229.
USA.

Holotype ♂: Globe, Arizona III-20 1937 Parker./
HOLOTYPE Mactomyia fracida Reinhard.

Allotype ♀: Ventura 13-V-44 Cal./ ALLOTYPE
Mactomyia fracida Reinhard.
Paratype: 1♂, New Mexico.

Maracasimyia antennalis Thompson, 1968, Mem. ent. Soc. Can.
56: 141. TRINIDAD.

Holotype ♀: HOLOTYPE Maracasimyia antennalis
Thomp. CNC No. 9874/ Maracas V X.27.60
Trinidad,BWI/ 1374A 1.30.64.

Paratypes: 2♀♀, Trinidad.

Remarks: Holotype is missing terminalia.

Masiphymomyia alearis Reinhard, 1944b, J. Kans. ent. Soc. 17: 66.
USA.

Holotype ♂: College Station May 15, 1943 Tex/ H J
Reinhard Collector/ HOLOTYPE Masiphymomyia
learis Reinhard/ Masiphymomyia alearis R'43 Rnh.
Allotype ♀: Same data as holotype except with allotype
label and dated "May 7, 1938".

Masiphymomyia comosa Reinhard, 1944b, J. Kans. ent. Soc. 17: 67.
USA.

Holotype ♂: Boulder Colo VI-30 1932 M.T.James/
6600'/ HOLOTYPE Masiphymomyia comosa
Reinhard/ Masiphymomyia comosa R43 Rnh.
Paratypes: 1♂, California; 1♂, Alberta.

Masiphymomyia longicornis Reinhard, 1944b, J. Kans. ent. Soc. 17:
67. USA.

Holotype ♀: Brewster Co VIII-8-37 Tex/ HOLOTYPE
Masiphymomyia longicornis Reinhard/
Masiphymomyia longicornis R'43 Rnh.

Allotype ♂: Same data as holotype except with allotype label.
 Paratype: 1♀, Texas.

Masiphyomyia paralis Reinhard, 1944b, J. Kans. ent. Soc. 17: 67.
 USA.
 Holotype ♂: Otero Co. N.M. VII-17-1926/ HOLOTYPE
Masiphyomyia paralis Reinhard/ *Masiphyomyia paralis* R.'43 Rnh.
 Paratype: 1♂, New Mexico.

Masistylum stenommatum Wood, 1974a, Can. Ent. 106: 178.
 CANADA.
 Holotype ♀: HOLOTYPE *Masistylum stenommatum* [sic] CNC No./ Summit Lake,B.C. Mi392 Alaska Hwy 15. VII. 1959 4700' R.E.Leech/ IDEMA illustration.
 Allotype ♂: Summit Lake,B.C. Mi392 Alaska Hwy 8. VII. 1959 4500' R.E. Leech/ ALLOTYPE *Masistylum stenommatum*/ IDEMA illustration/ 194 IDEMA illustration.

Masphyoidea chaetosa Thompson, 1963d, Can. Ent. 95: 1313.
 TRINIDAD.
 Holotype ♀: HOLOTYPE *Masphyoidea chaetosa* Thomp. CNC No. 9811/ St. Augustine, Trinidad,BWI. OCT 30.1953/ Collector F.J.Simmonds/ *Masphyoidea* [sic] *chaetosa* n.sp./ HOLOTYPE CNCNo./ 1004a QQ/ Type/ QQ.
 Paratype: 1♀, Trinidad.

Mastigiomyia delusa Reinhard, 1964b, Acta zool. mex. 7: 18.
 MEXICO.
 Holotype ♂: Cuernavaca,Mex. March 22 1934 S. E. Jones,Coll./ HOLOTYPE *Mastigiomyia delusa* Reinhard/ *Mastigiomyia delusa* R Rnh.
 Paratypes: 2♂♂, Mexico.

Mauromyia finitina Reinhard, 1967, J. Kans. ent. Soc. 40: 97.
 CANADA.
 Holotype ♀: RENNIE MAN. P. 12.9.61 Coll/ 49/ HOLOTYPE *Mauromyia finitina* Reinhard/ *Mauromyia finitina* R Rnh.
 Allotype ♂: RENNIE MAN. P. 28.8.61 Coll/ 47/ ALLOTYPE *Mauromyia finitina* Reinhard.
 Paratypes: 4♀♀, Manitoba.
 Remarks: There are five females labeled as paratypes, but only four paratypes are mentioned in the description.

Medina abdominalis Mesnil, 1971a, Entomophaga 16: 72. JAPAN.
 Holotype ♀: Napporo Japan Natural forest 26.8.1961/
Medina abdominalis Mesn. L.P. Mesnil det., 1964/
 TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Meigenia bellina Mesnil, 1967, Mushi 41: 44. JAPAN.
 Holotype ♂: Hikosan Kyushu-Japon 27.VI.1961/
Meigenia bellina Mesn. L.P. Mesnil det., 1964/
 TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Meigenia uncinata Mesnil, 1967, Mushi 41: 44. SWITZERLAND.

Holotype ♂: Suisse-Neuchâtel Bevaix 12 VIII 64 J. de Beaumont/ *Meigenia uncinata* Mesn. L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Melanopalpus meracula Reinhard, 1974, Can. Ent. 106: 1166.
 COLOMBIA.
 Holotype ♀: Pmo. Guerrero, Cundinamarca, Alt. 3240 K67 Fech. X-47 Col. [collector's name illegible]/ HOLOTYPE *Melanopalpus* [sic] *meracula* Reinhard/ *Melanopalpus meracula* R Rnh.

Meledonus albiceps Reinhard, 1956c, Ent. News 67: 126. USA.
 Holotype ♂: TrinityCoCal Eagle Creek VI.1 1941/
 HOLOTYPE *Meledonus albiceps* Reinhard.

Meledonus lindensis Reinhard, 1953c, Pan-Pacif. Ent. 29: 57. USA.
 Holotype ♂: Lind WASH 1 June 1919 R.W.Carlson/
 HOLOTYPE *Meledonus lindensis* Reinhard/
Meledonus lindensis R. Rnh.
 Allotype ♀: Lind WASH 12 June 1919 R.W.Carlson/
 HOLOTYPE *Meledonus lindensis* Reinhard.

Meleterus nuperus Reinhard, 1956c, Ent. News 67: 123. USA.
 Holotype ♂: Yonah Mt. Ga., 8-20-35/ HOLOTYPE
Meleterus nuperus Reinhard/ *Meleterus nuperus* R Rnh.
 Paratype: 1♂, Georgia.

Meriania septentrionalis Curran, 1930a, Jl N.Y. ent. Soc. 38: 75.
 CANADA.
 Holotype ♂: Vernon,B.C. 28. IV 1924 E.R.Buckell/
Meriania TYPE *septentrionalis* [sic] Curran ♂ No./
 TYPE ♂ *Meriania septentrionalis* No. 3215 Curran.
 Allotype ♀: Creston,B.C. 19/5 1924 C.S.Lallamand/
Meriania TYPE *septentrionalis* Curran. No. ♀/
 TYPE ♀ *Meriania septentrionalis* 3215 No.Curran.

Mericia alberta Curran, 1924a, Ent. News 35: 248. CANADA.
 Holotype ♂: HOLOTYPE *Mericia alberta* Curr No.798/ Banff, Alta. 30 VI 1922 C.B.D.Garrett 6000 ft.
 Allotype ♀: TYPE *M.alberta* Curr. No.798/ Banff,Alta. 16.VIII 1922 C.B.D.Garrett/ *M.alberta* Det.A.Brooks-42.
 Remarks: Holotype terminalia are glued to locality label. The CNC type number was published as 698.

Mericia campestris Curran, 1924a, Ent. News 35: 249. CANADA.
 Holotype ♂: TYPE *Mericia campestris* Curran No. 795/ P N. Vroom Aweme, Man. 20-V-1920.
 Paratype: 1♂, Manitoba.

Mericia cobala Reinhard, 1953c, Pan-Pacif. Ent. 29: 55. USA.
 Holotype ♂: OTERO CO. N.MEX./ Jun 20/ 8000 ft.
 Elevation/ E.M.& R.H.Painter Collectors/
 HOLOTYPE *Mericia cobala* Reinhard/
 ALLOTYPE *Mericia cobala* Reinhard/ *Mericia cobala* R Rnh.
 Allotype ♀: Mounted on same pin as holotype.

Mericia fasciventris Curran, 1924a, Ent. News 35: 248. CANADA.

- Holotype ♂: AylmerQue. 1-VI-1923 C.H.Curran/
HoloTYPE Mericia fasciventris Curr No. 797.
Paratype: 1♂, British Columbia.
Remarks: Holotype terminalia are glued to locality
label.
- Mericia hamilla* Reinhard, 1953c, Pan-Pacif. Ent. 29: 54. USA.
Holotype ♂: MtRainierWn [Washington] Sunrise6380'/
J.Wilcox Coll. VII-31-32/ HOLOTYPE Mericia
hamilla Reinhard.
Allotype ♀: Puyallup, Wn. VI-1 1933 J.Wilcox, Coll./
ALLOTYPE Mericia hamilla Reinhard.
Paratype: 1♂, Washington.
- Mericia manitoba* Brooks, 1943, Can. Ent. 75: 77. CANADA.
Holotype ♂: Aweme,Man. N.Criddle 10 VII 1928/
HoloTYPE ♂ Mericia manitoba No. Brks 5328.
Allotype ♀: Teulon Man. 17.VIII 1923 A.J.Hunter/
AlloTYPE ♀ Mericia manitoba No. Brooks 5328.
- Mericia occidentalis* Brooks, 1943, Can. Ent. 75: 77. CANADA.
Holotype ♂: Creston,B.C. 28.V. 1926 A.A.Dennys/
HoloTYPE ♂ Mericia occidentalis No. Brooks
5329.
Allotype ♀: Hedley,B.C. 29.VIII 1923 C.B.Garrett/
AlloTYPE ♀ Mericia occidentalis No. Brooks
5329.
- Mericia triangularis* Curran, 1924a, Ent. News 35: 247. CANADA.
Holotype ♂: HoloTYPE Mericia triangularis Curr No.
799/ P.N.Vroom Aweme, Man. 5-24-1920.
Remarks: Holotype terminalia are glued to locality
label, and puparium is pinned below specimen.
- Messiomyia triconis* Reinhard, 1955a, J. Kans. ent. Soc. 28: 125.
USA.
Holotype ♂: TEX./ HOLOTYPE Messiomyia triconis
Reinhard/ Messiomyia triconis Rnh. det.
H.J.Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratype: 1♂, Texas.
- Metagonistylum sesamiae* Mesnil, 1968b, Stuttg. Beitr. Naturk.
187: 4. UGANDA.
Holotype ♀: 1203 SERERE Kidetok 16.Jan. 1957
W.R.Ingram ex.S.orialua/ COLL. NO 3168
IDENT.NO/ Metagonistylum sesamiae Mesn. L.P.
Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.
Paratype: 1♀, Uganda.
Remarks: Holotype puparium is pinned below
specimen.
- Metamyia auranticauda* Arnaud, 1963, Am. Mus. Novit. 2157: 10.
USA.
Holotype ♂: Upper Lake VIII.8.59 Cal/ Light Trap
Collecting/ R.E.Dolphin Collector/ Cacozelus
sannionis R Rnh./ HOLOTYPE Metamyia ♂
auranticauda P.H.Arnaud/ Metamyia ♂
auranticauda det Arn. PHArnaud '62/ Reinhard
Collection.
- Metaphorocera maracasi* Thompson, 1968, Mem. ent. Soc. Can.
56: 18. TRINIDAD.
HoloTYPE Metaphorocera maracasi
Thomp. CNC No. 9842/ CUREPE TRINIDAD 7-7
1964/ F.D.BENNETT Collector.
Allotype ♀: U. Manzanilla I. 26. 61. Trinidad, BWI/
ALLOTYPE Metaphorocera maracasi Thomp.
CNC No. 9842.
Paratype: 1♀, Trinidad.
- Metaplagia brevicornis* Brooks, 1945b, Can. Ent. 77: 81.
CANADA.
Holotype ♂: Teulon Man. 11.VII 1923 A. J. Hunter/
HoloTYPE ♂ Metaplagia brevicornis No. Brks
5519.
Paratypes: 1♂, Prince Edward Island; 1♂,
Saskatchewan; 1♂, Northwest Territories; 1♂,
Massachusetts.
- Metavoria facialis* Reinhard, 1956c, Ent. News 67: 123. USA.
Holotype ♀: Fruitland,Ut. Aug 24, 1948 G.F.
Knowlton/ HOLOTYPE Metavoria facialis
Reinhard.
- Metavoria latifrons* Reinhard, 1956c, Ent. News 67: 121. USA.
Holotype ♂: Big Bend Pk VII-14 37 Tx/ HOLOTYPE
Metavoria latifrons Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratype: 1♂, Texas.
- Metopomuscopteryx fatigantis* Reinhard, 1958d, J. Kans. ent. Soc.
31: 284. USA.
Holotype ♂: Temescal Canyon Los Angeles Co Cal.
II-7-54/ W.McDonald Collector/ HOLOTYPE
Metopomuscopteryx fatigantis Reinhard.
Remarks: There may be a paratype among the CNC
material, but it is not labeled as such.
- Metopomuscopteryx incurata* Reinhard, 1958d, J. Kans. ent. Soc.
31: 283. USA.
Holotype ♂: Oak Creek Can. ARIZONA Mar. 31, 1956
FWerner,GButler/ HOLOTYPE
Metopomuscopteryx incurata Reinhard/
Metopomuscopteryx incurata R. Rnh.
Allotype ♀: Same data as holotype except with allotype
label and without last label.
- Microchaetina setifacies* Reinhard, 1942b, Can. Ent. 74: 88. USA.
Holotype ♂: Manhattan, Ks. Oct 15,1926/ R.H.Painter
Coll/ HOLOTYPE Microchaetina setifacies
Reinhard/ Microchaetina setifacies det.H.J.Rein
Rein/ Microchaetina mexicana R Tns.
Paratype: 1♂, Kansas.
Remarks: Date of collection of holotype was published
as 16 October 1926.
- Microleskia longipes* Thompson, 1968, Mem. ent. Soc. Can. 56:
143. TRINIDAD.
HoloTYPE Microleskia longipes Thomp.
CNC No. 9875/ Simla Trinidad/ Microleskia
flavipes n sp.
- Microphthalma ascita* Reinhard, 1953a, Bull. Brooklyn ent. Soc.
48: 89. MEXICO.

Holotype ♂: Acatlipa Morelos IX-16-50
Coll.WGDowns/ On rocky Hillside/ HOLOTYPE
Microphthalma ascita Reinhard/ Microphthalma
ascita R Rnh.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 3♂♂, Mexico.

Microphthalma europaea var. *nigeriensis* Villeneuve, 1935b,
Revue Zool. Bot. afr. 27: 137. NIGERIA.
Holotype ♂: Ikot-ekpene. S.Nigeria. J.J.Simpson.
19.5.1910/ Microphthalma disjunctum W. var.
nigeriense n. var. Villen. typ. [underside of label]
26 Mai 1935/ Microphthalma nigeriense Vill. L.P.
Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Microphthalma europaea numidica Mesnil, 1967, Mushi 41: 49.
ALGERIA.
Holotype ♂: ex Phyllognathus silenus Mostaganem/
Microphthalma numidica Mesn L.P. Mesnil det.,
1967/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.

Microphthalma pedalis Reinhard, 1953a, Bull. Brooklyn ent. Soc.
48: 90. MEXICO.
Holotype ♂: Chipinque, N.L. 22/IX/50 L.Navarro Coll.
WGDowns alt. 1440/ HOLOTYPE Microphthalma
pedalis Reinhard/ Microphthalma pedalis R. Rnh.
Paratype: 1♂, Mexico.

Microphthalma phyllophagae Curran, 1925d, Ent. News 36: 16.
CANADA.
Holotype ♂: HoloTYPE Microphthalma phyllophagae
Curr No. 1224/ Hemmingford, Que. 15.VIII. 1924
G.H.Hammond.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 44♂♂, 24♀♀, Quebec.

Milonioides scordalus Reinhard, 1955a, J. Kans. ent. Soc. 28: 127.
USA.
Holotype ♂: Dilley, Tex V-21-1920/ HOLOTYPE
Milonioides scordalus Reinhard.

Mimologus effetus Reinhard, 1955a, J. Kans. ent. Soc. 28: 130.
USA.
Holotype ♂: Tanbark Flat Los Angeles Co
Calif.VII-12-1952/ D.E. Barcus Collector/
HOLOTYPE Mimologus effectus [sic] Reinhard/
Mimelogus [sic] effetus R Rnh.
Paratypes: 3♂♂, California.

Mimologus emulatus Reinhard, 1962a, Ent. News 73: 177.
MEXICO.
Holotype ♀: WLNutting&FGWerner
10Mi.E.Navajoa,SON. MEX.VIII-13-1959./
HOLOTYPE Mimologus emulatus Reinhard.
Allotype ♂: Same data as holotype except with allotype
label.

Minella americana Reinhard, 1943b, J. Kans. ent. Soc. 16: 14.
USA.

Holotype ♂: Amherst, Ohio V-31 1931/ H J Reinhard
Collector/ HOLOTYPE Minella americana
Reinhard/ Minella americana det.H.J.Rein Rein/
IDEMA illustration.
Remarks: Collector of holotype was published as A.J.
Barckert.

Minthodes rhodesiana Villeneuve, 1942a, Bull. Soc. ent. Fr. 47: 54.
ZIMBABWE.

Syntypes: 2♀♀.
♀: GOTA GOTA URUNGWE Dept. Agric. S.
Rhodesia 16/9/1938/ W. L. Williams Collector/
Minthodes rhodesiana Typ. Villen./ Minthodes
rhodesiana Vill. L.P. Mesnil det., 1970/ TYPE/
CNC Syntype Minthodes rhodesiana Villeneuve
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

♀: GOTA GOTA URUNGWE Dept. Agric. S.
Rhodesia 23/9/1938/ W. L. Williams Collector/
Minthodes rhodesiana Vill. L.P. Mesnil det., 1970/
CNC Syntype Minthodes rhodesiana Villeneuve
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

Remarks: Second syntype is missing head and
abdomen.

Minthodexiopsis elegans Reinhard, 1964a, J. Kans. ent. Soc. 37: 37.
USA.

Holotype ♂: Cameron County 11-4 1930 Tex/ H J
Reinhard Collector/ HOLOTYPE Minthodexiopsis
elegans Reinhard/ Minthodexiopsis elegans R
Rnh.
Allotype ♀: Jalapa, Ver., Mex. VIII/1-6/61 R & K
Dreisbach/ ALLOTYPE Minthodexiopsis elegans
Reinhard.
Paratype: 1♀, Arizona.

Minthozelia argentina Reinhard, 1946c, J. Kans. ent. Soc. 19: 55.
USA.

Holotype ♂: El Paso,Tx VIII-6-1919/ HOLOTYPE
Minthozelia argentina Reinhard/ Minthozelia
argentina Rnh. Reinhard Coll'n/ Minthozelia
argentina R. Rnh.
Allotype ♀: BEXAR CO., TEX 6-8 , 1928 Parks/ 12/
ALLOTYPE Minthozelia argentina Reinhard/
Minthozelia montana A.) Tns./ Minthozelia
argentina Rnh. Reinhard Coll'n.
Paratype: 1♂, Texas.

Minthozelia metalis Reinhard, 1946c, J. Kans. ent. Soc. 19: 57.
USA.

Holotype ♂: Amherst, Ohio VII-8 1922/ AJ Barckert
Coll./ HOLOTYPE Minthozelia metalis Reinhard/
Minthozelia metalis R. Rnh.
Remarks: There may be a paratype among the CNC
material, but it is not labeled as such.

Minthozelia mira Reinhard, 1946c, J. Kans. ent. Soc. 19: 56. USA.

Holotype ♀: College Station 6.19.20 Texas/ H J
Reinhard Collector/ HOLOTYPE Minthozelia
mira Reinhard/ Minthozelia mira R. Rnh.

Minthozelia ruficauda Reinhard, 1946c, J. Kans. ent. Soc. 19: 58.
USA.

Holotype ♀: Forest Hill, L.I. July 4, 27 N.Y./
 HOLOTYPE *Minthozelia ruficauda* Reinhard/
Minthozelia ruficauda R. Rnh.
 Paratype: 1♀, New York.

Mochlosoma demissum Reinhard, 1958c, Can. Ent. 90: 101.
 MEXICO.
 Holotype ♂: Rio Frio, Mex. Sept. 9, 1947/ F. A. Cowan
 M. R. Wheeler Collectors/ HOLOTYPE
Mochlosoma demissum Reinhard/ *Mochlosoma demissum* R Rnh.
 Allotype ♀: Same data as holotype except with allotype label and without last label.
 Paratypes: 3♂♂, Mexico.

Mochlosoma duplare Reinhard, 1958c, Can. Ent. 90: 100.
 MEXICO.
 Holotype ♂: Cuernavaca, 7-25-47 Mex/ F. A. Cowan
 M. R. Wheeler Collectors/ HOLOTYPE
Mochlosoma duplare Reinhard/ *Mochlosoma duplare* R Rnh.
 Allotype ♀: Same data as holotype except with allotype label and without last label.
 Paratypes: 7♂♂, 7♀♀, Mexico; 1♂, 1♀, Arizona.

Mochlosoma illocale Reinhard, 1958c, Can. Ent. 90: 102. USA.
 Holotype ♂: Antelope Mt. Grant Co., Or. Aug. 6, 1932/
 Elevation 5000 ft./ DKFrewing Coll./ HOLOTYPE
Mochlosoma illocale Reinhard/ *Mochlosoma illocale* R Rnh.
 Allotype ♀: Bly, Oreg. Klamath Co. Oct. 13, 1943
 R.E.Rieder.col./ ALLOTYPE *Mochlosoma illocale* Reinhard.
 Paratypes: 2♂♂, Oregon; 1♂, Ohio; 3♂♂, Montana; 1♂,
 Wyoming; 1♂, California; 5♀♀, Utah; 1♂, 1♀,
 British Columbia.

Mochlosoma indutile Reinhard, 1958c, Can. Ent. 90: 104. USA.
 Holotype ♂: Martinez, Calif. Contra Costa Co.
 IX-14-46 H.E.Cott/ HOLOTYPE *Mochlosoma indutile* Reinhard/ *Mochlosoma indutile* R Rnh.
 Paratype: 1♀, Arizona.

Mochlosoma laudatum Reinhard, 1958c, Can. Ent. 90: 106.
 MEXICO.
 Holotype ♂: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan
 M. R. Wheeler Collectors/ HOLOTYPE
Mochlosoma laudatum Reinhard/ *Mochlosoma laudatum* R Rnh.
 Allotype ♀: Xochimilco, D.F. W.G.Downs, Coll./
 ALLOTYPE *Mochlosoma laudatum* Reinhard.
 Paratypes: 12♂♂, 1♀, Mexico.

Mochlosoma mentitum Reinhard, 1964a, J. Kans. ent. Soc. 37: 35.
 USA.
 Holotype ♀: Amherst Ohio Aug. 1958 H.J.Reinhard/
 HOLOTYPE *Mochlosoma mentitum* Reinhard/
Mochlosoma mentitum R Rnh.

Mochlosoma russulum Reinhard, 1958c, Can. Ent. 90: 105.
 MEXICO.
 Holotype ♂: Mixquic, D.F. Aug. 18 '49 Coll.
 WGDowns/ HOLOTYPE *Mochlosoma russulum* Reinhard/ *Mochlosoma russulum* R Rnh.

Allotype ♀: Mixquic, D.F. Aug. 18, 1949 Coll.
 WGDowns/ VIII/18/49-1 genitalia mounted/
 ALLOTYPE *Mochlosoma russulum* Reinhard.
 Paratypes: 1♂, 1♀, Mexico.
 Remarks: Allotype terminalia have not been located.

Mochlosoma serotinum Reinhard, 1958c, Can. Ent. 90: 101.
 MEXICO.
 Holotype ♂: Cuernavaca, 7-25-47 Mex/ F. A. Cowan
 M. R. Wheeler Collectors/ HOLOTYPE
Mochlosoma serotinum Reinhard/ *Mochlosoma serotinum* R Rnh.
 Paratypes: 4♂♂, Mexico.

Montserratia ovipara Thompson, 1964, Studia ent. 7: 149.
 TRINIDAD.
 Holotype ♀: HOLOTYPE *Montserratia ovipara* Thomp.
 CNC No. 9826/ III.19.64 Trinidad, WI/ 1423
 III.18.64/ *Montserratia ovipara* n.g. n.sp.

Mormonomya brunnicosa Villeneuve, 1935a, Bull. Soc. ent. Fr. 40: 252. SOUTH AFRICA.
 Holotype ♂: Port Elizabeth Sôd africás 15, 95 Dr.
 Brauns/ G. Mormonomyia/ Mormonomyia brunnicosa n.sp./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Mormonomya claripennis Villeneuve, 1935a, Bull. Soc. ent. Fr. 40: 253. SOUTH AFRICA.
 Syntype: 1♂.
 ♂: reducta n.sp./ Brit.E.Exp Nakuta Jan 1913 Dr.
 B.L. vanSomeren/ TYPE/ Mormonomyia claripennis n.sp./ CNC Syntype Mormonomyia claripennis Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.

Mormonomya fumosa Villeneuve, 1935a, Bull. Soc. ent. Fr. 40: 253. ZIMBABWE.
 Syntype: 1♂.
 ♂: Bulawayo S. Rhodesia 7. 10. 1923
 Coll.R.Stevenson/ Mormonomyia fumosa n.sp./ TYPE/ CNC Syntype Mormonomyia fumosa Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.

Mormonomya leucodes Villeneuve, 1935a, Bull. Soc. ent. Fr. 40: 252. SOUTH AFRICA.
 Holotype ♂: 641/ Mormonomyia leucodes n.sp./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Mormonomya nigrofimbriata Villeneuve, 1935a, Bull. Soc. ent. Fr. 40: 252. SOUTH AFRICA.
 Holotype ♂: Transvaal Kluksdorp 12. 90./
Mormonomya capensis Schir./ Mormonomyia nigrofimbriata n.sp./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
 Remarks: Holotype head is glued to the species label.

Mormonomya umbrosa Villeneuve, 1935a, Bull. Soc. ent. Fr. 40: 252. SOUTH AFRICA.
 Holotype ♂: Pretoria, Tvl. 2.i.1922./ Mormonomyia umbrosa n.sp./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Murana pauciseta Mesnil, 1980, Fliegen palaearkt. Reg. 64f: 11.
FRANCE.

Holotype ♂: Hautes-Pyrénées VII - VIII/ Murana
pauciseta Mesn. L.P. Mesnil det., 1978/ TYPE/
EX L.-P. MESNIL COLLECTION 1985.

Muscopteryx hiemalis Reinhard, 1944a, Ann. ent. Soc. Am. 37:
355. USA.

Holotype ♂: Madison VI-11-25 Wis/ HOLOTYPE
Muscopteryx hiemalis Reinhard.

Muscopteryx hilaris Reinhard, 1944a, Ann. ent. Soc. Am. 37: 355.
USA.

Holotype ♂: College Station 3-15-37 Texas/ H J
Reinhard Collector/ HOLOTYPE *Muscopteryx*
hilaris Reinhard/ *Muscopteryx hilaris* R Rnh.
Paratype: 1♂, Texas.

Muscopteryx longiseta Reinhard, 1944a, Ann. ent. Soc. Am. 37:
355. USA.

Holotype ♂: Calif. V-19-1914/ HOLOTYPE
Muscopteryx longiseta Reinhard.

Muscopteryx parilis Reinhard, 1944a, Ann. ent. Soc. Am. 37: 356.
USA.

Holotype ♂: Hartley Co VI-14-34 Tx/ HOLOTYPE
Muscopteryx parilis Reinhard/ *Muscopteryx*
parilis R. Rnh./ *Muscopteryx parilis* Rnh.
Reinhard Coll'n.

Muscopteryx petensis Reinhard, 1958d, J. Kans. ent. Soc. 31: 280.
USA.

Holotype ♂: Catalina Mts IV-30-41 Ariz/ HOLOTYPE
Muscopteryx petensis Reinhard/ *Muscopteryx*
petensis R. Rnh.

Allotype ♀: Same data as holotype except with allotype
label and without last label.

Paratypes: 3♂♂, 1♀, Arizona.

Myiobia longirostris Villeneuve, 1937a, Revue Zool. Bot. afr. 29:
210. SOUTH AFRICA.

Holotype ♀: Knyzna C.C. Oct. 1916. L. Peringuey/
longirostris Villen. Typ./ Fischeria longirostris
Villen. L.P. Mesnil det., 1970/ TYPE/
HOLOTYPE ♀ of *Myiobia longirostris* Vill.
Examined 1970 R.W.Crosskey.

Myiobia siphonina Villeneuve, 1937a, Revue Zool. Bot. afr. 29:
209. GUYANA.

Syntypes: 2♀♀.

♀: In Borings of Diatraea/ Pln Golden Fleece
Essequibo. B. Guiana 13. iii .13/ GE Bodkin
Collector/ Sugar Cane/ *Myiobia siphonina* Typ.
Villen./ TYPE/ CNC Syntype *Myiobia siphonina*
Villeneuve Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.
♀: In Borings of Diatraea/ Pln Golden Fleece
Essequibo. B. Guiana May 1913/ GE Bodkin
Collector/ Sugar Cane/ *Myiobia* (*Stomatodexia*)
siphonina Cotyp. Villen./ CNC Syntype *Myiobia*
siphonina Villeneuve Label affixed 1994/ EX
L.-P. MESNIL COLLECTION 1970.

Remarks: Each syntype has its puparium pinned below
specimen.

Myioclonia albertana Reinhard, 1945, Can. Ent. 77: 29. CANADA.

Holotype ♀: Lethbridge, Alta JUNE 6 1923 H.E.Gray/
Holotype M. *albertana* Reinhard/ HOLOTYPE ♀
Myioclonia albertana No. Rnh. 5498/ *Myioclonia*
albertana mss.Rnh.
Paratype: 1♀, Alberta.

Myioclonia erythrocerata Reinhard, 1945, Can. Ent. 77: 29. USA.

Holotype ♀: Smoky Mts June 33 Tenn./ HOLOTYPE
Myioclonia erythrocerata Reinhard/ *Myioclonia*
erythrocerata det.H.J.Rein Rnh.
Paratype: 1♀, Ontario.

Myioclonia nigricornis Reinhard, 1945, Can. Ent. 77: 30. USA.

Holotype ♀: Electron, Wn. [Washington] VI.2 1933/ J.
Wilcox, Coll./ HOLOTYPE *Myioclonia*
nigricornis Reinhard/ *Myioclonia nigricornis*
det.H.J.Reinhard Rnh.

Myiopharus canadensis Reinhard, 1945, Can. Ent. 77: 31.
CANADA.

Holotype ♂: HoloTYPE ♂ *Myiopharus canadensis* No.
Rnh. 5500/ Holotype M. *canadensis* Reinhard/
Berthierville Que. 4-7 1936 L.Daviault.
Allotype ♀: Parasite No 6 adulte Calli. bigbyana/
Berthierville Que. 9-7 1936 L.Daviault/
ALLOTYPE M. *canadensis* Reinhard/ AlloTYPE
♀ *Myiopharus canadensis* No. Rnh. 5500.
Paratypes: 1♂, 1♀, Quebec; 2♂♂, Ontario.

Myiopharus securis Reinhard, 1945, Can. Ent. 77: 30. USA.

Holotype ♂: TEXAS MAY 16.'27/ HOLOTYPE
Myiopharus securis Reinhard/ *Myiopharus securis*
Rein. det. H.J.Reinhard.
Allotype ♀: Taylor Tex 5-13-1929/ H J Reinhard
Collector/ ALLOTYPE *Myiopharus securis*
Reinhard.

Myiophasia harpi Reinhard, 1974, Can. Ent. 106: 1168. USA.

Holotype ♂: Junction, Texas Kimble County
Aug.-Sept., 1969 ex. Curculio caryae S. J. Harp/
HOLOTYPE *Myiophasia harpi* Reinhard/
Myiophasia harpi R Rnh.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 5♂♂, 7♀♀, Texas.

Myiophasia lasia Reinhard, 1959a, Ent. News 70: 225. USA.

Holotype ♂: Signal Mt., Wyo. Aug. 3, 1947/ F. A.
Cowan M. R. Wheeler Collectors/ HOLOTYPE
Myiophasia lasia Reinhard.
Paratypes: 1♂, Washington; 1♂, Oregon; 1♂, Utah.

Myiophasia sigilla Reinhard, 1959a, Ent. News 70: 226. USA.

Holotype ♂: Sny [Snowy] Range Wyo 11600ft 8/13/57/
H J Reinhard Collector/ HOLOTYPE *Myiophasia*
sigilla Reinhard.

Myiothyria benoisti Mesnil, 1959, Stuttg. Beitr. Naturk. 23: 20.

MOROCCO.
Holotype ♂: 10 MAI 28/ *Myiothyria benoisti* Mesnil
L.P. Mesnil det., 1967/ TYPE/ Ceracia
mucronifera Rond. L.P. Mesnil det., 1970/ EX
L.-P. MESNIL COLLECTION 1970.

Mystacomyoidea mirabilis Thompson, 1963d, Can. Ent. 95: 1318.
TRINIDAD.

Syntype: 1♀.

♀: HOLOTYPE *Mystacomyoidea mirabilis* Thomp. CNC No. 9813/ MARACAS V. Trinidad,BWI. FEB.16.1954/ Collector F.J.Simmonds/ 89/ HOLOTYPE CNCNo./ *Mystacomyoidea mirabilis* n.g. n.sp./ CNC Syntype *Mystacomyoidea mirabilis* Thompson Label affixed 1994.

Myxarchiclops caffer Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 495. SOUTH AFRICA.

Syntypes: 1♂, 1♀.

♂, ♀: ♀ ♂/ Cape Town G Peringuey Sep 1913/ *Myxarchiclops caffer* Typ. Villen./ *Myxarchiclops caffer* Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype *Myxarchiclops caffer* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Both syntypes are pinned to top label.

Myxarchiclops maculosus Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 496. SOUTH AFRICA.

Holotype ♀: Springbok Oct 90 R.M.L./ *Myxarchiclops maculosus* Typ. Villen./ *Myxogaedia maculosus* Vill. L.P. Mesnil det., 1969/ TYPE.

Myxarchiclops major Villeneuve, 1930a, Bull. Annls Soc. ent. Belg. 69 (1929): 353. SOUTH AFRICA.

Syntypes: 2♀♀.

♀: [blank pink label]/ IX 1925 Somerset Dr. Brauns. W.Strand Kapland/ *Myxarchiclops major* Villen. Typ./ *Myxarchiclops major* Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype *Myxarchiclops major* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
♀: [blank pink label]/ Somerset W. Strand Dr. Brauns. I.X 1925/ *Myxarchiclops major* Villen./ *Myxarchiclops major* Villen. L.P. Mesnil det., 1969/ CNC Syntype *Myxarchiclops major* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Second syntype is missing abdomen.

Myxexoristops sellersi Thompson, 1966, Studia ent. 8 (1965): 356. TRINIDAD.

Holotype ♂: HOLOTYPE *Myxexoristops sellersi* Thomp. CNC No. 9827/ CENTENO TRINIDAD BWI DEC. 1952/ Collector F.J.Simmonds/ 99/ FROM MANGO WOOD./ *Myxexoristops sellersi* nsp.

Myxexoristops (Myxexoristops) grandicornis Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 21. JAPAN.

Holotype ♂: TOKYO/ 169/ *Myxexoristops grandicornis* Mesn L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Myxexoristops (Myxexoristops) hertigi Mesnil, 1955b, Fliegen palaearkt. Reg. 64g: 450. GERMANY.

Holotype ♂: ex *Acantholyda erythrocephala* (Tenthru.) Haltern (Westf.-Albm.) 1951/ *Myxexoristops hertigi* Mesnil L.P. Mesnil det., 1967/ TYPE.

Nanoplagia nigrescens Mesnil, 1953b, Bull. Annls Soc. ent. Belg. 89: 172. "PALESTINE".

Holotype ♀: 13.II.40/ 92/ PALESTINE Dr. A. Shulov/ Dead Sea 13.2.1940/ Mesnilovoria nigrescens Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Nearchus duplaris Reinhard, 1964a, J. Kans. ent. Soc. 37: 46. USA.

Holotype ♂: WYO: Cody 8.VII.42 S. Camras/ HOLOTYPE *Nearchus duplaris* Reinhard/ *Nearchus duplaris* R Rnh.
Allotype ♀: Signal Mt., Wyo. Aug. 3, 1947/ F. A. Cowan M. R. Wheeler Collectors/ ALLOTYPE *Nearchus duplaris* Reinhard.
Paratype: 1♂, New Mexico.

Nearchus ningulus Reinhard, 1964a, J. Kans. ent. Soc. 37: 47. USA.

Holotype ♀: Beaver, Utah AUG.4.1954 G. F. Knowlton/ HOLOTYPE *Nearchus ningulus* Reinhard/ *Nearchus ningulus* R Rnh.

Nemoraea elegantula Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 59. JAPAN.

Holotype ♀: Sapporo 12-IX. 1942 Y.Nishijima/ 254/ *Dexiomima elegantula* Mesn. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Nemoraea infoederata Villeneuve, 1916a, Annls Soc. ent. Fr. 85: 199. ZAIRE.

Syntype: 1♂.
♂: R 2500 20.IV.14/ *Nemoraea infoederata* Typ. Villen./ TYPE/ *Dexiomima infoederata* Vill. L.P. Mesnil det., 1970/ Syntype CNC Syntype *Nemoraea infoederata* Villeneuve Label affixed 1994.

Nemoraea patellipalpis Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 58. JAPAN.

Holotype ♀: Mt. Hira Shiga JAPAN [underside of label] 12/VII'53 T.Hovio/ 252/ *Dexiomima patellipalpis* Mesn. L.P. Mesnil det., 1970/ TYPE/ *Dexiomima takanoi* Bar. L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970.

Nemoraea piligena Mesnil, 1971b, Fliegen palaearkt. Reg. 64g: 990. ALGERIA.

Holotype ♂: ex lepidopt/ Hussey Dey Alger [underside of label] 24-III-10 Hall/ *Nemoraea piligena* Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Nemosturmia japonica Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 9. JAPAN.

Holotype ♂: JAPAN T.Okutani [underside of label] 12/VI'49(L) 21/IV'50(A) Mitaka/ 328/ TYPE/ *Nemosturmia japonica* Mesnil L.P. Mesnil det., 1967/ EX L.-P. MESNIL COLLECTION 1970.

Nemosturmia laeta Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg.

39(24): 5. FRANCE.

Holotype ♂: La Celle St Cloud 21 Mai 1939/ Near Versailles France/ *Nemosturmia laeta* Mesnil L.P.

Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL
COLLECTON 1970.

Neocuphocera aurifacies Guimarães, 1963, Mems Inst. Oswaldo Cruz 61: 60. BRAZIL.
Syntypes: 2♂♂, 2♀♀.
♂, 2♀♀: São Paulo: Rio Paraná PORTO CABRAL IIIeIV-1944 L. Travassos col./ PARATYPUS/ *Neocuphocera aurifacies* J.H.Guimarães det. Guim./ CNC Syntype *Neocuphocera aurifacies* Guimarães Label affixed 1994.
♂: Paratypus/ BR-S. P. : R. Paraná PORTO CABRAL 1-25.IV-1944 Trav. Fo. & Carrera & E. Dente/ *Neocuphocera aurifacies* Guim.
J.H.Guimarães det./ CNC Syntype *Neocuphocera aurifacies* Guimarães Label affixed 1994.

Neominthoidea trinidadensis Thompson, 1968, Mem. ent. Soc. Can. 56: 30. TRINIDAD.
Holotype ♂: HOLOTYPE *Neominthoidea trinidadensis* Thomp. CNC No. 9845/ Brazil 8 6. 60 Trinidad,BWI.
Allotype ♀: ALLOTYPE *Neominthoidea trinidadensis* Thomp. CNC No. 9845/ S. Cruz V. II 14. 64 Trinidad,WI.
Paratypes: 1♂, 1♀, Trinidad.

Neotrafoia trinitatis Thompson, 1963a, Can. J. Zool. 41: 463.
TRINIDAD.
Syntypes: 1♂, 2♀♀.
♀: HOLOTYPE *Neotrafoia trinitatis* Thomp. CNC No. 9781/ Brazil II. 18. 61 Trinidad,BWI/ 1165 II.20.61/ *Neotrafoia incarum* TT./ CNC Syntype *Neotrafoia trinitatis* Thompson Label affixed 1994.
♂, ♀: Brazil II.26.61 Trinidad,BWI/ CNC Syntype *Neotrafoia trinitatis* Thompson Label affixed 1994.

Neoxynopoidea claripalpis Thompson, 1968, Mem. ent. Soc. Can. 56: 149. TRINIDAD.
Holotype ♂: HOLOTYPE *Neoxynopoidea claripalpis* Thomp. CNC No. 9876/ Maracas V. TRINIDAD W.I. X.21.60 1961 W.R.Thompson/ 1379 II.10.64.
Allotype ♀: ALLOTYPE *Neoxynopoidea claripalpis* Thomp. CNC No 9876/ CUREPE Trinidad, BWI. OCT.6.1952/ Collector F. D. Bennett/ 1010 W. V.13.59.
Paratype: 1♂, Trinidad.
Remarks: Holotype is missing terminalia.

Nicetria globosa Reinhard, 1953b, J. Kans. ent. Soc. 26: 100.
MEXICO.
Holotype ♀: HOLOTYPE *Nicetria globosa* Reinhard/ Vol. Sa Martin Las Tuxtla, Ver./ 29-V-51 L.Navarro/ Calodexia ? aldrichi R Cn./ HOMOTYPE *Euoestrogaster fuscus* Tnsd. Compared by Wood '79.

Nicetria peponis Reinhard, 1953b, J. Kans. ent. Soc. 26: 99.
MEXICO.
Holotype ♀: Vol. Sn Martin Las Tuxtla, Ver./ 29-V-51 L.Navarro/ HOLOTYPE *Nicetria peponis*

Reinhard/ Calodexia major R Cn/ Calodexia ? major Cn. Det.D.M.Wood 1979.
Paratype: 1♀, Mexico.

Nilea (Lylibaea) anatolica Mesnil, 1954a, Fliegen palaearkt. Reg. 64g: 362. TURKEY.
Holotype ♀: Akschehir Tal Anatolia1500m Lindner14.VI.34./ *Nilea anatolica* Mesn. L.P. Mesnil det., 19/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Nimiocauda erilis Reinhard, 1943c, Bull. Brooklyn ent. Soc. 38: 79. USA.
Holotype ♀: Wading River L.I., N.Y. VII-1924/ HOLOTYPE *Nimiocauda erilis* Reinhard/ *Nimiocauda erilis* R Rnh./ *Panacemyia erilis* R. Rnh.
Paratype: 1♀, New York.

Nimioglossa planicosta Reinhard, 1945, Can. Ent. 77: 36. USA.
Holotype ♂: Trout Creek, Utah 5-7- 1934 H.B. Stafford/ HOLOTYPE *Nimioglossa planicosta* Reinhard/ *Nimioglossa planicosta* Rnh. Reinhard Coll'n/ *Nimioglossa planicosta* R.'44 Rnh.
Paratype: 1♂, Utah.

Nimioglossa raviga Reinhard, 1945, Can. Ent. 77: 36. USA.
Holotype ♂: LINCOLN CO. N MEX. 21 VI 1929/ 9000 FT./ DAVID G. HALL COLLECTOR/ HOLOTYPE *Nimioglossa raviga* Reinhard/ *Nimioglossa raviga* Rnh. Reinhard Coll'n/ *Nimioglossa raviga* R.44 Rnh.
Paratype: 1♂, New Mexico.

Oberomyia palpalis Reinhard, 1964b, Acta zool. mex. 7: 2.
MEXICO.
Holotype ♂: Jalapa, Ver., Mex. VIII/1-6/61 R&K Dreisbach/ HOLOTYPE *Oberomyia palpalis* Reinhard/ *Oberomyia palpalis* R Rnh.
Paratype: 1♂, Mexico.

Ocyptera aberrans Villeneuve, 1936b, Bull. Mus. r. Hist. nat. Belg. 12(12): 2. ZAIRE.
Holotype ♀: Moho Lesse, 9.VI.14/ n.sp., 2 soies scut., 1 dc/ *Ocyptera aberrans* Type Villen.

Ocyptera ambulatoria Villeneuve, 1944, Bull. Soc. ent. Fr. 48 (1943): 144. TAIWAN.
Lectotype ♂: Formosa Sauter/ Takao 1907. XI.8./ LECTOTYPE ♂ *Ocyptera ambulatoria* n.sp./ Designated by R.W.Crosskey, 1974/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
Paratype: 1♂, Formosa.
Remarks: For lectotype designation see Crosskey (1976: 272).

Ocyptera cibrata Villeneuve, 1936b, Bull. Mus. r. Hist. nat. Belg. 12(12): 3. SOUTH AFRICA.
Syntype: 1♀.
♀: Algoa-Bay, Capland. Dr. H. Brauns 1 3 92/ 2 dc/ *Ocyptera cibrata* Typ. Villen./ TYPE/ CNC Syntype *Ocyptera cibrata* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.

Ocyptera deserta Villeneuve, 1936b, Bull. Mus. r. Hist. nat. Belg. 12(12): 2. NIGERIA.
Holotype ♂: N. Nigeria, Dr.J.W. Scott Macfie, 19.9.1910./ *Ocyptera deserta* Type Villen.

Ocyptera luciflua Villeneuve, 1944, Bull. Soc. ent. Fr. 48 (1943): 144. TAIWAN.
Lectotype ♂: Formosa, Sauter/ Kosempo, 1908. VI./ *Ocyptera luciflua* n.sp.. Lectotype ♂, designated by R.W. Crosskey 1974.
Paralectotypes: 1♂, 1♀, Taiwan.
Remarks: For lectotype designation see Crosskey (1976: 273).

Ocyptera retroflexa Villeneuve, 1944, Bull. Soc. ent. Fr. 48 (1943): 145. UGANDA.
Holotype ♀: Uganda Kampala 14. VII. 1936 H. Hargreaves/ Coll. 7777/ *Ocyptera retroflexa* Typ. Villen/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Ocyptera rufimana Villeneuve, 1944, Bull. Soc. ent. Fr. 48 (1943): 144. TAIWAN.
Lectotype ♂: Formosa Sauter/ Koroton 907.IX.1-15./ LECTO-TYPE/ *Ocyptera rufimana* n.sp./ Cylindromyia fuscipennis (Wied) det. R.W.Crosskey, 1974/ LECTOTYPE ♂ *Ocyptera rufimana* Villeneuve Designated by R.W.Crosskey, 1974/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
Paralectotype: 1♂, Taiwan.
Remarks: For lectotype designation see Crosskey (1976: 273).

Ocyptera scapularis Villeneuve, 1944, Bull. Soc. ent. Fr. 48 (1943): 145. ZIMBABWE.
Syntype: 1♂.
♂: Coll. 8699/ Vumba Mts., S. Rhodesia, 12.12.1936 Major Drysdale/ *Ocyptera scapularis* Villen. n.sp./ TYPE/ CNC Syntype *Ocyptera scapularis* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.

Oedematoocera optata Reinhard, 1942a, Ent. News 53: 107. USA.
Holotype ♂: New York VII-8-35/ HOLOTYPE Oedematoocera optata Reinhard/ Oedematoocera optata Rein. det.H.J.Reinhard/ Gilvelia optata Rn. Reinhard Coll'n./ Gilvelia optata R (Rnh.).
Allotype ♀: Cold Spr.Har. Long Island 8/20/1927 NY/ H C Hallock Collector/ ALLOTYPE Oedematoocera optata Reinhard/ Gilvelia optata Rn. Reinhard Coll'n.

Oestrocharis lutescens Villeneuve, 1927, Revue zool. afr. 15: 119. SOUTH AFRICA.
Holotype ♂: Willow-more Cape Dr. Brauns. I. 1926/ *Oestrocharis lutescens* Typ. Villen./ *Oestrocharis lutescens* Villen. L.P. Mesnil det., 1970/ Glaurocara flava Towns. L.P. Mesnil det., 1975/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Okanagania setifrons Brooks, 1943, Can. Ent. 75: 71. CANADA.
Holotype ♂: Osoyoos,B.C. 24.IV.1927 E.R.Buckell/ HoloTYPE ♂ *Okanagania setifrons* No. Brooks 5330.

Allotype ♀: Nicola B.C. June24.1925 H.L.Seamans/ AlloTYPE ♀ *Okanagania setifrons* No. Brooks 5330.
Paratypes: 1♂, Idaho; 1♀, Saskatchewan.

Onychogonia magna Brooks, 1944a, Can. Ent. 75 (1943): 227. USA.

Holotype ♂: Allen Canyon, Ut.Aug.21 1942/ G. F. Knowlton S. L. Wood Collectors/ Holotype ♂ *Onychogonia magna* No. Brks.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 3♂♂, 5♀♀, Utah; 4♀♀, Washington.

Opsoleskia major Thompson, 1968, Mem. ent. Soc. Can. 56: 89. TRINIDAD.

Holotype ♂: St. Augustine, Trinidad,BWI. July 20. 1935/ HOLOTYPE *Opsoleskia major* Thomp. CNC No. 9861/ On Hibiscus Ruth O'Connor Collector 317/ I.C.T.A. 4738/ *Opsoleskia major* n.sp./ HOMOTYPE *Telothyrionoma polybia* Cn. Compared by D.M.W. 76.

Opsotheresia nigricornis Reinhard, 1939, Bull. Brooklyn ent. Soc. 34: 67. USA.

Holotype ♂: Wisconsin VI-25-27/ HOLOTYPE *Opsotheresia nigricornis* Reinhard/ *Opsotheresia nigricornis* det.H.J.Rein Rein/ *Opsotheresia bigelowi* (Cn.) Reinhard Coll'n.
Allotype ♀: Iowa 7 VII 32/ ALLOTYPE *Opsotheresia nigricornis* Reinhard/ *Opsotheresia bigelowi* (Cn.) Reinhard Coll'n.
Remarks: Date of collection of holotype was published as 25 June 1937.

Orasturmia vallicola Reinhard, 1947, J. Kans. ent. Soc. 20: 22. USA.

Holotype ♂: Brownsville V-4-46 Tex./ HOLOTYPE *Orasturmia vallicola* Reinhard.
Paratypes: 3♂♂, Texas.

Orphanotrophus orbitalis Reinhard, 1943c, Bull. Brooklyn ent. Soc. 38: 83. USA.

Holotype ♂: Amherst, Ohio July, 1933/ H J Reinhard Collector/ HOLOTYPE *Orphanotrophus orbitalis* Reinhard/ Euhalidaya genalis R (Coq.)/ *Orphanotrophus orbitalis* Rein. det.H.J.Reinhard.
Allotype ♀: Same data as holotype except with allotype label.
Paratype: 1♀, Ohio.

Oswaldia bicoloripes Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 23. JAPAN.

Holotype ♂: Obihiro S. Takano [underside of label] 26/VI'50/ TYPE/ 337/ *Drinomyia bicoloripes* Mesnil L.P. Mesnil det., 1969/ EX L.-P. MESNIL COLLECTION 1970.

Oswaldia hirsuta Mesnil, 1970, Mushi 44: 115. JAPAN.

Holotype ♀: Nukabira Hokkaido 1.VIII.1961 S.Takano/ 382/ *Oswaldia hirsuta* Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Oswaldia micronychia Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 22. JAPAN.

Holotype ♂: Obihiro S.Takano [underside of label] 13.VII.51/ 234/ *Oswaldia micronychia* Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Pachystylum albescens Pandellé, 1896, Revue Ent. 15: 86. FRANCE.

Syntype: 1♂.
♂: ♂ 1226/ Tarbes./ *Pachystylum albescens* types Pand./ *Chaetomera fumipennis* B.B. L.P. Mesnil det., 1969/ CNC Syntype *Pachystylum albescens* Pandellé Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Pacidianus hirsutus Reinhard, 1943c, Bull. Brooklyn ent. Soc. 38: 89. USA.

Holotype ♂: AG. COLL. MICH. 21 Apr. 1922/ L.G. Gentner Collection/ HOLOTYPE *Pacidianus hirsutus* Reinhard.

Pacidianus persimilis Reinhard, 1943c, Bull. Brooklyn ent. Soc. 38: 90. USA.

Holotype ♂: Babylon,L.I.,N.Y. IX-4 1936 Blanton & Borders/ HOLOTYPE *Pacidianus persimilis* Reinhard/ *Pacidianus persimilis* R'43 Rnh. Paratype: 1♂, Ontario.

Pales contristans Villeneuve, 1938c, Bull. Mus. r. Hist. nat. Belg. 14(38): 1. SOUTH AFRICA.

Syntype: 1♂.
♂: Algoa-Bay. Capland. Dr. H. Brauns 10 12 16/ *Pales contristans* Typ. Villen./ *Pales contristans* Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype *Pales contristans* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Pales murina Mesnil, 1970, Mushi 44: 90. PAKISTAN.

Holotype ♂: Ex.hairy caterpillar on Q. dilatata/ Pakistan Ghaviai 20.6.1967/ *Pales murina* Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Palesisa nudioculata Villeneuve, 1929b, Bull. Annls Soc. ent. Belg. 69: 101. TURKMENIA.

Holotype ♂: Transasp Imam-baba/ *Palesisa nudioculata* Villen./ *Palesisa nudioculata* Vill. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Pallasia amplifrons Brooks, 1946, Can. Ent. 77 (1945): 225. CANADA.

Holotype ♀: Morrin,Alta, 27-VI -1939 P.J.G.Rock/ 96/ HoloTYPE ♀ *Pallasia amplifrons* Brks No. 5344/ SLIDE O77 No.
Remarks: Holotype terminalia are mounted on a microscope slide.

Palpexorista meridionalis Thompson, 1968, Mem. ent. Soc. Can. 56: 21. TRINIDAD.

Holotype ♂: Mt. ST. BENIDICT, [Benedict] Trinidad, BWI JULY.11.1953./ COLLECTOR W.R.THOMPSON/ HOLOTYPE *Palpexorista meridionalis* Thomp. CNC No. 9843/ *Palpexorista*

meridionalis nsp/ 78/ Sp. K.A. Phorocera s.lat./ 10159m♂ Sp. KA.

Remarks: Holotype is missing terminalia, and legs are glued to bottom label.

Palpexorista minor Thompson, 1968, Mem. ent. Soc. Can. 56: 24. TRINIDAD.

Holotype ♀: HOLOTYPE *Palpexorista minor* Thomp. CNC No. 9844/ Santa Cruz V. 8. 2. 60 Trinidad,BWL/ [blue label].
Paratype: 1♀, Trimad.

Panacemyia pallipes Reinhard, 1953d, Proc. ent. Soc. Wash. 55: 246. USA.

Holotype ♀: College Station May 16,1946 Tex/ H J Reinhard Collector/ HOLOTYPE *Panacemyia pallipes* Reinhard/ *Panacemyia pallipes* R. Rnh.
Paratype: 1♀, Texas.

Panacemyia trinitatis Thompson, 1963a, Can. J. Zool. 41: 459. TRINIDAD.

Lectotype ♀: I.C.T A ST AUGUSTINE TRINIDAD AUG 1959/ Ex coll. OF SAMAAN FLOWERS/ F.D.Bennett Collector/ LECTOTYPE ♀ of *Panacemyia trinitatis* Thompson designated 1984 D.M.Wood.

Parlectotypes: 2♀ ♀, Trinidad.

Remarks: For lectotype designation see Wood (1985: 103). Lectotype is missing terminalia and most abdominal tergites.

Panacemyia verticalis Reinhard, 1953d, Proc. ent. Soc. Wash. 55: 247. USA.

Holotype ♂: Amherst, Ohio VI-15 1924/ H.J.Reinhard Collector/ HOLOTYPE *Panacemyia verticalis* Reinhard/ *Panacemyia verticalis* R Rnh./ IDEMA illustration.

Pandelleia otiorynchi Villeneuve, 1922c, Annls Sci. nat. Zool. (10) 5: 338. GERMANY.

Lectotype ♂: Traben Mosel Rheinl./ auf Otiorynchus sulcatus/ *Pandelleia otiorynchi* Villen. L.P. Mesnil det., 1975/ TYPE/ CNC Syntype *Pandelleia otiorynchi* Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1975: 1363. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.

Parlectotype: 1♀, Germany.

Remarks: We regard Mesnil's (1975b: 1363) reference to a holotype as a lectotype fixation for this species.

Pandelleia pschorni Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 52. JAPAN.

Holotype ♀: Nagano C. Honshu, VI-8 K. KAMIJO 1959/ *Afrophasia pschorni* Mesnil L.P. Mesnil det., 1975/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Pantagathus curulis Reinhard, 1943b, J. Kans. ent. Soc. 16: 18. USA.

Holotype ♀: Baton Rouge 8-25-35 La./ HOLOTYPE *Pantagathus curulis* Reinhard/ *Pantagathus curulis* det. H.J.Reinhard Rein/ Pelecotheca curulis R Rnh.

Paradidyma merista Reinhard, 1953b, J. Kans. ent. Soc. 26: 95.
MEXICO.

Holotype ♀: AMANALCO MÉX. 5-V-50 Coll.
WGDowns/ HOLOTYPE *Paradidyma merista*
Reinhard/ *Paradidyma merista* R Rnh.
Paratype: 1♀, Mexico.

Paradidyma recincta Reinhard, 1964b, Acta zool. mex. 7: 19.
MEXICO.

Holotype ♂: Cuernavaca, Mor., Mex. 7-29-61 7000'
R&K Dreisbach/ HOLOTYPE *Paradidyma*
recincta Reinhard/ *Paradidyma recincta* R Rnh.

Paradmontia picticornis Reinhard, 1967, J. Kans. ent. Soc. 40: 96.
USA.

Holotype ♀: Sta.Catalina Mts.ARIZ. Mi.12 July 21,
1958/ HOLOTYPE *Paradmontia picticornis*
Reinhard/ Reinhard Collection.

Paradrino solitaris Thompson, 1966, Studia ent. 8 (1965): 420.
TRINIDAD.

Holotype ♂: HOLOTYPE *Paradrino solitaris* Thomp.
CNC No. 9889/ Maracas V. TRINIDAD W.I.
XI.28 1960 W.R.Thompson/ *Paradrino solitaris*
n.sp.

Remarks: Date of collection of holotype was published
as 28 November 1961.

Paragonia portentosa Mesnil, 1950b, Fliegen palaearkt. Reg. 64g:
106. AUSTRALIA.

Holotype ♂: AUSTRALIE, OCCIDENT. WAROONA
G.F.BERTHOUD 1910/ Waroona, W.A. 1. 11. 09
G. F. Berthoud/ *Paragonia portentosa* Mesnil L.P.
Mesnil det., 1967/ TYPE/ HOLOTYPE ♀ of
Paragonia portentosa Mesnil Examined 1970
R.W.Crosskey.

Remarks: Holotype terminalia are glued to locality
label.

Paralispe aldrichi Curran, 1926c, Can. Ent. 58: 217. CANADA.
Holotype ♂: *Paralispe aldrichi* ♂ Curran No.
1965/ Aylmer,Que. 8.VIII.1924 C.H.Curran/ ♂.

Paranoxynops curepeiensis Thompson, 1968, Mem. ent. Soc. Can.
56: 64. TRINIDAD.

Holotype ♂: CUREPE TRINIDAD BWI DEC.29.1952/
Collector F.J.Simmonds/ 47/ HOLOTYPE
Paranoxynops curepeiensis Thomp. CNC No.
9854.

Paratype: 1♂, Trinidad.

Paraphasia albipennis Brooks, 1945a, Scient. Agric. 25: 657.
CANADA.

Holotype ♂: Saskatoon,Sask. 2-V-1940 A. R. Brooks/
HoloTYPE ♂ *Paraphasia albipennis* No. Brks.
5588/ *Phasia fenestrata* (Bigot 1889) ♂ Det.
Xuekui Sun, 1993.

Paraphasiopsis trinitatis Thompson, 1963d, Can. Ent. 95: 1311.
TRINIDAD.

Holotype ♂: HOLOTYPE *Paraphasiopsis trinitatis*
Thomp. CNC No. 9810/ PIARCO Trinidad,BWI.
OCT.28.1953./ Collector F.J.Simmonds/
HOLOTYPE CNCNo./ *Paraphasiop-sis trinitatis*
n.sp.

Paratypes: 4♂♂, Trinidad.

Paraphasmophaga dissita Reinhard, 1962b, Pan-Pacif. Ent. 38:
221. USA.

Holotype ♀: Yuma, ARIZONA June 25, 1955/
HOLOTYPE *Paraphasmophaga dissita* Reinhard/
Paraphasmophaga dissita R Rnh.

Paratypes: 1♀, Arizona; 2♀♀, California.

Paraphasmophaga pictipennis Reinhard, 1935c, Ann. ent. Soc.
Am. 28: 167. USA.

Holotype ♀: Castolon 11-8-29 Tex/ E.R.Tinkham
Collector/ Castolon Tex Nov 8.1929/ Holotype P.
pictipennis Reinhard/ HOLOTYPE
Oraphasmophaga pictipennis Reinhard/
Paraphasmophaga pictipennis det. H.J.Reinhard
Rein/ *Oraphasmophaga pictipennis* R Rnh.

Paraphorantha auricaudata Brooks, 1945a, Scient. Agric. 25: 661.
USA.

Holotype ♂: Milton, Oregon June 22, 1938 Coll. K.
Gray J. Schuh/ HoloTYPE ♂ *Paraphorantha*
auricaudata No. Brks/ Reinhard Collection/
Paraphorantha auricaudata Det. Brks A.R.Brooks/
Phasia pollinosa (Brooks 1945) ♂ Det. Xuekui
Sun, 1993.

Allotype ♀: Cuchara,Colo. 6-10000 Ft.Elev. F.Snyder
8-7 '40/ Allo TYPE ♀ *Paraphorantha auricaudata*
No. Brks/ Reinhard Collection/ *Phasia pollinosa*
(Brooks 1945) ♀ Det. Xuekui Sun, 1993.

Paraprosena marmorata var. *africana* Villeneuve, 1935b, Revue
Zool. Bot. afr. 27: 138. NIGERIA.

Syntype: 1♂.
♂: Capt. Blair Ilesha, S.Nigeria. 2/11 Billaea
(*Phorostoma Rond*) *africana* type Villen./ TYPE/
CNC Syntype *Paraprosena marmorata* var.
africana Villeneuve Label affixed 1994/ EX L.-P.
MESNIL COLLECTION 1985.

Pararondania multipunctata Villeneuve, 1916c, Ann. S. Afr. Mus.
15: 498. SOUTH AFRICA.

Holotype ♀: S. W. Distr. Cape Col./ *Pararondania*
multipunctata Typ. ♀ Villen./ TYPE/ *Pararondania*
(=Steleoneura Stein) *multipunctata* Vill. L.P.
Mesnil det., 1975/ EX L.-P. MESNIL
COLLECTION 1970.

Paratamiclea pallida Villeneuve, 1936c, Bull. Mus. r. Hist. nat.
Belg. 12(41): 1. ZIMBABWE.

Syntype: 1♂.
♂: Salisbury Rhodesia Dept. Agric. 30.XI.1935/
Paratamiclea pallida Typ. Villen./ *Eutrixopsis*
pallida Vill L.P. Mesnil det., 1970/ TYPE/ CNC
Syntype *Paratamiclea pallida* Villeneuve Label
affixed 1994/ EX L.-P. MESNIL COLLECTION
1970.

Paratrixa takanoi Mesnil, 1970, Mushi 44: 115. JAPAN.

Holotype ♀: Nukabira Hokkaido 21.VIII.1964
S.Takano/ 372/ *Paratrixa takanoi* Mesnil L.P.
Mesnil det., 1969/ TYPE.

Paratryphera longicornis Mesnil, 1970, Mushi 44: 117. INDIA.

Holotype ♂: Inside tunnel of Toon twig made by Hypsipyla robusta/ PASHOK W. BENGAL 9.VI.1965/ CIBC -IS KPG.SS./ 3/ Paratryphera longicornis Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Holotype puparium is pinned below specimen.

Paratryphera mesnili Herting, 1977, Stuttg. Beitr. Naturk. (A) 295: 2. ALBANIA.

Holotype ♂: Alban.Exp.1918 Bicaj 14.,15.VI./ Albany Nr. Yugoslavia/ Paratryphera handlirschi B.B. L.P. Mesnil det., 1969/ Paratryphera mesnili n.sp. B. Herting det. Type/ EX L.-P. MESNIL COLLECTION 1970.

Parepalpus labeosus Reinhard, 1957, Ent. News 68: 102. MEXICO.

Holotype ♂: Cuernavaca,Mex. March 22 1934 S. E. Jones.Coll./ HOLOTYPE Parepalpus labeosus Reinhard/ Parepalpus labeosus R Rnh./ not discors comp. to syntype Det. D.M.Wood 1989.

Paratype: 1♀, Mexico.

Parerigone takanoi Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 62. JAPAN.

Holotype ♂: TYPE/ Obihiro, S.Takano, [underside of label] 15/VI '50/ 278./ Parerigone takanoi Mesnil, L.P. Mesnil det. 1967.

Patelloa nigripalpis Thompson, 1963b, Studia ent. 6: 368. TRINIDAD.

Holotype ♂: TRINIDAD W.I. W.R.Thompson/ Patelloa nigripalpis n.sp/ HOLOTYPE Patelloa nigripalpis Thomp. CNC No.9794.

Pelatachina orillia Curran, 1927c, Can. Ent. 59: 23. CANADA.

Holotype ♀: Pelatachina TYPE orillia ♀ Curran No. 2238/ Orillia,Ont. 5-VI-1925 C.H.Curran.

Pelecotheca flavipes Thompson, 1968, Mem. ent. Soc. Can. 56: 46. TRINIDAD.

Holotype ♀: HOLOTYPE Pelecotheca flavipes Thomp. CNC No. 9850/ MARACAS V. Trinidad,BWI. NOV.12-1952./ Collector F.J.Simmonds/ 1015 V.29-59/ Pelecotheca sp Det.Th.

Paratypes: 3♀ ♀, Trinidad.

Pelecotheca trinidadensis Thompson, 1968, Mem. ent. Soc. Can. 56: 44. TRINIDAD.

Holotype ♀: HOLOTYPE Pelecotheca trinidadensis Thomp. CNC No. 9849/ FLORADELLE Trinidad,BWI. FEB.12.1954./ Collector F.J.Simmonds/ Pelecotheca nsp. trinidadensis [underside of label] Pelecotheca Det. n.sp.

Peleteria aclista Reinhard, 1956b, Pan-Pacif. Ent. 32: 109. MEXICO.

Holotype ♂: Mt. Popo.3000m. W.G. Downs/ XI-27-51/ HOLOTYPE Peleteria aclista Reinhard/ Peleteria aclista Rnh det.H.J.Reinhard.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 2♂♂, Mexico.

Remarks: Holotype terminalia are glued to locality label.

Peleteria alberta Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 234. CANADA.

Holotype ♂: HOLOTYPE Peleteria alberta No. Curr. 1231/ Banff,Alta. 2 VIII 1922 C.B.D.Garrett 7600 ft.

Peleteria angulata Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 236. CANADA.

Holotype ♂: HOLOTYPE Peleteria angulata No. Cur. 1233/ Hedley,B.C. 25.VII 1923 C.B.Garrett. Paratypes: 1♂, Alberta; 1♂, British Columbia; 1♀, Washington.

Peleteria arctica Malloch, 1919, Rep. Can. arct. Exped. 3: 57c. CANADA.

Holotype ♀: Cockburn Pt Can. Arctic Coast/ TYPE No. 492/ Canadian Arctic Expedition Sep.5 1914/ F. Johansen Coll.-/ 544/ Peleteria aenea (Staeger) det. L. Richards 1969.

Peleteria bryanti Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 250. CANADA.

Holotype ♂: HOLOTYPE Peleteria bryanti Curran No. 1242/ Slave Lake/ AUG 25'24.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 6♂♂, 2♀ ♀, Alberta; 1♂, Saskatchewan; 2♂♂, Colorado; 1♂, New Mexico.

Remarks: There is one additional specimen labeled as a paratype but it is from a locality not mentioned in the original description.

Peleteria campestre Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 247. CANADA.

Holotype ♂: P.N.Vroom Aweme, Man. 5.VIII. 1920/ HOLOTYPE Peleteria campestris Curr No. 1243.

Allotype ♀: Same data as holotype except with allotype label and collected "12.VIII.1920".

Paratypes: 6♂♂, 3♀ ♀, Manitoba; 6♂♂, 5♀ ♀, Alberta; 4♂♂, British Columbia; 1♂, Oregon; 2♂♂, 1♀, Wyoming; 10♂♂, 2♀ ♀, Arizona; 3♂♂, 1♀, California; 5♂♂, 3♀ ♀, New Mexico; 2♂♂, 1♀, Mexico.

Remarks: There are seven additional specimens labeled as paratypes but they are from localities not mentioned in the original description.

Peleteria carnata Reinhard, 1953a, Bull. Brooklyn ent. Soc. 48: 92. MEXICO.

Holotype ♀: Zamora, Mich. Aug. 27, 1947/ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE Peleteria carnata Reinhard.

Paratype: 1♀, Mexico.

Peleteria clara Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 239. CANADA.

Holotype ♂: HOLOTYPE Peleteria clara No. Curr. 1234/ Lethbridge Alta Aug 9-1921 E.R.Strickland.

Allotype ♀: ALLOTYPE P. clara Curran No. 1234/ Lethbridge,Alta. Aug 18 1922 H.E.Gray.

Paratypes: 8♂♂, 12♀ ♀, Alberta; 2♂♂, Colorado.

- Peleteria confusa* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 253. CANADA.
 Holotype ♂: Holo.TYPE *Peleteria confusa* No. Curr 1241/ Jordan, Ont 18.VII, 1919 C.H.Curran.
 Allotype ♀: Allo TYPE P. *confusa* No. Curr 1241./ P.N.Vroom Aweme, Man. 12.VIII. 1920.
 Paratypes: 3♂♂, New Brunswick; 3♂♂, 1♀, Quebec; 1♂, 3♀♀, Ontario; 3♂♂, 4♀♀, Manitoba; 2♂♂, 1♀, Alberta; 1♀, Massachusetts.
 Remarks: There are ten additional specimens labeled as paratypes but they are from localities not mentioned in the original description.
- Peleteria conjuncta* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 235. CANADA.
 Holotype ♂: HoloTYPE *Peleteria conjuncta* No. Curran/ Nicola, B.C. 17.V. 1923 P.N. Vroom/
Peleteria conjuncta Cn. Reinhard Coll'n.
 Paratypes: 2♂♂, 2♀♀, Colorado.
- Peleteria cornigera* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 232. CANADA.
 Holotype ♂: TYPE *Peleteria cornigera* No. Curr. 1228/ Lillooet, B.C. A.W. Phair.
 Paratypes: 1♂, 1♀, Colorado.
- Peleteria cornuta* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 232. CANADA.
 Holotype ♂: HoloTYPE *Peleteria cornuta* Curr. No. 1229/ Pincher, Alta. July 18 1913 E. H. Strickland.
 Allotype ♀: AlloTYPE P. *cornuta* Curr. No. 1229./ Calgary 10 Jul. 03.
 Paratypes: 1♂, 1♀, Alberta.
 Remarks: Holotype right wing is glued to the holotype label, and left wing is pinned below locality label.
- Peleteria cornuticaudata* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 234. CANADA.
 Holotype ♂: Holo TYPE *Peleteria cornuticaudus* [sic] Curr No. 1232/ Banff, Alta. 2 VIII 1922 C.B.D.Garrett 7600 ft.
 Allotype ♀: Banff, Alta. 20.IX 1922 C.B.D.Garrett/ Allo.TYPE *Peleteria cornuticaudatus* No. Curr. 1232.
- Peleteria eronis* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 238. CANADA.
 Holotype ♂: Holo.TYPE *Peleteria eronis* No. Curr. 1235/ William's Lake, B.C. 24.viii. 1919 W.B.Anderson.
 Allotype ♀: AlloTYPE *Peleteria eronis* No. Curr. 1235./ Lardo B.C. 26.IX 1919 C. G. Hewitt.
 Paratypes: 2♂♂, Alberta; 1♂, Manitoba; 10♂♂, 1♀, Colorado; 5♂♂, 2♀♀, Arizona; 2♂♂, 1♀, New Mexico.
 Remarks: There are five additional specimens labeled as paratypes but they are from localities not mentioned in the original description.
- Peleteria mimica* Villeneuve, 1913, Revue zool. afr. 3: 26. ZAIRE.
 Holotype ♂: Sankisia 25.VIII-1911/ *Peleteria mimica* Villen. Dr Villeneuve det. Typ./ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

- Peleteria obsoleta* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 251. USA.
 Holotype ♂: TYPE *Peleteria obsoleta* Curran. No. 1498/ Magdalena Mts N.M. Aug. '94 Snow.
 Paratype: 1♂, Costa Rica.
- Peleteria phairi* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 233. CANADA.
 Holotype ♂: HoloTYPE *Peleteria phairi* Curr No. 1230/ Lillooet, B.C. A.W.A.Phair 29.7.18 8000ft.
 Paratype: 1♂, Alberta.
- Peleteria texensis* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 246. USA.
 Holotype ♂: HoloTYPE *Peleteria texensis* Curr No. 1239/ College Station 5-21-19Tex/ H.J.Reinhard Collector.
 Allotype ♀: Allo TYPE *Peleteria texensis* Curr. No. 1239/ College Station 5-19-19Tex/ H.J.Reinhard Collector.
 Paratypes: 2♂♂, 3♀♀, Texas; 1♀, Mexico.
- Peleteria torta* Reinhard, 1943b, J. Kans. ent. Soc. 16: 21. USA.
 Holotype ♂: Roosevelt Lake, Ariz. 19/ Mar DKDuncan/ HOLOTYPE *Peleteria torta* Reinhard/ *Peleteria torta* det.H.J.Reinhard Rein/ *Peleteria torta* Rnh. Reinhard Coll'n.
 Remarks: Holotype terminalia are glued to locality label.
- Peleteria valida* Curran, 1925c, Trans. R. Soc. Can., Ser. 3, 19: 255. USA.
 Holotype ♂: S.C. 31.VII-13/ TYPE *Peleteria valida* Curr No. 1240.
 Remarks: Holotype terminalia are glued to locality label.
- Peleteria (Sphyromyia) neotexensis* Brooks, 1949, Can. Ent. 81: 23. USA.
 Holotype ♂: College Station Apr.21,1944 Tex/ H. J. Reinhard Collector/ Holo TYPE ♂ *Peleteria neotexensis* Brks No. 5773.
 Allotype ♀: Same data as holotype except with allotype label and collected "May 9, 1944".
 Paratypes: 3♂♂, 2♀♀, Texas.
- Peleteria multispinosa* Thompson, 1963a, Can. J. Zool. 41: 413. TRINIDAD.
 Holotype ♀: HOLOTYPE *Peleteria multispinosa* Thomp. CNC No.9776/ Santa Cruz V 3.5.60 Trinidad, BWI.
- Peleteria trinitatis* Thompson, 1963a, Can. J. Zool. 41: 415. TRINIDAD.
 Syntype: 1♀.
 ♀: HOLOTYPE *Peleteria trinitatis* Thomp. CNC No. 9777/ Hollis Res. II.25.61. Trinidad, BWI/ Type ♀/ CNC Syntype *Peleteria trinitatis* Thompson Label affixed 1994.
 Remarks: Date of collection of syntype was published as 25 March 1961.
- Periprosena dispar* Villeneuve, 1938c, Bull. Mus. r. Hist. nat. Belg. 14(38): 14. ZAIRE.

- Holotype ♀: Mokoto XII-14/ Periprosena dispar Typ. Villen./ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Periscepsia pyrenaica* Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 50. FRANCE.
Holotype ♂: URDOS Sompor 18.VII.19 B. [Basses] P. [Pyrénées]/ Periscepsia pyrenaica Mesnil L.P. Mesnil det., 1970/ TYPE/ Ramona plorans Rond. L.P. Mesnil det., 1975/ EX L.-P. MESNIL COLLECTION 1985.
- Peteina ontario* Curran, 1926d, Can. Ent. 58: 87. CANADA.
Holotype ♂: Peleina TYPE ♂ ontario Curran No. 1870/ N.K.Bigelow Macdiarmid Lake Nipigon,Ont. 4.VII.1923.
Allotype ♀: Same data as holotype except without ♂ symbol.
- Petinops labradorensis* Brooks, 1945b, Can. Ent. 77: 92. CANADA.
Holotype ♂: HoloTYPE ♂ Petinops labradorensis No. Brks 5534/ Hopedale,Labr. 5. VI 1928 W.W.Perrett.
Allotype ♀: Same data as holotype except with allotype label and collected "11.VI 1928".
Paratype: 1♂, Newfoundland.
Remarks: Date of collection of holotype was published as 3 June 1928.
- Pexomyia (Erythrocera) picta* Villeneuve, 1936a, Bull. Mus. r. Hist. nat. Belg. 12(4): 7. NIGERIA.
Holotype ♂: Oshogbo, S.Nigeria. Dr.T.F.G.Mayer Nov1910/ Erythrocera (Pexomyia) picta Typ. Villen./ TYPE/ HOLOTYPE ♂ of Pexomyia (Erythrocera) picta Vill. Examined 1970 R.W.Crosskey.
- Pexopsis pilosa* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 14. JAPAN.
Holotype ♂: IV-22.1949 Hakusan LHATTORI/ 273/ Pexopsis pilosa Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Pexopsis rasa* Mesnil, 1970, Mushi 44: 107. PHILIPPINES.
Holotype ♀: Luzon. Banahao IV. 1914/ Pexopsis rasa Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: The type locality was published as Lucon.
- Peyritschia roederi* Villeneuve, 1932a, Bull. Annls Soc. ent. Belg. 71 (1931): 241. GREECE.
Holotype ♀: Attica/ Röder/ Peyritschia Roderi Villen Type/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
- Phaenopsis venusta* Reinhard, 1946b, Can. Ent. 78: 120. USA.
Holotype ♂: College Station 5-18.23 Texas/ H J Reinhard Collector/ HOLOTYPE Phaenopsis venusta Reinhard/ Phaenopsis venusta R. Rnh.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 2♂♂, 3♀♀, Texas.
- Phasiopsis triangularis* Thompson, 1963d, Can. Ent. 95: 1306. TRINIDAD.
Holotype ♀: HOLOTYPE Phasiopsis triangularis Thomp. CNC No. 9808/ FLORADALE Trinidad,BWI. FEB.12.1954/ Collector F.J.Simmonds/ 1278 XII.1962/ HOLOTYPE CNCNo./ Phasiopsoidae triangularis n.sp.
- Phasiopsis trinitatis* Thompson, 1963d, Can. Ent. 95: 1304. TRINIDAD.
Holotype ♂: HOLOTYPE Phasiopsis trinitatis Thomp. CNC No. 9807/ I.C.T.A. Trinidad,BWI. FEB.18.1953/ Collector F.J.Simmonds/ HOLOTYPE CNCNo./ Phasiopsis trinitatis n.sp.
Allotype ♀: Same data as holotype except with allotype label and dated "MARCH 1954".
Paratypes: 4♂♂, 4♀♀, Trinidad.
- Phebellia monochaeta* Mesnil, 1970, Mushi 44: 111. RUSSIA.
Holotype ♂: [locality in Russian] 15.VIII.957/ Phebellia monochaeta Mesn L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Phebellia nigricauda* Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 14. JAPAN.
Holotype ♀: Kagoshima Kyushu, 13 - V K. KAMIO 1959/ Phebellia nigricauda n.sp. L.P. Mesnil det., 1962/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Philocalia tenuirostris* Reinhard, 1939, Bull. Brooklyn ent. Soc. 34: 71. USA.
Holotype ♀: Bozeman VII.18.36 Mont./ 37/ HBHoeffler coll./ HOLOTYPE Philocalia tenuirostris Reinhard/ Philocalia tenuirostris det.H.J.Reinhard Rein.
- Philocorus montanum* Cortés, 1975, Boln téc Estac. exp. agron. Fac. Agronomía Univ. Chile 40: 13. CHILE.
Holotype ♂: ♂/ HOLOTIPO/ Banos El Toro Coquimbo,CHILE 3300-4000 m. 6-12.I.66.Pena/ Philocorus ♂ montanum n.sp. R.Cortés det. 1974.
Paratypes: 2♂♂, Chile.
- Phobetromyia dumalis* Reinhard, 1964b, Acta zool. mex. 7: 10. MEXICO.
Holotype ♀: Rio Frio, Mex. Sept. 9, 1947/ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE Phobetromyia dumalis Reinhard.
Paratype: 1♀, Mexico.
- Phorinia cinctella* Mesnil, 1971a, Entomophaga 16: 70. UGANDA.
Holotype ♂: UGANDA KAWANDA 21. V -1969 A.I.MOHYUDDIN/ Ex Lepid. larvae On Loranthus Loranthus-49/ Phorinia cinctella Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
Remarks: Holotype puparium is stored in a gelatine capsule pinned below specimen.
- Phorocera anassa* Reinhard, 1959b, Pan-Pacif. Ent. 35: 160. USA.

Holotype ♂: L.Waha Ida VIII-8-41/ HOLOTYPE
Phorocera anassa Reinhard/ *Cyzenis incrassata*
 (Smith) Det. J. O'Hara 1992.
 Allotype ♀: Same data as holotype except with allotype
 label.

Phorocera arnaudi Reinhard, 1956b, Pan-Pacif. Ent. 32: 106. USA.
 Holotype ♂: Chino Canyon Palm Springs, Cal.
 December 22, 1950 P. H. Arnaud/ On foliage of
Hyptis emeryi ♂/ HOLOTYPE *Phorocera arnaudi*
 Reinhard.
 Paratypes: 2♂♂, California; 1♂, Nebraska.
 Remarks: There are nine additional specimens labeled
 as paratypes but they are from localities not
 mentioned in the original description.

Phorocera clunalis Reinhard, 1956b, Pan-Pacif. Ent. 32: 107. USA.
 Holotype ♂: Mission, Tex. June 29, 1947/ H J Reinhard
 Collector/ HOLOTYPE *Phorocera clunalis*
 Reinhard/ P. *clunalis* Rnh.

Phorocera crassipalpis Villeneuve, 1938c, Bull. Mus. r. Hist. nat.
 Belg. 14(38): 2. ZAIRE.
 Syntype: 1♀.
 ♀: Congo-belge Bomputu-XII-1935 J. Ghesquière
 1075/ Parasite Lepido 1048/ R. Mus. Hist. Nat.
 Belg. 10482/ *Phorocera crassipalpis* typ. ♀ Villen./
P. crassipalpis Villen. L.P. Mesnil det., 1969/
 CNC Syntype *Phorocera crassipalpis* Villeneuve
 Label affixed 1994/ EX L.-P. MESNIL
 COLLECTION 1970.

Phorocera longirostris Villeneuve, 1938c, Bull. Mus. r. Hist. nat.
 Belg. 14(38): 2. SOUTH AFRICA.
 Holotype ♂: Marley nov.15 1824 Cranz K./
Chlorolydella longirostris Villen. L.P. Mesnil det.,
 1969/ TYPE/ EX L.-P. MESNIL COLLECTION
 1970.

Phorocera noera Reinhard, 1957, Ent. News 68: 109. MEXICO.
 Holotype ♂: Cuernavaca, VIII-6-42 Mex./ HOLOTYPE
Phorocera noera Reinhard/ *Phorocera noera* Rnh./
 =linearis comp. to Ht♂ Det. D.M.Wood 1982.

Phorocera pellecta Reinhard, 1957, Ent. News 68: 106. USA.
 Holotype ♂: Brown'sCn. Baboquivari Mts. ARIZ Aug
 18 1955 FGWerner & GDButler/ HOLOTYPE
Phorocera pellecta Reinhard/ Palpexorista pellecta
 R (Rnh)/ *Phorocera pellecta* R Rnh.
 Allotype ♀: Same data as holotype except with allotype
 label.
 Paratypes: 5♀♀, Arizona.
 Remarks: Date of collection of holotype and allotype
 was published as 19 August 1955.

Phorocera stolida Reinhard, 1957, Ent. News 68: 107. USA.
 Holotype ♂: Amherst, Ohio VIII-28 1930/ H J Reinhard
 Collector/ HOLOTYPE *Phorocera stolida*
 Reinhard.
 Paratypes: 2♂♂, Ohio.

Phorocera (Parasetigena) agilis takaoi Mesnil, 1960b, Fliegen
 palaearkt. Reg. 64g: 637. JAPAN.
 Holotype ♂: IV. 4. 1931 Mt. Takao T. IIIDA/ 187/
Parasetigena takaoi Mesnil L.P. Mesnil det., 1970/
 TYPE.
 Remarks: The collector's name was published as T.
 Lida.

Phorocera (Pseudotachinomyia) convexa Wood, 1972, Can. Ent.
 104: 484. CANADA.
 Holotype ♂: Old Chelsea, QUE. Summit King Mtn.
 22.V.1968 D.M.Wood/ HOLOTYPE *Phorocera*
(Pseudotachinomyia) convexa Wood CNC No.
 10931.
 Allotype ♀: Summit King Mt. Old Chelsea, QUE. 1150'
 17.VI.70 J.R.Vockeroth/ ALLOTYPE P. *convexa*
 CNCNo. Wood.
 Paratypes: 5♂♂, Quebec; 1♂, Ontario; 1♂, 2♀♀, Texas;
 1♂, New York.

Phorocera (Pseudotachinomyia) exigua Wood, 1972, Can. Ent.
 104: 484. CANADA.
 Holotype ♂: Ninette, Manitoba 20.V. 1958 J. F.
 McAlpine/ HOLOTYPE *Phorocera*
(Pseudotachinomyia) exigua Wood CNC No.
 10932.
 Allotype ♀: Same data as holotype except with allotype
 label.
 Paratypes: 51♂♂, 7♀♀, Manitoba; 2♀♀, Ontario; 1♂,
 2♀♀, Quebec.

Phorostoma rutilans Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 504.
 SOUTH AFRICA.
 Syntype: 1♂.
 ♂: 562/ [top side of label illegible] [underside of
 label] Guigoni VY Marley 4.15/ *Phorostoma*
rutilans Typ. ♂ Villen./ TYPE/ CNC Syntype
Phorostoma rutilans Villeneuve Label affixed
 1994/ EX L.-P. MESNIL COLLECTION 1985.

Phorostoma takanoi Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 67.
 JAPAN.
 Holotype ♂: Obihiro S. Takano/ [underside of label]
 6.VII.'52/ 78/ TYPE.

Phryno aprica Villeneuve, 1912b, Bull. Mus. natn. Hist. nat., Paris
 18: 509. ALGERIA.
 Lectotype ♂: Alger. IV 52418/ *Phryno aprica* Villen./
A. aprica Vill. L.P. Mesnil det., 1969/ TYPE/
 CNC Syntype *Phryno aprica* Villeneuve Label
 affixed 1994/ Lectotype fixation by Mesnil, 1956:
 483. Label affixed 1995/ EX L.-P. MESNIL
 COLLECTION 1970.
 Remarks: We regard Mesnil's (1956a: 483) reference to
 a holotype as a lectotype fixation for this species.

Phryno katoi Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 11.
 JAPAN.
 Holotype ♂: V.7. 1950 Nikkō S.KATO/ 346/ *Phryno*
katoi Mesnil L.P. Mesnil det., 1960/ TYPE/ EX
 L.-P. MESNIL COLLECTION 1970.
 Remarks: Holotype terminalia are stored in glycerine in
 a plastic microvial pinned below specimen.

Phylacteropoda nigra Brooks, 1945b, Can. Ent. 77: 93. CANADA.
 Holotype ♀: Transcona, Man. 29.VII.24 G. S. Brooks/
 HoloTYPE ♀ Phylacteropoda nigra No. Brks
 5535.

Phyllomyia fuscicosta Curran, 1927a, Bull. Brooklyn ent. Soc. 22:
 147. CANADA.
 Holotype ♂: Phyllomyia TYPE fuscicosta Curran. No./
 Seton Lake Lillooet, B.C. 28.V 1926
 J.McDunnough.

Phyllomyia nobilis Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 71.
 JAPAN.
 Holotype ♀: JAPAN S.Takano [underside of label]
 12-13/vi 54/ 227/ Phyllomyia nobilis Mesnil L.P.
 Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL
 COLLECTION 1970.

Phyllomyia takanoi Mesnil, 1970, Mushi 44: 119. JAPAN.
 Holotype ♂: Nukabira Hokkaido 25.VII.1964
 S.Takano/ 377/ Phyllomyia takanoi Mesnil L.P.
 Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL
 COLLECTION 1985.

Phytomyptera lacteipennis Villeneuve, 1934b, Revue Zool. Bot. afr. 26: 71. EGYPT.
 Holotype ♀: 3e Tour. Route Suez. 23.3.23/
 Coll.Efflatoun EGYPT/ Phytomyptera
 lacteipennis Vill. L.P. Mesnil det., 1975/ TYPE/
 HOLOTYPE ♀ Phytomyptera lacteipennis Mesnil
 det. Stig Andersen/ EX L.-P. MESNIL
 COLLECTION 1970.
 Remarks: Although no holotype was designated in the
 description, this specimen is believed to have been
 the only one before Villeneuve at the time of
 description.

Phytomyptera rufescens Villeneuve, 1936a, Bull. Mus. r. Hist. nat. Belg. 12(4): 3. SOUTH AFRICA.
 Holotype ♀: M fongosi Zulu L. W E Jones/
 Phytomyptera rufescens n.sp./ TYPE/ EX L.-P.
 MESNIL COLLECTION 1970.

Phytomyptera walleyi Brooks, 1945b, Can. Ent. 77: 91. CANADA.
 Holotype ♀: Kazubazua, Que. 16.VIII 1927 G.S.
 Walley/ HoloTYPE ♀ Phytomyptera walleyi Brks
 No. 5532.
 Paratype: 1♀, Ontario.

Phytomyzoneura aristalis Villeneuve, 1936a, Bull. Mus. r. Hist. nat. Belg. 12(4): 2. SOUTH AFRICA.
 Holotype ♀: S. W. Distr Cape Col./ Phytomyzoneura
 aristatis Villen. L.P. Mesnil det., 1970/ TYPE/ EX
 L.-P. MESNIL COLLECTION 1970/ Elfia
 aristalis ♀ (Villeneuve, 1936) 1984 Det. D.
 Barraclough.

Picconia incurva acanthina Mesnil, 1961b, Fliegen palaearkt. Reg. 64g: 667. CROATIA.
 Holotype ♂: 20/5-13/ Picconia acanthina Mesnil L.P.
 Mesnil det., 1969/ TYPE.

Plagia hilfii sinaica Villeneuve, in Hermann & Villeneuve, 1909,
 Verh. naturw. Ver. Karlsruhe 21: 157. EGYPT.
 Holotype ♀: Sinai/ var. sinaica Villen/ Plagia Hilfii
 Strobl/ Plagiomima sinaica Vill. L.P. Mesnil det.,
 1975/ TYPE/ EX L.-P. MESNIL COLLECTION
 1970.

Plagiocoma crassiseta Villeneuve, 1916c, Ann. S. Afr. Mus. 15:
 474. SOUTH AFRICA.
 Holotype ♀: port Elizabeth Kaapland 23.VII.1912/
 Plagiocoma crassiseta ♀ Villen./ Linnaemyia
 crassiseta Villen. L.P. Mesnil det., 1970/ TYPE/
 EX L.-P. MESNIL COLLECTION 1970.

Plagiomima euethes Reinhard, 1957, Ent. News 68: 104. USA.
 Holotype ♂: 10miN.Sedona,ARIZ Sept. 13, 1955 G. D.
 Butler/ HOLOTYPE Plagiomima euethes
 Reinhard/ Plagiomima euethes R Rnh.
 Allotype ♀: 15miN.Sedona,ARIZ Sept. 14-16,1955 G.
 D. Butler/ ALLOTYPE Plagiomima euethes
 Reinhard.
 Paratypes: 2♂♂, 1♀, Arizona.

Plagiomima faceta Reinhard, 1957, Ent. News 68: 105. USA.
 Holotype ♂: Plainview Tex 9-25-48/ F. A. Cowan
 Collector/ HOLOTYPE Plagiomima faceta
 Reinhard.
 Allotype ♀: Same data as holotype except with allotype
 label.

Plagiprospherysa trinitatis Thompson, 1968, Mem. ent. Soc. Can. 56: 91. TRINIDAD.
 Holotype ♂: HOLOTYPE Plagiprospherysa trinitatis
 Thomp. CNC No. 9862/ Trinidad 1964/
 Plagiprospherysa trinitatis n. sp.
 Allotype ♀: ALLOTYPE Plagiprospherysa trinitatis
 Thomp. CNC No. 9862/ 1964 Trinidad, WI.

Platymya (Archimera) oncoperae Mesnil, 1954b, Fliegen
 palaearkt. Reg. 64g: 371. TASMANIA.
 Holotype ♂: 3/ Breadalbane Tasmania December 1933/
 Bred from Larvae of Oncopera/ Archimera
 oncoperae Mesn. L.P. Mesnil det., 1969/ TYPE/
 Austrophryno densa Walk. L.P. Mesnil det., 1969/
 EX L.-P. MESNIL COLLECTION 1970.

Platymya (Platymya) nemestrina genibarbis Mesnil, 1954b,
 Fliegen palaearkt. Reg. 64g: 375. GERMANY.
 Holotype ♀: Rehbrücke 23.IX.06/ Platymya genibarbis
 Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P.
 MESNIL COLLECTION 1970.

Pleropeleteria peringueyi Villeneuve, 1916c, Ann. S. Afr. Mus. 15:
 471. SOUTH AFRICA.
 Holotype ♂: [word illegible]/ Pleropeleteria ♂
 Peringueyi n.sp. Villen./ TYPE/ Peleteria litantrax
 Wied. L.P. Mesnil det., 1970/ HOLOTYPE of
 Pleropeleteria peringueyi Vill. Examined 1970
 R.W.Crosskey.

Plesiooestrus albifacies Villeneuve, 1914, Annls hist.-nat. Mus.
 natn. hung. 12: 441. ZAIRE.
 Holotype ♀: Kibombo 2-XI-1910/ Plesiooestrus
 albifacies Type Villen./ Plesiooestrus albifacies

- Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Plesiooestrus leonidei* Mesnil, 1964, Bull. Soc. ent. Fr. 69: 262. FRANCE.
Holotype ♂: P 199 J.C.LÉONiDE Ex Ephilliger ephilliger Fieb. Massif de la Sainte. Baume/ Plesiooestrus leonidei Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Policheta crassispinosa* Wood, 1985, Mem. ent. Soc. Can. 132: 77. USA.
Holotype ♀: USA Oregon Benton Co., Alsea Falls Rec. Area 16km.s.w. Alpine 31.VIII-1.IX.1980 J.E. O'Hara/ ♀/ HOLOTYPE Policheta crassispinosa Wood CNC No. 18432.
Paratype: 1♀, Oregon.
Remarks: Holotype is missing abdomen.
- Politomyia angulineura* Reinhard, 1935c, Ann. ent. Soc. Am. 28: 166. USA.
Holotype ♂: Puyallup Wash/ Wm W.Baker Coll VII.6 '32/ HOLOTYPE Politomyia angulineura Reinhard/ Euscopilia dakotensis det. H.J. REIN Tns.
- Procistogaster unicolor* Brooks, 1946, Can. Ent. 77 (1945): 229. USA.
Holotype ♀: College Sta. Tx. 11-4-19/ H.J. Reinhard Collector/ Holo.TYPE Procistogaster unicolor No. Brs/ Gymnoclytia unicolor R Brks./ Gymnoclytia unicolor (Brks) Reinhard Coll'n.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 6♂♂, 3♀♀, Texas.
- Prodemoticus orientalis* Villeneuve, 1919, Bull. Soc. ent. Fr. 1919: 265. HUNGARY.
Lectotype ♂: Ungarn 53346/ Prodemoticus orientalis Villen.Typ/ Prodemoticus orientalis Villen. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype Prodemoticus orientalis Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1973: 1156. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: We regard Mesnil's (1973a: 1156) reference to a holotype as a lectotype fixation for this species.
- Promedina japonica* Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 26. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label] 23.VII.54/ 13/ Promedina japonica Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Proxystomima claripennis* Villeneuve, 1925, Konowia 4: 51. NIGERIA.
Syntypes: 1♂, 1♀.
♂: Rutschuru XI-1914/ Proxystomima claripennis Typ. Villen./ Plesiooestrus claripennis Vill. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype Proxystomima claripennis Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- ♀: Capt. Blau Ilesha, S.Nigeria. 3/11/ Proxystomima claripennis Vill. L.P. Mesnil det., 1970/ Proxystomima claripennis Typ. Villen./ CNC Syntype Proxystomima claripennis Villeneuve Label affixed 1994.
- Prozenillia distans* Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 488. SOUTH AFRICA.
Syntypes: 2♂♂.
♂: Gardens Bay Lightfoot Jan 1915/ Prozenillia distans Villen Typ./ Aplomyia distans Villen. L.P. Mesnil det., 1969/ TYPE/ ?THIS SPECIMEN NO TYPE-STATUS R.W.Crosskey det./ CNC Syntype ? Prozenillia distans Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
♂: Gardens Bay Lightfoot Jan 1915/ S.A.M./ Prozenillia distans Typ. Villen./ Aplomyia distans Villen. L.P. Mesnil det., 1969/ CNC Syntype ? Prozenillia distans Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: These specimens are not known with certainty to belong to the syntype series.
- Psalidopteryx brunnea* Brooks, 1942, Can. Ent. 74: 147. CANADA.
Holotype ♂: HoloTYPE ♂ Psalidopteryx brunnea Brs No.5311/ Lethbridge, Alta. Jun 6 1923 H.L.Seamans.
Allotype ♀: Medicine Hat AltaCan/ Oct 1911/ JRMalloch Coll/ AlloTYPE ♀ Psalidopteryx brunnea Brs No. 5311.
Paratypes: 2♂♂, Alberta.
- Psalidopteryx fuscisquamis* Brooks, 1942, Can. Ent. 74: 149. CANADA.
Holotype ♂: HoloTYPE ♂ Psalidopteryx fuscisquamis No. Brs 5314/ WatertonAlta. JUL 14 1923 H.L.Seamans.
Remarks: Date of collection of holotype was published as 11 July 1923.
- Psalidopteryx macdunnoughi* Brooks, 1942, Can. Ent. 74: 146. CANADA.
Holotype ♂: HoloTYPE ♂ Psalidopteryx macdunnoughi No. Brooks 5307./ Alberton, P.E.I. 28 III -1940 J.McDunnough/ Host. Scythris.
Allotype ♀: Same data as holotype except with allotype label and a label with "Host. Scythris sp".
Paratypes: 4♂♂, 6♀♀, Prince Edward Island.
- Psalidopteryx nuda* Brooks, 1942, Can. Ent. 74: 147. CANADA.
Holotype ♂: HoloTYPE Psalidopteryx nuda Brs No. 5310/ W.G.Mathers Vancouver, B.C. IV-18-1936/ ?Parasite of Griselda radicana?/ 17580 Loc7.
- Psalidopteryx pollinosa* Brooks, 1942, Can. Ent. 74: 148. CANADA.
Holotype ♂: HoloTYPE ♂ Psalidopteryx pollinosa No.5312 Brs/ Chilcotin, B.C. 12. V. 1920 E.R.Buckell.

Allotype ♀: Trinity Valley, B.C. Aug. 3 1937 K.
 Graham/ AlloTYPE ♀ *Psalidopteryx pollinosa* No.
 Brs 5312.
 Paratype: 1♀, British Columbia.

Psalidopteryx pseudonuda Brooks, 1942, Can. Ent. 74: 147.
 CANADA.

Holotype ♂: HoloTYPE *Psalidopteryx pseudonuda* No.
 5309 Brs./ Clinton, B.C. Ex Peronea varicana/
 Rec:B.C. 1497.A F.I.Survey 1940 Em. 1.III. 1941.
 Remarks: Holotype puparium is pinned below
 specimen.

Psalidopteryx psilocorsiphaga Brooks, 1942, Can. Ent. 74: 149.
 CANADA.

Holotype ♂: HoloTYPE ♂ *Psalidopteryx*
psilocorsiphaga No. Brs 5315/ Berthierville Que.
 20-6 1932 L.Davault/ Host. Psilocorsis sp.
 -Birch-.
 Allotype ♀: Same data as holotype except with allotype
 label and dated "15-6 1932".
 Paratypes: 1♂, 4♀ ♀, Quebec.

Pseudalsomyia piligena Mesnil, 1968a, Bull. Annls Soc. r. ent.
 Belg. 104: 178. PAKISTAN.

Holotype ♂: 1204/ Ex stem borers of *Pinus excelsa*/
 Paras 6. 2. 63/ C.I.B.C. PG.2/63-325/
Pseudalsomyia piligena Mesn. L.P. Mesnil det.,
 1967/ TYPE/ EX L.-P. MESNIL COLLECTION
 1970.

Pseudapinops rogalis Reinhard, 1955c, Bull. Brooklyn ent. Soc.
 50: 132. USA.

Holotype ♀: Marion Mt Cmp San Jacinto Mts
 Cal.VII-1-1952/ E. M. Evans Collector/
 HOLOTYPE *Pseudapinops rogalis* Reinhard/
Pseudapinops rogalis R. Rnh.

Pseudocelatoria robusta Thompson, 1968, Mem. ent. Soc. Can. 56:
 190. TRINIDAD.

Holotype ♀: MARACAS V. Trinidad, BWI.
 NOV.24.1952/ Collector F.J.Simmonds/ 48/
Pseudocelatoria robusta n.sp./ 1015b. V.27.59
 [underside of label] Ottawa, Ont. 195
 W.R.Thompson/ Species New sp. U.
 Det.Thompson/ HOLOTYPE *Pseudocelatoria*
robusta Thomp CNC No.

Pseudochaeta canadensis Brooks, 1945b, Can. Ent. 77: 93.
 CANADA.

Holotype ♀: No. 12139-27/ C.E. Atwood 7. VIII 1940
 Laniel, Que./ Lexis sp./ HoloTYPE ♀
Pseudochaeta canadensis No. Brks 5536/ SLIDE
 0506 No.
 Allotype ♂: Lepidoptera pupa./ C.E. Atwood 25.VII
 1940 Laniel, Que./ Allo TYPE ♂ *Pseudochaeta*
canadensis No. Brks 5536.
 Paratypes: 3♀ ♀, Quebec; 1♀, Ontario; 1♀, Manitoba;
 2♀ ♀, British Columbia.

Remarks: Holotype and allotype puparia are pinned
 below their respective specimens. The holotype
 slide has not been located; the holotype is intact
 and the puparium is missing only the

cephalopharyngeal skeleton, so the slide might
 contain that structure.

Pseudochaeta clurina Reinhard, 1946b, Can. Ent. 78: 119. USA.
 Holotype ♂: College Station Apr. 6, 1944 Tex/ H J
 Reinhard Collector/ HOLOTYPE *Pseudochaeta*
clurina Reinhard/ *Pseudochaeta clurina* R. Rnh.
 Paratype: 1♂, Texas.

Pseudochaeta curepei Thompson, 1964, Studia ent. 7: 114.
 TRINIDAD.

Holotype ♀: HOLOTYPE *Pseudochaeta curepei*
 Thomp. CNC No. 9818/ I.C.T.A. Trinidad, BWI.
 APRIL.7.1954./ Collector F.J.Simmonds/ N.
 Paratypes: 4♀ ♀, Trinidad.

Pseudochaeta finalis Reinhard, 1946b, Can. Ent. 78: 114. USA.
 Holotype ♀: College Station Apr 28, 1943 Tex/ H J
 Reinhard Collector/ HOLOTYPE *Pseudochaeta*
finalis Reinhard/ *Pseudochaeta finalis* R. Rnh.

Pseudochaeta flavigalpis Thompson, 1964, Studia ent. 7: 101.
 TRINIDAD.

Holotype ♂: HOLOTYPE *Pseudochaeta flavigalpis*
 Thomp. CNC No. 9814/ St. Augustine,
 Trinidad, BWI. NOV.18.1952/ Collector
 F.J.Simmonds/ 1319 VIII-13.63/ P. *flavigalpis*
 n.sp/ sp. B.
 Allotype ♀: ALLOTYPE *Pseudochaeta flavigalpis*
 Thomp. CNC No. 9814/ I.C.T.A. Trinidad, BWI.
 APRIL.1.1954./ Collector F.J.Simmonds/ Sp.F
 1217.II.8 62.

Remarks: Holotype is missing terminalia.

Pseudochaeta frontalis Reinhard, 1946b, Can. Ent. 78: 114. USA.

Holotype ♂: College Station May 15, 1943 Tex/ H J
 Reinhard Collector/ HOLOTYPE *Pseudochaeta*
frontalis Reinhard/ *Pseudochaeta frontalis* R. Rnh.
 Allotype ♀: Same data as holotype except with allotype
 label and dated "May 26, 1943".

Pseudochaeta latitarsus Thompson, 1964, Studia ent. 7: 111.
 TRINIDAD.

Holotype ♀: HOLOTYPE *Pseudochaeta latitarsus*
 Thomp. CNC No. 9817/ ST. BENEDICT.
 Trinidad, BWI. MARCH.5 1954./ Collector
 F.J.Simmonds/ sp.2X Det./ 109/ *Pseudochaeta*
latitarsus n.sp.

Pseudochaeta marginalis Reinhard, 1946b, Can. Ent. 78: 118.
 USA.

Holotype ♂: College Station 6.23.19 Texas/ H J
 Reinhard Collector/ HOLOTYPE *Pseudochaeta*
marginalis Reinhard/ *Pseudochaeta marginalis* R.
 Rnh.
 Allotype ♀: College Station 7-25-23 Texas/ H J
 Reinhard Collector/ 13361- X.7.63 3/ ALLOTYPE
Pseudochaeta marginalis Reinhard.
 Paratype: 1♂, Texas.

Pseudochaeta nitens Thompson, 1964, Studia ent. 7: 108.

TRINIDAD.
 Holotype ♂: HOLOTYPE *Pseudochaeta nitens* Thomp.
 CNC No. 9816/ Orange Grove TRINIDAD W.I.

- IX 27 60 W.R.Thompson/ sp C/ Pseudochaeta nitens n.sp.
 Paratypes: 2♂♂, Trinidad.
 Remarks: The date "1961" appears after "IX 27 60" on the locality label but has been crossed out.
- Pseudochaeta perdecora* Reinhard, 1946b, Can. Ent. 78: 115. USA.
 Holotype ♀: College Station May 21,1943 Tex/ H J Reinhard Collector/ HOLOTYPE Pseudochaeta perdecora Reinhard/ Pseudochaeta perdecora R. Rnh.
 Paratypes: 4♀♀, Texas.
 Remarks: There are five specimens from College Station, Texas labeled as paratypes but only four are listed in the original description.
- Pseudochaeta siminina* Reinhard, 1946b, Can. Ent. 78: 117. USA.
 Holotype ♂: Amherst, Ohio July, 1933/ H J Reinhard Collector/ HOLOTYPE Pseudochaeta siminina Reinhard/ Pseudochaeta siminina R. Rnh.
 Allotype ♀: Same data as holotype except with allotype label.
- Pseudochaeta syngamiae* Thompson, 1964, Studia ent. 7: 105. TRINIDAD.
 Holotype ♂: HOLOTYPE Pseudochaeta syngamiae Thomp. CNC No. 9815/ June 1963 CUREPE TRINIDAD/ ex Syngamia oulantana/ F.D.BENNETT Collector.
 Allotype ♀: Same data as holotype except with allotype label.
 Paratypes: 4♂♂, 2♀♀, Trinidad.
- Pseudochaeta trinitatis* Thompson, 1964, Studia ent. 7: 117. TRINIDAD.
 Holotype ♀: HOLOTYPE Pseudochaeta trinitatis Thomp. CNC No. 9819/ CARONI Trinidad,BWI. SEPT.30 1952/ Collector F. D. Bennett/ P. nsp trinitatis.
 Paratypes: 2♀♀, Trinidad.
- Pseudogonia madagascariensis* Villeneuve, 1915a, Revue zool. afr. 4: 192. MADAGASCAR.
 Holotype ♀: Madagascar Tananarive Lamberton 1912/ Pseudogonia madagascarien-sis Villen. Typ./ Isomera madagascariensis L.P. Mesnil det., 1969 Vill./ TYPE.
 Remarks: We consider this specimen to be a holotype, under the assumption that there were no other specimens (syntypes) in the type series.
- Pseudolomyia scissilis* Reinhard, 1962a, Ent. News 73: 176. USA.
 Holotype ♂: Amherst Ohio Jul-Aug 1958 H.J.Reinhard/ HOLOTYPE Pseudolomyia scissilis Reinhard.
- Pseudomasiphya petiolata* Thompson, 1963d, Can. Ent. 95: 1316. TRINIDAD.
 Holotype ♀: HOLOTYPE Pseudomasiphya petiolata Thomp. CNC No. 9812/ MARACAS V. Trinidad,BWI. NOV.17.1952./ Collector F.J.Simmonds/ HOLOTYPE CNCNo.
 Paratype: 1♀, Trinidad.
- Pseudoperichaeta pilosa* Villeneuve, 1942a, Bull. Soc. ent. Fr. 47: 52. ZIMBABWE.
 Syntype: 1♂.
 ♂: GOTA GOTA URUNGWE Dep. Agric. S. Rhodesia 29/7/1938/ W.L. Williams Collector/ Coll.8591/ Pseudoperichaeta pilosa n.sp. Cotyp. Villen./ CNC Syntype Pseudoperichaeta pilosa Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Pseudoperichaeta (Achaetoneurilla) madecassa* Mesnil, 1939b, Bull. Annls Soc. ent. Belg. 79: 210. MADAGASCAR.
 Syntypes: 3♂♂, 2♀♀.
 2♂♂, ♀: MADAGASCAR BEKILY REG. SUD DE L'ILE/ XI. 33 A. SEYRIG/ Achaetoneurilla madecassa Mesnil L.P. Mesnil det., 1969/ CNC Syntype Pseudoperichaeta (Achaetoneurilla) madecassa Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
 ♂, ♀: MADAGASCAR BEKILY REG. SUD DE L'ILE/ Achaetoneurilla madecassa Mesnil L.P. Mesnil det., 1985/ CNC Syntype Pseudoperichaeta (Achaetoneurilla) madecassa Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1985.
- Pseudoprospherysa claripalpis* Thompson, 1968, Mem. ent. Soc. Can. 56: 79. TRINIDAD.
 Holotype ♀: HOLOTYPE Pseudoprospherysa claripalpis Thomp. CNC No. 9858/ ARIPO V. TRINIDAD 10-7-64/ Pseudoprospherysa claripalpis n sp.
 Remarks: Date of collection of holotype was published as 7 July 1964.
- Pseudoprospherysa santacruzi* Thompson, 1968, Mem. ent. Soc. Can. 56: 74. TRINIDAD.
 Holotype ♂: HOLOTYPE Pseudoprospherysa santacruzi Thomp. CNC No. 9856/ Santa Cruz V. 3. 2. 60 Trinidad,BWI/ 1487 VIII.3.66.
 Allotype ♀: ALLOTYPE Pseudoprospherysa santacruzi Thomp. CNC No. 9856/ LABA JAR TRINIDAD W.I. 3-3-1961 N.Gopaul Coll./ 1460 VII.5.65.
 Paratype: 1♂, Trinidad.
 Remarks: Holotype is missing terminalia.
- Pseudoprospherysa similis* Thompson, 1968, Mem. ent. Soc. Can. 56: 77. TRINIDAD.
 Holotype ♀: HOLOTYPE Pseudoprospherysa similis Thomp. CNC No. 9857/ St Augustine Trinidad WI/ 1090 III.14.60/ Pseudoprospherysa similis n.sp/ [white label with left wing and three left legs glued to it].
- Pseudorrhincta rubricornis* Thompson, 1968, Mem. ent. Soc. Can. 56: 120. TRINIDAD.
 Holotype ♂: HOLOTYPE Pseudorrhincta rubricornis Thomp. CNC No. 9868/ III.2.64 Brazil Trinidad, WI.
- Pseudosiphonsturmia aberrans* Thompson, 1966, Studia ent. 8 (1965): 415. TRINIDAD.

- Holotype ♀: HOLOTYPE Pseudosiphosturmia aberrans
Thomp. CNC No.9838/ Champs fleurs
TRINIDAD W.I. IX 28 '60 W.R.Thompson/ 1253
X.19.62/ Pseudosiphosturmia aberrans n.g; nsp.
Allotype ♂: ALLOTYPE Pseudosiphosturmia aberrans
Thomp. CNC No. 9838/ JAN 8 1960 AT LIGHT
St. Augustine, Trinidad,BWI/ F.D.Bennett
Collector/ 1277 XII.7.62.
Paratype: 1♀, Trinidad.
Remarks: Year of collection of holotype was published
as 1961.
- Pseudosturmia clavipalpis* Thompson, 1966, Studia ent. 8 (1965):
424. TRINIDAD.
Holotype ♂: HOLOTYPE Pseudosturmia clavipalpis
Thomp. CNC No. 9840/ 1369 II.15.63/
Pseudosturmia nsp clavipalpis/ S. Cruz V. II.14.64
Trinidad, WI.
Allotype ♀: ALLOTYPE Pseudosturmia clavipalpis
Thomp. CNC No. 9840/ American Rd. Mauvan
Hill XII.20.60. Trinidad,BWI/ 1325 IX.23.63.
- Pseudotachinomyia aequalis* Reinhard, 1935b, Ent. News 46: 133.
USA.
Holotype ♂: College Station 4-19-20 Texas/ H J
Reinhard Collector/ HOLOTYPE
Pseudotachinomyia aequalis Reinhard/
Pseudotachinomyia aequalis Rein.
det.H.J.Reinhard.
Paratypes: 3♀ ♀, Texas.
- Pseudotachinomyia compascua* Reinhard, 1935b, Ent. News 46:
135. USA.
Holotype ♀: Moses Coulee, Wn. [Washington] IV-2
'33/ J. Wilcox, Coll./ HOLOTYPE
Pseudotachinomyia compascua Reinhard/
Pseudotachinomyia compascua Rein.
det.H.J.Reinhard.
- Pseudotachinomyia longiforceps* Brooks, 1945b, Can. Ent. 77: 89.
CANADA.
Holotype ♂: Ottawa Ont. 23.IV. 1938 G.E.Shewell/
HoloTYPE ♂ Pseudotachinomyia longiforceps No.
Brks 5515.
Allotype ♀: Aylmer, Que. 11-V 1925 G.S.Walley/
AlloTYPE ♀ Pseudotachinomyia longiforceps No.
Brks 5515.
Paratype: 1♀, Quebec.
- Psilopleura arida* Reinhard, 1943a, Can. Ent. 75: 166. USA.
Holotype ♀: Salt Lake City, Ut. 6-15-37/ G. F.
Knowlton Collector/ HOLOTYPE Psilopleura
arida Reinhard/ Psilopleura arida Rnh. Reinhard
Coll'n/ Psilopleura arida Rnh det.H.J.Reinhard.
Paratype: 1♀, New Mexico.
- Ptilodexia agilis* Reinhard, 1943b, J. Kans. ent. Soc. 16: 22. USA.
Holotype ♂: College Station Oct.8,1933 Tex/ H J
Reinhard Collector/ HOLOTYPE Ptilodexia agilis
Reinhard.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 7♂♂, 32♀ ♀, Texas.
- Ptilodexia westi* Wilder, 1979, Proc. Calif. Acad. Sci. 42: 27. USA.
- Holotype ♂: Imperial Tex. IV-10-54 LD Beamer/
Reinhard Collection/ HOLOTYPE Ptilodexia
westi Wilder.
- Punaclista fidelis* Reinhard, 1967, J. Kans. ent. Soc. 40: 98. PERU.
Holotype ♂: LIMA 30. XII. 63. Raven, Coll./ SNA
63-63/ HOLOTYPE Punaclista fidelis Reinhard.
Allotype ♀: LIMA 10. XII. 63. K. Raven, Coll./ SNA
18-63/ ALLOTYPE Punaclista fidelis Reinhard.
Paratype: 1♀, Peru.
Remarks: Holotype puparium is pinned below
specimen.
- Rhacodinella aurata* Mesnil, 1970, Mushi 44: 108. JAPAN.
Holotype ♂: Omogokei Shikoku 6.VI.1964 S.Takano/
407/ Rhacodinella aurata Mesnil L.P. Mesnil det.,
1969/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.
- Rhedia nigra* Brooks, 1944a, Can. Ent. 75 (1943): 225. CANADA.
Holotype ♀: Swift Current, Sask. 15 V -1938
A.R.Brooks/ HoloTYPE ♀ Rhedia nigra No. Brks
5451.
Allotype ♂: Lillooet, B.C. 27. III 1941 A.Phair/
AlloTYPE ♂ Rhedia nigra No. Brks 5451.
Paratype: 1♀, British Columbia.
- Rhedia setifacies* Brooks, 1944a, Can. Ent. 75 (1943): 225.
CANADA.
Holotype ♂: Pike Lake, Sask. 28.IV -1940 A. R
Brooks/ HoloTYPE ♂ Rhedia setifacies No. Brks
5450.
Allotype ♀: Same data as holotype except with allotype
label.
Paratypes: 2♂♂, 1♀, Manitoba; 1♂, Saskatchewan; 1♀,
Idaho.
- Rhinotachina ancyrensis* Villeneuve, 1942b, Bull. Soc. ent. Fr. 47:
134. TURKEY.
Holotype ♂: [small red blank label]/ Rhinotachina
ancyrensis Typ. Villen./ Bithia ancyrensis Vill.
L.P. Mesnil det., 1975/ TYPE.
- Rhodogyne canadensis* Brooks, 1946, Can. Ent. 77 (1945): 220.
CANADA.
Holotype ♀: Queen's Park, Aylmer, Q 27.8. 1926
C.B.Hutchings/ Holo TYPE ♀ Gymnosoma
canadense Brks No. 5345.
Allotype ♂: Sherrington P.Q. 22.VIII. 1927 G. H.
Hammond/ AlloTYPE ♂ Gymnosoma canadense
No. Brks 5345.
Paratypes: 5♂♂, 6♀ ♀, Quebec; 3♂♂, Ontario.
- Rhodogyne emdeni* Mesnil, 1950c, Bull. Annls Soc. ent. Belg. 86:
114. ZIMBABWE.
Holotype ♂: Gatooma S. Rhodesia, Nov. 1934
A.Cuthbertson/ 3834 EX. Pentatomid Agonoscelis
versicolor F./ Gymnosoma emdeni Mess. L.P.
Mesnil det., 1971/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.
Remarks: Holotype puparium is pinned to card mount.
- Rhodogyne persica* Mesnil, 1952c, Bull. Annls Soc. ent. Belg. 88:
152. IRAN.

Holotype ♂: TYPE/ IRAN Keredj, 40km. de Teheran, 14.IV.1936, coll. F.H. Brandt/ Rhodogyne persica n. sp., L. Mesnil det. 1952.

Rhodogyne rungsi Mesnil, 1952c, Bull. Annls Soc. ent. Belg. 88: 151. MOROCCO.
Holotype ♂: 48/ Rabat 17.VIII.46 Larve à Aetia/ Rhodogyne rungsi n.sp. L. Mesnil det., 1952/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Rhodogyne verbekei Mesnil, 1952c, Bull. Annls Soc. ent. Belg. 88: 150. FRANCE.
Holotype ♀: Reuilbeau 30 Juillet 1941/ Rhodogyne verbekei n.sp. L. Mesnil det., 1952/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.
Remarks: The type locality was published as Marchais-Béton, Yonne.

Rileymia triseta Brooks, 1947, Can. Ent. 78 (1946): 181. CANADA.
Holotype ♂: VICTORIA B.C. APRIL 26 1916 R. C. Treherne/ Holotype ♂ Rileymyia triseta Brks No. 5692.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 23♂♂, 10♀♀, British Columbia; 3♂♂, 1♀, Oregon; 1♀, Alberta.

Rondaniooestrus apivorus Villeneuve, 1916b, Ann. S. Afr. Mus. 15: 467. SOUTH AFRICA.
Holotype ♂: Port Elizabeth, Cape Prov., Nov. 17, 1915/ Collected by F.W. Fitzsimmons, bred from honey bee./ Rondaniooestrus apivorus, Type ♂, Villen.

Roubaudia pruinosa Roubaud & Villeneuve, 1914, Revue zool. afr. 4: 123. SENEGAL.
Syntype: 1♀.
♀: Hte.Gambie/ Polistes B/ Roubaudia (Anacamptomyia) pruinosa Villen./ A. pruinosa Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Roubaudia pruinosa Roubaud & Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Roubaudia rufescens Villeneuve, 1910b, Wien. ent. Ztg 29: 249. CONGO.
Syntype: 1♂.
♂: ♂/ Roubaudia rufescens type Villen./ Anacamptomyia eod.nom./ A. rufescens Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Roubaudia rufescens Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Roubaudia rufescens var. *pallida* Roubaud & Villeneuve, 1914, Revue zool. afr. 4: 124. AFRICA.
Syntype: 1♂.
♂: M.fongoss Zulu L. W E Jones/ Roubaudia rufescens var. pallida Villen/ A. pallida Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Roubaudia rufescens var. pallida Roubaud & Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Zaire, Nigeria, Zimbabwe and Senegal are listed as type localities in the original description.

Roubaudia (*Pararoubaudia*) *bisetosa* Roubaud & Villeneuve, 1914, Revue zool. afr. 4: 125. SENEGAL.

Syntype: 1♂.
♂: ♂/ Pararoubaudia bisetosa Villen./ A. bisetosa Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Roubaudia (*Pararoubaudia*) *bisetosa* Roubaud & Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Salia mageritensis Villeneuve & Mesnil, 1936, Bull. Soc. ent. Fr. 41: 274. SPAIN.

Holotype ♂: [white label with holotype terminalia glued to it.]/ Madrid J. GIL/ Chaetogena mageritensis Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Salia rondaniana Villeneuve, 1931, Konowia 10: 57. FRANCE.

Holotype ♀: Digne VI-VII/ Digne S. France VI - VII/ Chaetogena rondaniana Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: We consider this specimen to be a holotype, under the assumption that there were no other specimens (syntypes) in the type series.

Santacruzia dubiosa Thompson, 1968, Mem. ent. Soc. Can. 56: 125. TRINIDAD.

Holotype ♀: HOLOTYPE Santacruzia dubiosa Thomp. CNC No. 9870/ Santa Cruz V. XII.20.60/ Trinidad, BWI/ Santa Cruzia/ 1491 X.14.66.

Santacruziopsis claripennis Thompson, 1968, Mem. ent. Soc. Can. 56: 165. TRINIDAD.

Holotype ♀: HOLOTYPE Santacruziopsis claripennis Thomp. CNC No. 9879/ St. Augustine, Trinidad, BWI. 2. 1. 60.

Allotype ♂: ALLOTYPE Santacruziopsis claripennis Thomp. CNC No. 9879/ 1966 1.25.64.

Paratypes: 2♀♀, Trinidad.

Santacruziopsis fumipennis Thompson, 1968, Mem. ent. Soc. Can. 56: 163. TRINIDAD.

Holotype ♀: HOLOTYPE Santacruziopsis fumipennis Thomp. CNC No. 9878/ SANTA CRUZ Trinidad, BWI. FEB.12.1953./ Collector F.J.Simmonds/ 10 10X.(X) V.14.59 [underside of label] W.R.Thompson Ottawa, Ont. 195 W.R.Thompson 1010.

Sarrorrhina pupilla Villeneuve, 1936a, Bull. Mus. r. Hist. nat. Belg. 12(4): 2. SOUTH AFRICA.

Syntypes: 2♂♂, 1♀.
♂: Wint-hoek [Wind-hoek] Tulbagh 3600f./ Sarrorrhina pupilla Type n.sp./ Sarrorrhina pupillata [sic] Vill. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype Sarrorrhina pupilla Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
♂: Wind. hoek Tulbagh 3600 ft./ Sarrorrhina pupillata [sic] Vill. L.P. Mesnil det., 1970/ CNC Syntype Sarrorrhina pupilla Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
♀: tarses ant. dilates ♀/ Wind. hoek Tulbagh 3600 ft/ Sarrorrhina pupillata [sic] Vill. L.P. Mesnil det.,

1970/ CNC Syntype Sarrorrhina pupilla Villeneuve
Label affixed 1994/ EX L.-P. MESNIL
COLLECTION 1970.

Saundersiops harpeza Reinhard, 1952b, Bull. Brooklyn ent. Soc. 47: 8. MEXICO.
Holotype ♂: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE
Saundersiops harpeza Reinhard/ Saundersiops harpeza R. Rnh.
Allotype ♀: Rio Frio, Mex. Sept. 3, 1947/ F. A. Cowan M. R. Wheeler Collectors/ ALLOTYPE
Saundersiops harpeza Reinhard/ close or = to rufitibia Wlp Det.D.M.Wood 1989.
Paratypes: 2♂♂, 3♀♀, Mexico.

Scaphomyia takanoi Mesnil, 1967, Mushi 41: 43. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label] 3/IX'52/ 362/ Scaphomyia takanoi Mesnil L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Siphoclytia pavonacea Reinhard, 1939, Bull. Brooklyn ent. Soc. 34: 72. USA.
Holotype ♂: Amherst, Ohio 6. 24 1935/ A.J.Barckert collector/ HOLOTYPE Siphoclytia pavonacea Reinhard.
Paratypes: 2♀♀, Ohio.

Siphokesia loriola Reinhard, 1955a, J. Kans. ent. Soc. 28: 126.
USA.
Holotype ♀: Brownsville VIII-14-45 Tex/ F. A. Cowan Collector/ HOLOTYPE Siphokesia loriola Reinhard/ Siphokesia loriola R Rnh./ Siphokesia loriola Rnh. Reinhard Coll'n.
Allotype ♂: Brownsville VIII-14-45 Tex/ F. A. Cowan Collector/ ALLOTYPE Siphokesia loriola Reinhard/ Siphokesia loriola Rnh. Reinhard Coll'n.
Paratypes: 3♀♀, Mexico; 1♀, Texas.

Siphona akidnomyia O'Hara, 1983, Quaest. Ent. 18 (1982): 311. MEXICO.
Holotype ♂: MEX.Chis.9600ft. Zontehults,nr.S.Crist. 25 June 1969 W.R.M.Mason/ HOLOTYPE
Siphona akidnomyia O'Hara CNC No. 19086/
HOLOTYPE ♂ Siphona akidnomyia O'Hara.
Allotype ♀: MEXICO,Chiapas Mt. Tzontehultz 9400'12mi.NE San Cristobal 27.V. 1969 B.V.Peterson/ ALLOTYPE Siphona akidnomyia O'Hara CNC No. 19086/ ALLOTYPE ♀ Siphona akidnomyia O'Hara.
Paratypes: 3♂♂, 1♀, Mexico.
Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.

Siphona boreata Mesnil, 1960c, Bull. Annls Soc. r. ent. Belg. 96: 190. GERMANY.
Holotype ♂: Arnsbg. Wald 15.6.56/ Arnsberg Germany 15. 6. 1956/ Siphona boreata Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Siphona brunnea O'Hara, 1983, Quaest. Ent. 18 (1982): 308. MEXICO.

Holotype ♂: MEX.Chis.9600ft. Zontehults,nr.S.Crist. 17 May 1969 W.R.M.Mason/ HOLOTYPE
Siphona brunnea O'Hara CNC No. 19082/
HOLOTYPE ♂ Siphona brunnea O'Hara.
Allotype ♀: Same data as holotype except with allotype label.
Paratypes: 1♂, 1♀, Mexico.

Siphona confusa Mesnil, 1961a, Bull. Annls Soc. r. ent. Belg. 97: 201. SWEDEN.

Holotype ♂: Gränna Suède 22.V.1956/ Siphona confusa Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970/ HOLOTYPE ♂ Siphona confusa Mesnil det. Stig Andersen/ Siphona confusa Mesnil = mesnili Andersen det. Stig Andersen 1988.

Remarks: There may be a female paratype in the CNC material, but it is not labeled as such. Holotype terminalia are stored in glycerine in a plastic tube pinned below specimen.

Siphona efflatouni Mesnil, 1960c, Bull. Annls Soc. r. ent. Belg. 96: 188. EGYPT.

Holotype ♀: MARIOUT 16. 3. 23/ Coll.Efflatoun EGYPT/ Siphona efflatouni Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Siphona hokkaidensis Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 36. JAPAN.

Holotype ♀: 295/ Obihiro S.Takano [underside of label] 2-3/VII'49/ Siphona hokkaidensis Mesnil L.P. Mesnil det., 1970/ TYPE/ HOLOTYPE ♀ Siphona hokkaidensis Mesnil det. S. Andersen/ EX L.-P. MESNIL COLLECTION 1970.

Siphona longissima O'Hara, 1983, Quaest. Ent. 18 (1982): 311. MEXICO.

Holotype ♂: MEX.Chis.10mi NE. San Cristobal 13.V.69 7500' H.J.Teskey/ HOLOTYPE Siphona longissima O'Hara CNC No. 19087/ HOLOTYPE Siphona ♂ longissima O'Hara.

Paratype: 1♂, Mexico.
Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.

Siphona lurida Reinhard, 1943b, J. Kans. ent. Soc. 16: 20. USA.

Holotype ♂: Rainier, Ore. III-3 30/ R.E.Dimick Coll./ HOLOTYPE Siphona lurida Reinhard/ Siphona lurida R'43 Rnh.
Paratypes: 2♂♂, Oregon.

Siphona maculata griseola Mesnil, 1970, Mushi 44: 118. ISRAEL.

Holotype ♂: Tel Aviv Israel 18.III 1968 leg. Kugler/
Siphona griseola Mesnil L.P. Mesnil det., 1970/
TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Siphona medialis O'Hara, 1983, Quaest. Ent. 18 (1982): 303.

CANADA.

Holotype ♂: Cranberry I. Lockeport,N.S. 25.VII.1958 J.R.Vockeroth/ HOLOTYPE Siphona medialis O'Hara CNC No.19085/ HOLOTYPE Siphona medialis ♂ O'Hara.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 1♂, 3♀♀, Utah.

Siphona multifaria O'Hara, 1983, Quaest. Ent. 18 (1982): 293.
CANADA.

Holotype ♂: Mi. 87, Y.T. Dempster Hwy.
8-12.VII.1973 G.& D.M.Wood/ HOLOTYPE
Siphona multifaria O'Hara CNC No. 19083/
HOLOTYPE *Siphona* ♂ *multifaria* O'Hara.

Allotype ♀: Same data as holotype except with allotype label and dated "16-17.VII.1973".

Paratypes: 2♀♀, Newfoundland; 5♂♂, 6♀♀, New Brunswick; 10♂♂, 7♀♀, Quebec; 5♂♂, 3♀♀, Ontario; 15♂♂, Alberta; 15♂♂, 8♀♀, British Columbia; 40♂♂, 33♀♀, Yukon; 2♀♀, Northwest Territories; 2♂♂, 10♀♀, Alaska; 2♂♂, 1♀, Arizona; 9♂♂, 5♀♀, New York; 5♂♂, 6♀♀, Texas; 9♂♂, 3♀♀, Colorado; 1♂, New Hampshire; 1♀, Michigan; 1♂, Pennsylvania; 1♀ North Carolina.

Siphona oligomyia O'Hara, 1983, Quaest. Ent. 18 (1982): 297.
CANADA.

Holotype ♂: Keremeos, B.C. 20.VI 1923 C.B.Garrett/
HOLOTYPE *Siphona oligomyia* O'Hara CNC No.
19084/ HOLOTYPE *Siphona* ♂ *oligomyia* O'Hara.

Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.

Siphona pisinnia O'Hara, 1983, Quaest. Ent. 18 (1982): 298. USA.

Holotype ♂: USA N.M. Grant Co. Gila Nat. For.Cherry Ck.Campgrd. 21km.n. Silver City 2250m.
3-4.VIII.1980 J.E. & W.M. O'Hara/ HOLOTYPE
Siphona pisinnia O'Hara CNC No. 19081/
HOLOTYPE *Siphona* ♂ *pisinnia* O'Hara.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 10♂♂, Mexico; 2♂♂, 3♀♀, Arizona; 1♀, California; 17♂♂, 5♀♀, New Mexico.

Siphona silvarum Herting, 1967, Stuttg. Beitr. Naturk. 173: 9.
GERMANY.

Holotype ♂: ex larvae *Tipula irrorata* Mg./ Moorswald Freiburg.L.B. IV. 1958/ *Siphona silvarum* Hert. L.P. Mesnil det., 1970/ TYPE/ HOLOTYPE ♂ *Siphona silvarum* Herting det. Stig Andersen 1986/ EX L.-P. MESNIL COLLECTION 1970/ *Siphona hokkaidensis* Mesnil det. Stig Andersen 1985.

Remarks: Two puparia are glued to paper mount below specimen.

Siphophyto politura Reinhard, 1946a, Proc. ent. Soc. Wash. 48: 85.
CANADA.

Holotype ♂: Laniel,Que 5 vii-1944 A.R.Brooks/
Holotype *Siphophyto politura* Reinhard/
HOLOTYPE ♂ *Siphophyto politura* No. Rnh 5623.
Paratypes: 18♂♂, 1♀, Quebec; 1♂, 1♀, Michigan.

Siphosturmia maltana Reinhard, 1951, Bull. Brooklyn ent. Soc. 46: 7. USA.

Holotype ♂: Malta, Mont. July 26, 1947/ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE

Siphosturmia maltana Reinhard/ *Siphosturmia maltana* R. Rnh.

Allotype ♀: Same data as holotype except with allotype label.

Paratypes: 10♂♂, 2♀♀, Montana.

Siphosturmiosis rufiventris Reinhard, 1934c, Ent. News 45: 18.
USA.

Holotype ♀: Monrovia Canyon Calif. VII-26-1931 C.H & D.Martin/ HOLOTYPE *Siphosturmiosis rufiventris* Reinhard/ *Siphosturmiosis rufiventris* det.H.J.Reinhard Rein.

Sisyropa ghanii Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 176. PAKISTAN.

Holotype ♂: Ex Lepid larva. On *Ficus Palmata*./
.R.Pindi [Rawalpindi] 26. 7.62/ 908/ C.I.B.C. 62
FP31/-/ *Sisyropa ghanii* n.sp. L.P. Mesnil det.,
1967/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.

Sitellitergus aemulus Reinhard, 1964b, Acta zool. mex. 7: 4.
MEXICO.

Holotype ♂: Cuernavaca, IX-6.51 Mex./ HOLOTYPE
Sitellitergus aemulus Reinhard.

Allotype ♀: XI-27-51/ Paso Cortes, Mex alt. 3000 m
W.G. Downs, Coll./ ALLOTYPE *Sitellitergus aemulus* Reinhard/ *Sitellitergus aemulus* R Rnh.
Paratype: 1♂, Mexico.

Sitellitergus simiolus Reinhard, 1964b, Acta zool. mex. 7: 5.
MEXICO.

Holotype ♂: Jalapa, Ver. 9/28-X/3/61 R&K Dreisbach/
HOLOTYPE *Sitellitergus simiolus* Reinhard/
Sitellitergus simiolus R Rnh.

Smidtia magnicornis Mesnil, 1967, Mushi 41: 40. JAPAN.

Holotype ♂: Mt. Moiwa Sapporo 16.V.1966 S. Takano/
415/ TYPE/ *Smidtia magnicornis* Mesn L.P.
Mesnil det., 1967.

Solieria (Anthoica) aureola Mesnil, 1973a, Fliegen palaearkt. Reg. 64g: 1119. JAPAN.

Holotype ♀: Nopporo Japan Natural forest 6. 8. 1961/
Solieria aureola Mesnil L.P. Mesnil det., 1970/
TYPE/ EX L.-P. MESNIL COLLECTION 1985.
Paratype: 1♂, Japan.

Solieriopsis borealis Reinhard, 1967, J. Kans. ent. Soc. 40: 100.
USA.

Holotype ♂: Umiat,Alaska 12 VII.1959 J.E.H.Martin/
HOLOTYPE *Solieriopsis borealis* Reinhard/
Solieriopsis borealis R Rnh.

Remarks: Holotype left front leg is glued to locality label.

Spallanzania colludens Reinhard, 1958b, J. Kans. ent. Soc. 31: 231.
USA.

Holotype ♂: College Station Oct.19.1935 Tex/ H. J Reinhard Collector/ HOLOTYPE *Spallanzania colludens* Reinhard/ *Spallanzania colludens* R Rnh.

Allotype ♀: Same data as holotype except with allotype label and dated "Oct.29,1933".

- Paratypes: 4♂♂, 2♀♀, Texas; 3♂♂, Arizona; 1♀, South Carolina.
- Spanipalpus conspicuum* Cortés, 1975, Boln tèc Estac. exp. agron. Fac. Agronomía Univ. Chile. 40: 6. CHILE.
Holotype ♂: ♂/ HOLOTYPE/ Pichinahuel, Cord. Nahuelbuta Arauco, CHILE 28-31.I -1959 L. Pena/ Spanipalpus ♂ conspicuum n.sp. R.Cortés det. 1974.
Paratype: 1♀, Chile.
- Spathidexia cerussata* Reinhard, 1934a, Bull. Brooklyn ent. Soc. 29: 152. USA.
Holotype ♂: Amherst, Ohio July, 1933/ H J Reinhard Collector/ HOLOTYPE Spathidexia cerussata Reinhard/ Spathidexia cerussata det.H.J.Reinh Rein.
- Spathidexia nexa* Reinhard, 1953a, Bull. Brooklyn ent. Soc. 48: 94. MEXICO.
Holotype ♀: Mexico City,D.F. W.G.Downs,Coll. 15-VII-51/ HOLOTYPE Spathidexia nexa Reinhard/ Spathidexia nexa R. Rnh.
- Spathidexia rasilis* Reinhard, 1934a, Bull. Brooklyn ent. Soc. 29: 153. USA.
Holotype ♀: Madison 5-31-31, Wis/ HOLOTYPE Spathidexia rasilis Reinhard/ Spathidexia rasilis R'43 Rnh/ Reinhard Collection.
Paratype: 1♀, Ohio.
Remarks: Holotype is missing abdomen.
- Spathimeigenia aurifrons* Curran, 1930b, Can. Ent. 62: 246. CANADA.
Holotype ♂: Holo TYPE Spathimeigenia aurifrons No. 3197. Curran/ Mont Laurier Que. Emer. 6.XII.1929. M.Dunn./ Host. Neodiprion Sp. Infesting P. banksiana/ 18/ Spathimeigenia TYPE aurifrons ♂ Curran. No.
Allotype ♀: Same data as holotype except with allotype label, and numbered label with "20" instead of "18".
Paratypes: 3♀♀, Quebec.
- Spathimeigenia dolopis* Reinhard, 1958a, Proc. ent. Soc. Wash. 60: 208. USA.
Holotype ♂: JacksonCoOre Griffin Creek VII.5 1951/ C.Fitch Collector/ HOLOTYPE Spathimeigenia dolopis Reinhard/ Spathimeigenia dolopis R Rnh.
- Spathimeigenia dolosa* Reinhard, 1958a, Proc. ent. Soc. Wash. 60: 209. USA.
Holotype ♂: College Station Jun.19.1946 Tex/ H J Reinhard Collector/ HOLOTYPE Spathimeigenia dolosa Reinhard/ Spathimeigenia dolosa R Rnh.
Allotype ♀: Same data as holotype except with allotype label and dated "Apr. 24, 1946".
Paratypes: 3♂♂, 3♀♀, Texas.
- Spathimeigenia erronis* Reinhard, 1958a, Proc. ent. Soc. Wash. 60: 212. USA.
Holotype ♂: College Station 6-3-20 Texas/ H J Reinhard Collector/ HOLOTYPE Spathimeigenia erronis Rnhd. CNC No./ Spathimeigenia erronis R Rnh.
- Spoggosia raoi* Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 182. INDIA.
Holotype ♂: Gooty Anantpur [Anantapur] 28. 8. 1963 Bangalore Inde/ Ex Amsacta albistriga/ TYPE/ Spoggosia raoi Mesnil L.P. Mesnil det., 1967/ EX L.-P. MESNIL COLLECTION 1970.
- Steiniomyia takanoi* Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24): 48. JAPAN.
Holotype ♀: Osaka Hiyakata M. Asuha [underside of label] 1957/ 229/ Steinomyia takanoi Mesnil L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Steleoneura novemmaculata* Wood, 1985, Mem. ent. Soc. Can. 132: 81. CANADA.
Holotype ♂: Victoria Is.,N.W.T 71° 17'N, 114° W 23-28.VII.1975 G. & M. Wood/ HOLOTYPE Steleoneura novemmaculata Wood CNC No. 18433/ HOLOTYPE Steleoneura novemmacula [sic] CNC No.
Allotype ♀: Same data as holotype except with allotype label and dated "8-10.VII.1973".
Paratypes: 16♂♂, Northwest Territories; 29♂♂, Yukon Territory.
- Stenocheata claripalpis* Thompson, 1968, Mem. ent. Soc. Can. 56: 159. TRINIDAD.
Holotype ♂: HOLOTYPE Stenocheata claripalpis Thomp. CNC No. 9877/ Legerville Mt. XII.12.60. Trinidad,BWI.
- Stenoneura serotina* Reinhard, 1945, Can. Ent. 77: 33. USA.
Holotype ♂: College Station Apr.21,1944 Tex/ H J Reinhard Collector/ HOLOTYPE Stenoneura serotina Reinhard/ Stenoneura serotina Rnh.
Allotype ♀: Same data as holotype except with allotype label and dated "Jun.27, 1944".
Paratypes: 2♂♂, 1♀, Texas.
Remarks: Date of collection of holotype was published as 21 April 1943.
- Stictodexia crosi* Villeneuve, 1920c, Annls Soc. ent. Belg. 60: 203. ALGERIA.
Holotype ♀: Mascara (Oran) 31.X.12/ Stictodexia Crosi ♀ Typ. Villen./ TYPE.
- Stictodexia lesnei* Villeneuve, 1912a, Bull. Mus. natn. Hist. nat., Paris 18: 416. ALGERIA.
Syntype: 1♂, 1♀.
♂: Edough le Morkat 500m. 3 oct.08 P. Lesne/ Stictodexia Lesnei Villen./ TYPE/ CNC Syntype Stictodexia lesnei Villeneuve Label affixed 1994.
♀: Ste Croix 4 oct.08 750m alt. P. Lesne/ CNC Syntype Stictodexia lesnei Villeneuve Label affixed 1994.
- Stolatosoma cidaris* Reinhard, 1953b, J. Kans. ent. Soc. 26: 97. MEXICO.
Holotype ♂: Acatlipa, Morelos May 11 '50 Coll.WGDowns/ HOLOTYPE Stolatosoma cidaris Reinhard/ Stolatosoma cidaris R. Rnh.

Allotype ♀: Acatlipa, Morelos June 2 1950
 Coll.WGDowns/ ALLOTYPE Stolatosoma cidaris
 Reinhard.
 Paratypes: 3♂♂, Mexico.

Stomatomya metallica Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 475. ZIMBABWE.

Syntype: 1♂.
 ♂: 634/ 00/31,31 marley 26.9.15/ Stomatomyia metallica Typ. Villen./ Chlorolydella metallica Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Stomatomya metallica Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Stomina caliendrata varians Villeneuve, 1930b, Bull. Annls Soc. ent. Belg. 70: 43. TUNISIA.

Syntype: 1♂.
 ♂: Tunis. P- 52787./ Stomina varians Villen. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Stomina caliendrata varians Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Stroblomyia jepsoni Villeneuve, 1937d, Bull. Mus. r. Hist. nat. Belg. 13(35): 2. MAURITIUS.

Holotype ♂: Ace, No. Date 31.12.34 LES MARES MAURITIUS W.F. Jepson/ Peribaea jepsoni Villen. L.P. Mesnil det., 1970/ Stroblomyia jepsoni Typ. Villen./ TYPE/ HOLOTYPE ♂ of Stroblomyia jepsoni Vill. Examined 1970 R.W.Crosskey.

Strongygaster nishijimai Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 73. JAPAN.

Holotype ♂: Sapporo, Maruyama, 18-VI.1941, Y.Nishijima/ Strongygaster nishijimai Mesn./ [underside of label] Host: [host name in Japanese letters].

Sturmia aurifera Villeneuve, 1943a, Bull. Soc. ent. Fr. 48: 36. ZAIRE.

Syntypes: 2♂♂.
 ♂: Congo-belge Eala-XII-1935 J.Ghesquière/ Sturmia aurifera n.sp./ TYPE/ Drino aurifera Villen. L.P. Mesnil det., 1969/ SYN-TYPE/ Syntype/ CNC Syntype Sturmia aurifera Villeneuve Label affixed 1994.
 ♂: Congo-belge Eala-XI-1935 J.Ghesquière/ Drino aurifera Villen. L.P. Mesnil det., 1969/ SYN-TYPE/ Syntype/ CNC Syntype Sturmia aurifera Villeneuve Label affixed 1994.

Sturmia crescentis Reinhard, 1944b, J. Kans. ent. Soc. 17: 69. USA.

Holotype ♂: College Station 5-1-29 Texas/ H J Reinhard Collector/ HOLOTYPE Sturmia crescentis Reinhard/ Sturmia crescentis det.H.J.Reinhard Rein./ Drino crescentis R (Rnh.).
 Allotype ♀: College Station 5-I-29 Texas/ H J Reinhard Collector/ ALLOTYPE Sturmia crescentis Reinhard.
 Paratypes: 34♂♂, 13♀♀, Texas.

Sturmia dasyops Villeneuve, 1943a, Bull. Soc. ent. Fr. 48: 40. NIGERIA.

Holotype ♂: Degema S. Nigeria Oct-Dec. 1910
 Dr.J.W.Collectt/ Sturmia dasyops Type n.sp./ TYPE/ Exorista dasyops Villen. L.P. Mesnil det., 1970/ EX L.-P. MESNIL COLLECTION 1970.

Sturmia fascipes Villeneuve, 1943a, Bull. Soc. ent. Fr. 48: 37. ZAIRE.

Holotype ♂: Sankisia 25.VIII.1911/ Sturmia fascipes ♂ n.sp./ Blepharella fascipes Villen. L.P. Mesnil det., 1969/ TYPE/ HOLOTYPE ♂ of Sturmia fascipes Villeneuve Examined 1970 R.W.Crosskey.

Sturmia laetabilis Villeneuve, 1933, Revue Zool. Bot. afr. 23: 279. SIERRA LEONE.

Syntype: 1♂.
 ♂: KONTA Sierra Leone, 13.V. 1912 Jas. J. Simpson/ Sturmia laetabilis i.litt. Villen./ Blepharella laetabilis Villen L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Sturmia laetabilis Villeneuve Label affixed 1994.

Sturmia mayneana Villeneuve, 1930d, Revue Zool. Bot. afr. 20: 59. ZAIRE.

Syntypes: 1♂, 1♀.
 ♂: Eala 14-XII-1917 R. Mayné/ Sturmia mayneana n.sp./ Drino mayneana Villen. L.P. Mesnil det., 1969/ TYPE/ SYN-TYPE/ Syntype/ CNC Syntype Sturmia mayneana Villeneuve Label affixed 1994.
 ♀: 23 Eala 5-II-1918 R. Mayné/ Drino mayneana Villen. L.P. Mesnil det., 1969/ SYN-TYPE/ Syntype/ CNC Syntype Sturmia mayneana Villeneuve Label affixed 1994.

Sturmia nigroscutellata Mesnil, 1970, Mushi 44: 91. PAPUA NEW GUINEA.

Holotype ♂: New Guinea Maprik Sepik 7. 1959/ Ex pup. Taenaris sp. (Amathusiid) pest of Banana/ Sturmia nigroscutellata Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: There may be a female paratype among the CNC material, but it is not labeled as such.

Sturmia rubricosa Villeneuve, 1933, Revue Zool. Bot. afr. 23: 279. MALAWI.

Syntype: 1♂.
 ♂: Mt. Mlanje, Nyasaland, 10.IX. 1913. S. A. Neave./ Sturmia rubricosa Villen./ Blepharella rubricosa Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype Sturmia rubricosa Villeneuve Label affixed 1994.

Sturmia versatilis Villeneuve, 1910b, Wien. ent. Ztg 29: 253. CONGO.

Holotype ♂: Typus/ Sturmia versatilis Villen./ Blepharella versatilis Villen. L.P. Mesnil det., 1969/ TYPE.

Sturmia (Blepharipoda) semitestacea Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 477. SOUTH AFRICA.

Syntype: 1♂.
 ♂: Kimberley Bro Power Nov.1913/ Sturmia (Blepharipoda) semitestacea Cotyp. Villen./

- Pimelomyia semitestacea* Vill. L.P. Mesnil det., 1969/ CNC Syntype *Sturmia* (*Blepharipoda*) *semitestacea* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Sturmiellina trinitatis* Thompson, 1963b, *Studia ent.* 6: 376. TRINIDAD.
Syntypes: 1♂, 1♀.
♂: HOLOTYPE *Sturmiellina trinitatis* Thomp. CNC No. 9795/ Melajo R. III.18.60
Trinidad,BWI/ *Sturmiellina* ng. *trinitatis* nsp./ CNC Syntype *Sturmiellina trinitatis* Thompson Label affixed 1994.
♀: ALLOTYPE *Sturmiellina trinitatis* Thomp. CNC No. 9795/ Brazil Trinidad,BWI. III.6.60./ CNC Syntype *Sturmiellina trinitatis* Thompson Label affixed 1994.
- Sturmiopsoidea obscura* Thompson, 1966, *Studia ent.* 8 (1965): 359. TRINIDAD.
Holotype ♂: HOLOTYPE *Sturmiopsoidea obscura* Thomp. CNC No. 9828/ American Rd. Mauvan Hill XII.30.60. Trinidad,BWI/ *Sturmiopsoidea obscura* ng nsp.
Paratype: 1♂, Trinidad.
- Subfischeria flavogrisea* Villeneuve, 1937a, *Revue Zool. Bot. afr.* 29: 211. SOUTH AFRICA.
Holotype ♀: Windsaxton Grigualand Dr.Brauns./ Cape Colony 20.XII. 1920/ *Subfischeria flavogrisea* Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Suensonomyia nudinervia* Mesnil, 1957, *Mém. Soc. r. ent. Belg.* 28: 15. JAPAN.
Holotype ♂: VII-5 1949 Kaminochi S.KATO/ 208/ *Suensonomyia nudinervia* Mesn. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
- Syneplaca ghesquierei* Villeneuve, 1938c, *Bull. Mus. r. Hist. nat. Belg.* 14(38): 13. ZAIRE.
Syntype: 1♂.
♂: Congo - belge Eala - IX -1936 J.Ghesquière/ *Syneplaca ghesquieri* [sic] Typ. Villen./ *Syneplaca ghesquierei* Vill. L.P. Mesnil det., 1970/ TYPE/ *Sumpigaster diaristata* Towns. L.P. Mesnil det., 1975/ CNC Syntype *Syneplaca ghesquierei* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Synorbitalia flavipes* Thompson, 1968, *Mem. ent. Soc. Can.* 56: 32. TRINIDAD.
Holotype ♀: HOLOTYPE *Synorbitalia flavipes* Thomp. CNC No. 9846/ "Simla" Trinidad/ 1358 1.17.64/ *Synorbitalia* gn. *flavipes* n.sp.
Paratype: 1♀, Trinidad.
- Tachina chaoi* Mesnil, 1966, *Fliegen palaearkt. Reg.* 64g: 910. JAPAN.
Syntype: 1♂.
♂: MUSEUM PARIS NIPPON MOYEN ENV. DE TOKIO ET ALPES DE NIKKO J. HARMAND 1901/ *Tachina chaoi* Mesnil L.P. Mesnil det., 1970/ TYPE/ CNC Syntype *Tachina chaoi* Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Tachina duplaria* Villeneuve, 1916c, *Ann. S. Afr. Mus.* 15: 493. MALAWI.
Syntype: 1♂.
♂: Nyasaland Mlanje W.H. Brown/ TYPE/ *duplaria* Typ. Villen./ *Exorista duplaria* Villen. L.P. Mesnil det., 1970/ Syntype/ CNC Syntype *Tachina duplaria* Villeneuve Label affixed 1994.
- Tachina ebneri* Villeneuve, 1922b, *Revue zool. afr.* 10: 62. SUDAN.
Lectotype ♂: Kondofan Werner. eb.VIII.05 Ex. *Auchmophila Kordofensis* Rh./ *Tachina Ebneri* Typ. Villen./ TYPE/ LECTO-TYPE/ *Exorista ebneri* Villen. L.P. Mesnil det., 1970/ *Tachina ebneri* Vill. Mesnil fixation as LECTOTYPE/ CNC Syntype *Tachina ebneri* Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1960: 589. Label affixed 1995.
Paralectotype: 1♀, Sudan.
Remarks: We regard Mesnil's (1960a: 589) reference to a holotype as a lectotype fixation for this species.
- Tachina fallax* var. *pseudofallax* Villeneuve, 1920a, *Revue zool. afr.* 8: 151. SOUTH AFRICA.
Syntypes: 2♂♂.
♂: Capland Willowmore 12 1911 Dr. Brauns/ *Exorista pseudofallax* Villen. L.P. Mesnil det., 1970/ TYPE/ Syntype/ CNC Syntype *Tachina fallax* var. *pseudofallax* Villeneuve Label affixed 1994.
♂: Capland Willowmore 12 1911 Dr. Brauns/ *Tachina fallax* var. a yeux valus/ *Exorista pseudofallax* Villen. L.P. Mesnil det., 1970/ CNC Syntype *Tachina fallax* var. *pseudofallax* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- Tachinomima multisetosa* var. *fasciata* Villeneuve, 1936a, *Bull. Mus. r. Hist. nat. Belg.* 12(4): 8. UGANDA.
Holotype ♀: Afrique or. allemande KILIMANDJARO VERSAN SUD-EST ALLUAUD & JEANNEL/ PRAIRIES ALPINES BISMARCK-HUGEL 2600-2800m Mars-Avril 1912 St. 70/ *Tachinomima multisetosa* var. *albopilosa* Typ. ♀ Villen./ *Linnaemyia fasciata* Villen. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.
Remarks: Villeneuve apparently mislabeled the holotype as "var. *albopilosa*" instead of "var. *fasciata*". He did not describe a "var. *albopilosa*".
- Tachinomyia apicata* Curran, 1926a, *Trans. R. Soc. Can., Ser. 3,* 20: 171. CANADA.
Holotype ♂: TYPE *Tachinomyia apicata* Curran No. 1867/ Aylmer, Que. 24.V.1924 C.H.Curran.
Allotype ♀: Mounted on same pin as holotype.
Paratypes: 1♂, 1♀, Quebec; 1♂, 1♀, Alberta; 1♀, British Columbia; 1♀, Nova Scotia.
Remarks: We assume that Curran intended the male to be the holotype, as he consistently chose males as holotypes throughout his 1926 paper.

Tachinomyia occidentalis Curran, 1926a, Trans. R. Soc. Can., Ser. 3, 20: 170. CANADA.

Holotype ♂: Victoria,B.C. 13.V.1917 A.E.Cameron/ HoloTYPE Tachinomyia occidentalis Curran No. 1866.

Allotype ♀: Victoria,B.C. 2.VI 1923 K.F.Auden/ AlloTYPE Tachinomyia occidentalis Curran No. 1866.

Paratypes: 7♂♂, British Columbia; 2♂♂, Manitoba.

Tachinomyia variata Curran, 1926a, Trans. R. Soc. Can., Ser. 3, 20: 169. CANADA.

Holotype ♂: HoloTYPE Tachinomyia variata Curran No. 1865/ Pt.Pelee,Ont. 28.V.1925 G.S.Walley.

Allotype ♀: AlloTYPE Tachinomyia variata Curran No. 1865/ Onah, Man. 25 V 1922 H.A.Robertson.

Paratypes: 8♂♂, Ontario; 3♂♂, 2♀♀, Manitoba; 7♂♂, 4♀♀, Alberta; 2♂♂, Kansas.

Tachinophytopsis ghanii Mesnil, 1975a, Stuttg. Beitr. Naturk. (A) 277: 1. PAKISTAN.

Holotype ♂: Kahuta 7. VIII. 74/ Ex. Chrysomelid larva feeding on leaves of *Loranthus longiflorus* 1399/ C.I.B.C Lor. 8/74.25/ TYPE/ Tachinophytopsis ghanii Mesnil L.P. Mesnil det., 1975/ EX L.-P. MESNIL COLLECTION 1985.

Takanomyia scutellata Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 10. JAPAN.

Holotype ♀: XI-1.1949 Manazuru N.FUKUHARA/ 185/ Takanomyia scutellata Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Date of collection of holotype was published as 1 September 1949.

Talparomyia pollinosa Thompson, 1968, Mem. ent. Soc. Can. 56: 101. TRINIDAD.

Holotype ♀: HOLOTYPE Talparomyia pollinosa Thomp. CNC No. 9864/ SANTA CRUZ Trinidad,BWI. JULY. 15. 1954/ Collector F.J.Simmonds/ 1403 II.21.64/ 73/ 36b.

Tamanamyia trinitatis Thompson, 1963a, Can. J. Zool. 41: 466. TRINIDAD.

Holotype ♀: HOLOTYPE Tamanamyia trinitatis Thomp. CNC No. 9782/ Trinidad,BWI. Plum Mitan XII.22.60/ Biche/ HOLOTYPE CNCNo./ 1199 IX.14.61/ Tamanamyia trinitatis g.n. sp.n./ Stenoneura trinitatis Thomp.

Tarassophorus evexus Reinhard, 1964a, J. Kans. ent. Soc. 37: 39. USA.

Holotype ♀: Amherst Ohio Jul-Aug 1953 H.J.Reinhard/ HOLOTYPE Tarassophorus evexus Reinhard/ Tarassophorus evexus R Rnh.

Allotype ♂: War Bonnet Canyon Neb/ ALLOTYPE Tarassophorus evexus Reinhard.

Paratype: 1♀, Ohio.

Tarpessita fulgens Reinhard, 1967, J. Kans. ent. Soc. 40: 96. COLOMBIA.

Holotype ♂: Funza, Cund 5-VIII-54/ M. Revelo Maiz/ 3978/ HOLOTYPE Tarpessita fulgens Reinhard/ Tarpessita fulgens R Rnh.

Technamyia cinereola Reinhard, 1974, Can. Ent. 106: 1169. MEXICO.

Holotype ♀: 25Mi.W.LaPaz,BAJA CALIF.VIII-30-1959 KW Radford&FG Werner/ HOLOTYPE Technamyia cinereola Reinhard/ Technamyia cinereola R Rnh.

Telemus trossulus Reinhard, 1967, J. Kans. ent. Soc. 40: 106. USA.

Holotype ♀: Oakhurst, Cal. Madera Co. V-26 1942/ Arthur J Walz Collector/ HOLOTYPE Telemus trossulus Reinhard/ Telemus trossulus R Rnh/ Myatelemus trossulus R (Rnh).

Allotype ♂: Big Bend Nat. Pk. TEX. Chisos Basin Apr. 29 1959 Howden & Becker/ ALLOTYPE Telemus trossulus Reinhard/ IDEMA ILLUSTRATION.

Paratype: 1♀, Arizona; 1♀, Texas.

Tesseracephalus lenis Reinhard, 1955a, J. Kans. ent. Soc. 28: 124. MEXICO.

Holotype ♂: Acatlipa, Mor. VII/14/50 Coll. WGDowns/ HOLOTYPE Tesseracephalus lenis Reinhard/ Tesseracephalus lenis Rnh. Reinhard Coll'n/ Tesseracephalus lenis R Rnh.

Allotype ♀: Same data as holotype except with allotype label and without last label.

Paratype: 1♂, Mexico.

Thecocarcelia latifrons Mesnil, 1949a, Fliegen palaearkt. Reg. 64g: 56. MOZAMBIQUE.

Holotype ♂: MUSEUM PARIS ZAMBEZE NOVA CHOUPANGA PRÉS CHEMBA P. LESNE 1929/ JUIN/ T. latifrons Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Thecocarcelia robusta Mesnil, 1950a, Revue Zool. Bot. afr. 43: 22. ZAIRE.

Syntype: 1♂.

♂: Congo - belge Eala-12.IV - 1935 J.Ghesquière/ R. Mus. Hist. Nat. Belg. 1 0 4 8 2./ Cotype/ T. robusta Mesnil L.P. Mesnil det., 1969/ CNC Syntype Thecocarcelia robusta Mesnil Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Thecocarcelia trichops Herting, 1967, Stuttg. Beitr. Naturk. 173: 4. FRANCE.

Holotype ♂: Vaucluse Lagnes 13.7.65/ Thecocarcelia trichops Hert. L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Thelaira americana Brooks, 1945b, Can. Ent. 77: 88. CANADA.

Holotype ♂: Aylmer,Que. 24.VI.1924 C.H.Curran/ HOLOTYPE ♂ Thelaira americana No. Brks 5527.

Allotype ♀: Same data as holotype except with allotype label and dated "27. VI. 1924".

Paratypes: 1♀, Nova Scotia; 2♂♂, 2♀♀, Quebec; 5♂♂, 2♀♀, Ontario; 1♀, British Columbia; 3♂♂, 4♀♀, Massachusetts; 8♂♂, Ohio; 12♂♂, Texas.

Thelaira bryanti Curran, 1925a, Can. Ent. 57: 281. CANADA.

Holotype ♂: HOLOTYPE ♂ Thelaira bryanti No. Curran 1428/ Slave Lake/ AUG 17.'24.

Allotype ♀: Same data as holotype except with allotype label.

Thelaira ghanii Mesnil, 1968a, Bull. Annls Soc. r. ent. Belg. 104: 186. PAKISTAN.

Holotype ♂: Ex.larvae in web of *Prunus grunta*/ Murree 29-XII-61/ *Thelaira ghanii* Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Holotype puparium is pinned below specimen. Host plant was published as *Prunus cornuta*. There may be a paratype among the CNC material, but it is not labeled as such.

Thelairodoriopsis maracasi Thompson, 1968, Mem. ent. Soc. Can. 56: 54. TRINIDAD.

Holotype ♂: HOLOTYPE *Thelairodoriopsis maracasi* Thomp. CNC No. 9851/ Simla TRINIDAD W.I. III.16.61 W.R.Thompson/ 1365 1.24.64.

Allotype ♀: ST. JOSEPH Trinidad,BWI. NOV.10.1952./ Collector F.J.Simmonds/ ALLOTYPE *Thelairodoriopsis maracasi* Thomp. CNC No. 9851/ HOMOTYPE *Phorocera sobrina* Wulp Compared by Wood '83.

Remarks: Date of collection of holotype was published as 19 March 1961. Holotype is missing terminalia.

Thelairosoma fumosum Villeneuve, 1916c, Ann. S. Afr. Mus. 15: 500. MALAWI.

Syntype: 1♀.

♀: Mt. Mlanje, Nyasaland, 10.IX.1913 S. A. Neave./ *Thelairosoma fumosum* Typ. Villen./ *Thelairosoma fumosum* Vill. L.P. Mesnil det., 1969/ TYPE/ CNC Syntype *Thelairosoma fumosum* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Thelairosoma (Lespesiopsis) coerulescens Mesnil, 1954c, Annls Mus. r. Congo belge Sér. 4to (Zool.) 1: 471. TANZANIA.

Holotype ♂: Urw.hint.d.Randbg. d.N.W.Tanganika-S. 18-2200m. Graner/ Lespesiopsis coerulescens Mesn. L. Mesnil det., 1943/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Thelairosoma (Seyrigomyia) ingrami Mesnil, 1970, Mushi 44: 103. UGANDA.

Holotype ♂: 1619 SERERE Obulai 13.Dec.1958 W.R.Ingram ex. *Papilio pylades*/ *Thelairosoma ingrami* Mesnil L.P. Mesnil det., 1969/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Holotype puparium is pinned below specimen.

Thelyconychia (Thelyconychia) discalis Mesnil, 1957, Mém. Soc. r. ent. Belg. 28: 5. JAPAN.

Holotype ♂: Obihiro S.Takano [underside of label] 10/VI '49/ 164/ *Thelyconychia discalis* Mesnil L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Thelymyia disparis Reinhard, 1959a, Ent. News 70: 230. USA.

Holotype ♂: W.sl.Patagonia Mts Sta.Cruz Co. ARIZ Aug 9 1955 GDButler &FGWerner/ HOLOTYPE *Thelymyia dispar* [sic] Reinhard.

Paratypes: 2♂♂, Arizona.

Thematheca medeola Reinhard, 1961, J. Kans. ent. Soc. 34: 208. MEXICO.

Holotype ♀: Rio Tehuantepec, Oax. Sept. 6, 1947/ Oax./ F. A. Cowan M. R. Wheeler Collectors/ HOLOTYPE *Thematheca medeola* Reinhard/ *Thematheca medeola* R Rnh.

Paratypes: 2♀♀, Mexico; 1♀, Arizona.

Thyellina brevicornis Mesnil, 1949a, Fliegen palaearkt. Reg. 64g: 70. AUSTRALIA.

Holotype ♀: Herbezon Dodd XII. 1910 3700 Ft./coll. Liebtwardt/ *Thyellina brevicornis* Mesn. L.P. Mesnil det., 1967/ TYPE/ HOLOTYPE ♀ of *Thyellina brevicornis* Mesnil Examined 1970. R.W.Crosskey.

Trafoia gemina Herting, 1966, Stuttg. Beitr. Naturk. 146: 6. AUSTRIA.

Holotype ♂: Hauser Kaibling 23.6.61/ *Trafoia gemina* Herting L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Trichoparia (Admontia) cepelaki Mesnil, 1961b, Fliegen palaearkt. Reg. 64g: 674. SWITZERLAND.

Holotype ♂: Bernina Aout. 04/ *Trichoparia cepelaki* Mesnil L.P. Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Remarks: Date of collection of holotype was published as August 1914.

Trichosaundersia callithrix Reinhard, 1974, Can. Ent. 106: 1169. MEXICO.

Holotype ♂: Mex.City 10,000 ft. July-1936 H.G.Meyer/ HOLOTYPE *Trichosaundersia callithrix* Rnh. CNC No.

Triodontopyga trinitatis Thompson, 1963c, Can. Ent. 95: 992. TRINIDAD.

Holotype ♂: HOLOTYPE *Triodontopyga trinitatis* Thomp. CNC No. 9805/ I.C.T.A. Trinidad,BWI. APRIL.5.1954/ Collector F.J.Simmonds.

Allotype ♀: PIARCO Trinidad,BWI. OCT.29.1953/ Collector F.J.Simmonds/ ALLOTYPE

Triodontopyga trinitatis Thomp. CNC No. 9805.

Paratypes: 3♂♂, 4♀♀, Trinidad.

Trismegistus pumilis Reinhard, 1967, J. Kans. ent. Soc. 40: 101. USA.

Holotype ♀: Portal,Ariz. VIII-17-65/ H J Reinhard Collector/ HOLOTYPE *Trismegistus pumilis* Reinhard/ *Trismegistomyia pumulis* R (Rnh.).

Trixia pyrenaica Villeneuve, 1928, Bull. Annls Soc. ent. Belg. 68: 50. FRANCE.

Lectotype ♂: Lac de Caderoles H.G. VIII/ *Trixia pyrenaica* Vill. L.P. Mesnil det., 1978/ TYPE/ CNC Syntype *Trixia pyrenaica* Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1980: 14. Label affixed 1995/ EX L.-P. MESNIL COLLECTION 1985.

Paralectotype: 1♀, France.

Remarks: We regard Mesnil's (1980: 14) reference to a holotype as a lectotype fixation for this species.

Trochiloglossa aurea Thompson, 1963a, Can. J. Zool. 41: 350.
TRINIDAD.

Holotype ♂: HOLOTYPE *Trochiloglossa aurea* Thomp. CNC No. 9765/ St. Augustine, Trinidad, BWI. III.8.60/ P.M.Estate U.S.M D.W.FEWKES/ Emerged 12/1/60 Parasite of ?*Opogona* sp./ *Trochiloglossa aurea* n.sp.

Trypheromyia pallens Reinhard, 1945, Can. Ent. 77: 32. USA.
Holotype ♀: Amherst, Ohio July, 1933/ H J Reinhard Collector/ HOLOTYPE *Trypheromyia pallens* Reinhard/ *Trypheromyia pallens* R Rnh.
Paratype: 1♀, Iowa.

Uclesia varicornis Curran, 1927b, Can. Ent. 59: 300. USA.
Holotype ♀: Beach ND/ CNAinslie Collector/ *Uclesia* TYPE ♀ *varicornis* Curran. No. 2456/ *Uclesia* *varicornis* Det. Curran. C.H.Curran.

Uclesia zonalis Curran, 1927b, Can. Ent. 59: 299. CANADA.
Holotype ♀: *Uclesia* TYPE ♀ *zonalis* Curran No. 2455/ Oliver, B.C. 10.V 1923 E.R.Buckell/ *Uclesia* *zonalis* Det. Curran C.H.Curran.

Uramya hariola Reinhard, 1961, J. Kans. ent. Soc. 34: 206.
COLOMBIA.
Holotype ♂: Colombia S.A./ HOLOTYPE *Uramya* *hariola* Reinhard/ *Uramya hariola* R Rnh.

Uromacquartia trinitatis Thompson, 1963a, Can. J. Zool. 41: 352.
TRINIDAD.
Syntype: 1♂.
♂: HOLOTYPE *Uromacquartia trinitatis* Thomp. CNC No. 9766/ Mt. Lambert TRINIDAD, W.I. IX.21.60 W.R.Thompson/ Type ♂/ CNC Syntype *Uromacquartia trinitatis* Thompson Label affixed 1994.

Vibrissotheresia pechumani Reinhard, 1943c, Bull. Brooklyn ent. Soc. 38: 87. USA.
Holotype ♂: Brewster, N.Y. 2 July, 1937 L. L. Pechuman/ HOLOTYPE *Vibrissotheresia* *pechumani* Reinhard/ *Vibrissotheresia* *pechumani* det.H.J.Reinh Rein.

Viviania pedita Reinhard, 1959b, Pan-Pacif. Ent. 35: 158. USA.
Holotype ♂: Lake Placid Fla. March 3 1945 J.G.Needham/ HOLOTYPE *Viviania pedita* Reinhard/ *Sitophaga aurigera* R (Coq.).

Vivianoidea grisea Thompson, 1968, Mem. ent. Soc. Can. 56: 37.
TRINIDAD.
Holotype ♂: HOLOTYPE *Vivianoidea grisea* Thomp. CNC No. 9847/ Simla 1964 Trinidad, WI.
Paratypes: 2♂♂, Trinidad.
Remarks: Holotype is missing terminalia.

Voria pilibasis Villeneuve, 1922c, Annls Sci. nat. Zool. (10)5: 339.
YUGOSLAVIA.
Lectotype ♀: Alban.Exp.1918 Prizren 16.V./ *Voria* *pilibasis* ♀ Villen. Typ./ *Hypovoria pilibasis*

Villen. L.P. Mesnil det., 1970/ TYPE/ CNC
Syntype *Voria pilibasis* Villeneuve Label affixed 1994/ Lectotype fixation by Mesnil, 1974: 1280.
Label affixed 1995.

Remarks: We regard Mesnil's (1974: 1280) reference to a holotype as a lectotype fixation for this species.

Voria (Chaetovoria) antennata Villeneuve, 1920b, Annls Soc. ent. Belg. 60: 118. FRANCE.

Holotype ♂: type/ Lautaret (H.A) 28.VII.08/ *Voria* (*Chaetovoria*) *antennata* Typ. Villen./ *Uclesia* *antennata* Vill. L.P. Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Wagneria cornuta Curran, 1928a, Can. Ent. 60: 49. USA.

Holotype ♂: *Wagneria* TYPE ♂ *cornuta* Curran. No. 2669/ MelrsHghds 7/6/09 Mass/ WRThompson Collector.

Allotype ♀: *Wagneria* AlloTYPE *cornuta* ♀ Curran No. 2669/ Barber D N.B. 25-VI-1914 F. M.McKenzie.
Paratype: 1♀, New Brunswick.

Remarks: Holotype right front leg is glued to type label.

Wagneria misella Villeneuve, 1937c, Bull. Mus. r. Hist. nat. Belg. 13(34): 13. CHINA.

Holotype ♀: MtOmei SzechuenChina/ DCGraham Coll/ *Wagneria misella* Typ. Villen./ *Periscepsia misella* Villen. L.P. Mesnil det., 1970/ TYPE/
HOLOTYPE ♀ of *Wagneria misella* Villeneuve Examined 1970 R.W.Crosskey.

Wagneria ocellaris Reinhard, 1955b, J. Kans. ent. Soc. 28: 55.
USA.

Holotype ♂: Cayuse, Oregon May 13, 1938 Coll. K. Grey J. Schuh/ HOLOTYPE *Wagneria ocellaris* Reinhard/ *Wagneria ocellaris* det.H.J.Rein Rnh.
Allotype ♀: Monrovia Canyon Calif. Mar.27,1932/
ALLOTYPE *Wagneria ocellaris* Reinhard.
Paratypes: 5♂♂, California; 1♂, Oregon.

Wagneria pacata Reinhard, 1955b, J. Kans. ent. Soc. 28: 56. USA.

Holotype ♂: Amherst, Ohio 7-14 1934/ H J Reinhard Collector/ HOLOTYPE *Wagneria pacata* Reinhard/ *Wagneria pacata* Rnh det.H.J.Reinhard.
Allotype ♀: Amherst, Ohio 6-19 1939/ A J Barckert Coll./ ALLOTYPE *Wagneria pacata* Reinhard.
Paratypes: 1♂, Ohio; 1♂, Colorado; 1♀, Alberta.

Wagneria umbrinervis Villeneuve, 1937c, Bull. Mus. r. Hist. nat. Belg. 13(34): 13. CHINA.

Syntypes: 2♂♂.
♂: of Fundart coll. hugmayer/ *Wagneria* *umbrinervis* Typ. Villen./ *Periscepsia umbrinervis* Vill. L.P. Mesnil det., 1975/ TYPE/ CNC Syntype *Wagneria umbrinervis* Villeneuve Label affixed 1994.
♂: West tibet hugmayer/ *Wagneria umbrinervis* Villen./ *Periscepsia umbrinervis* Vill. L.P. Mesnil det., 1975/ CNC Syntype *Wagneria umbrinervis* Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Wagneria (Phorichaeta) alpina Villeneuve, 1910a, Wien. ent. Ztg 29: 87. FRANCE.

Syntypes: 10♂♂.

- ♂: Lautaret (H.A.) VII/ Wagneria alpina Type Villen./ Aphelogaster alpina Villen. L.P. Mesnil det., 1970/ TYPE/ CNC Syntype Wagneria (Phorichaeta) alpina Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- ♂: Lautaret (H.A.) VII/ ♂/ Aphelogaster alpina Villen. L.P. Mesnil det., 1970/ CNC Syntype Wagneria (Phorichaeta) alpina Villeneuve Label affixed 1994.
- 7♂♂: Lautaret (H.A.) VII/ Aphelogaster alpina Villen. L.P. Mesnil det., 1970/ CNC Syntype Wagneria (Phorichaeta) alpina Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.
- ♂: Lautaret (H.A.) 7/ Aphelogaster alpina Villen. L.P. Mesnil det., 1970/ CNC Syntype Wagneria (Phorichaeta) alpina Villeneuve Label affixed 1994/ EX L.-P. MESNIL COLLECTION 1970.

Wagneria (Wagneria) micronychia Mesnil, 1974, Fliegen palaearkt. Reg. 64g: 1289. SPAIN.

Holotype ♂: Sierra Palacio Spain 13. 4. 1924/ Wagneria micronychia Mesnil. L.P. Mesnil det., 1973/ TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Wardarina melancholica Mesnil, 1956a, Fliegen palaearkt. Reg. 64g: 481. YUGOSLAVIA.

Holotype ♂: Skopje Jugoslavien Dr.R.Meyer V.1937/ Wardarina melancholica Mesnil. L.P. Mesnil det., 1966/ TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Weberia (Chaetoweberia) rubiginans Villeneuve, 1932b, Bull. Soc. ent. Fr. 37: 270. TAIWAN.

Holotype ♀: Formosa, Sauter/ Kosempo, 1908 VI/ Chaetoweberia rubiginans Type Villen./ Holotype of Ch. rubiginans Vill. Examined 1975, R.W. Crosskey.

Winthemia analisella Thompson, 1963c, Can. Ent. 95: 978. TRINIDAD.

Holotype ♂: HOLOTYPE Winthemia analisella Thomp. CNC No. 9801/ ORANGE GROVE Trinidad,BWI. OCT-24-1952/ Collector F.J.Simmonds/ 1238D IX.7.62.

Paratypes: 3♂♂, Trinidad; 1♂, Brazil.

Remarks: Holotype is missing terminalia.

Winthemia andersoni Guimarães, 1972, Archos Zool. S Paulo 22: 50. USA.

Holotype ♂: Samuel Spr. Napa Co. Cal. IV.28 1964/ R.M.Bohart Collector/ Reinhard Collection/ HOLOTYPE Winthemia andersoni Guimaraes/ 14/ Winthemia andersoni Guimaraes det.Guimaraes 1969.

Paratype: 1♂, Arizona; 1♂, 1♀, California.

Remarks: Date of collection of holotype was published as 8 July 1955.

Winthemia aureonigra Thompson, 1963c, Can. Ent. 95: 978. TRINIDAD.

Holotype ♂: HOLOTYPE Winthemia aureonigra

Thomp. CNC No. 9802/ MARACAS V.

Trinidad,BWI. NOV.17.1952./ Collector

F.J.Simmonds/ 1292 III.15.63.

Allotype ♀: 2.17.60 Brazil Trinidad,BWI./ 1239A

IX.10.62/ ALLOTYPE Winthemia aureonigra

Thomp. CNC No. 9802.

Paratypes: 4♀♀, Trinidad.

Remarks: Holotype is missing terminalia.

Winthemia caledoniae Mesnil, 1969, Entomophaga 13: 203. NEW CALEDONIA.

Holotype ♂: ex chenille de Othreis fullonica L. Noumea (N.Cal.) 12.X.1964./ TYPE/ Winthemia caledoniae n.sp. L.P. Mesnil det., 1965/ EX L.-P. MESNIL COLLECTION 1970/ Holotype ♂ of Winthemia caledoniae Mesnil Det. B.K. Cantrell, 1987.

Remarks: Holotype puparium is glued to paper point pinned below specimen.

Winthemia communis Thompson, 1963c, Can. Ent. 95: 982. TRINIDAD.

Holotype ♀: HOLOTYPE Winthemia communis Thomp. CNC No. 9803/ FLORADELLE. Trinidad,BWI. OCT.7.1953./ Collector F.J.Simmonds.

Paratypes: 8♀♀, Trinidad.

Winthemia dubiosa Thompson, 1963c, Can. Ent. 95: 968. TRINIDAD.

Holotype ♂: HOLOTYPE Winthemia dubiosa Thomp. CNC No. 9799/ MARACAS V. Trinidad,BWI. MARCH 3. 1953./ Collector F.J.Simmonds/ 1238C IX.5.62/ Winthemia dubiosa n.sp.

Remarks: Holotype is missing terminalia, and left hind leg is glued to locality label.

Winthemia floridensis Guimarães, 1972, Archos Zool. S Paulo 22: 66. USA.

Holotype ♂: Tampa VII-20.34 Fla/ Reinhard Collection/ HOLOTYPE Winthemia floridensis/ Winthemia floridensis Guim. J.H.Guimarães det.

Remarks: Holotype terminalia are stored in glycerine in a plastic microvial pinned below specimen.

Winthemia fumiferanae Tothill, 1912, Can. Ent. 44: 2. CANADA.

Holotype ♀: Emerg. 15.VII.11 from. sp. [spruce] b. [bud] worm Loc. Duncan B.C. Coll./ TYPE No. 491/ Winthemia fumiferanae n.sp. J.D.T. Type.

Paratypes: 1♂, 1♀, Quebec; 4♂♂, 11♀♀, British Columbia.

Winthemia orbitalis Villeneuve, 1934b, Revue Zool. Bot. afr. 26: 68. ZAIRE.

Holotype ♂: Moho Lesse 9.VI.14/ Winthemia Dr Villeneuve det. orbitalis ♂ n.sp. Villen./ TYPE/ Blepharella erronea Mesnil. L.P. Mesnil det., 1969/ HOLOTYPE ♂ of Winthemia orbitalis Villeneuve Examined 1970 R.W.Crosskey.

Winthemia papuana Mesnil, 1969, Entomophaga 13: 204. PAPUA NEW GUINEA.

Holotype ♂: Aiyura Montagnes Est Nelle Guinée/ Prépupe de Saturnide sp. 22.2.1958/ Larve sortie

le 5.3.1958, adulte éclos 21.3.1958/ TYPE/
Winthemia papuana Mesnil L.P. Mesnil det.,
 1967/ Holotype ♂ of *Winthemia papuana* Mesnil
 det. B.K. Cantrell, 1987/ EX L.-P. MESNIL
 COLLECTION 1970.

Winthemia pollinosa Thompson, 1963c, Can. Ent. 95: 984.
 TRINIDAD.
 Holotype ♀: HOLOTYPE *Winthemia pollinosa* Thomp.
 CNC No. 9804/ Plum Mitan Rd. XII.20.60.
 Trinidad,BWI/ *Winthemia pollinosa* n.sp.

Winthemia trinitatis Thompson, 1963c, Can. Ent. 95: 971.
 TRINIDAD.
 Holotype ♂: HOLOTYPE *Winthemia trinitatis* Thomp.
 CNC No. 9800/ CHAGUANAS. Trinidad,BWI.
 OCT.30.1952/ Collector F.J.Simmonds.
 Allotype ♀: St. Augustine, Trinidad,BWI. JAN 1960/
 1029 I.12.60/ ALLOTYPE *Winthemia trinitatis*
 Thomp. CNC No. 9800.
 Paratypes: 9♂♂, 1♀, Trinidad.

Winthemia venustoides Mesnil, 1967, Mushi 41: 39. JAPAN.
 Holotype ♂: Tsukisappu nr/Sapporo 19.VIII.1961 Japon
 Hokkaido/ *Winthemia venustoides* Mesn. L.P.
 Mesnil det., 1964/ TYPE/ EX L.-P. MESNIL
 COLLECTION 1970.

Winthemia (Dorbinella) jacentkowskyi Mesnil, 1949a, Fliegen
 palaearkt. Reg. 64g: 85. CZECH REPUBLIC.
 Holotype ♂: Obrany 4.VI.42 19/ *Winthemia*
jacentkowskii [sic] Mes. L.P. Mesnil det., 1964/
 TYPE/ EX L.-P. MESNIL COLLECTION 1970.

Winthemia (Winthemia) albidiopilosa Mesnil, 1949a, Fliegen
 palaearkt. Reg. 64g: 83. INDONESIA.
 Holotype ♀: Sunda Exp. Rensch Ende Flores
 13.II.1927/ *Winthemia albidiopilosa* Mesn. L.P.
 Mesnil det., 1949/ TYPE/ HOLOTYPE ♀ of
Winthemia albidiopilosa [sic] Mesnil Examined
 1970 R.W.Crosskey.

Winthemia (Winthemia) speciosissima Mesnil, 1949a, Fliegen
 palaearkt. Reg. 64g: 95. FRANCE.
 Holotype ♂: Bois St Martin Versailles 27 Aout 1936/
Winthemia speciosissima Mesn. L.P. Mesnil det.,
 1944/ TYPE/ EX L.-P. MESNIL COLLECTION
 1970.

Xanthocera lucentis Reinhard, 1962a, Ent. News 73: 174. USA.
 Holotype ♀: Cranberry L.,N.Y. 10July 1951
 L.L.Pechuman/ HOLOTYPE *Xanthocera lucentis*
 Reinhard/ *Xanthocera lucentis* R Rnh./ *Xanthocera*
 atra R Tns.

Xanthomelanodes trivittatus Reinhard, 1955d, Ent. News 66: 233.
 MEXICO.
 Holotype ♂: 15 km.N.of Chapalilla, Nay.,Mex. 19 July
 1951/ Baccharis/ P. D. Hurd Collector/
 HOLOTYPE *Xanthomelanodes trivittata*
 Reinhard/ *Xanthomelanodes trivittata* R. Rnh.

Xanthooestrus fastuosus Villeneuve, 1914, Annls hist.-nat. Mus.
 natn. hung. 12: 440. TAIWAN.

Lectotype ♂: Formosa Toyenmongai/ *Xanthooestrus*
fastuosus ♂ Villen./ EX L.-P. MESNIL
 COLLECTION 1970/ CNC Lectotype
Xanthooestrus fastuosus Villeneuve Crosskey
 designation 1976 Label affixed 1994.

Remarks: For lectotype designation see Crosskey
 (1976: 278).

Xenolophosia diversipes Villeneuve, 1926c, Bull. Annls Soc. ent.
 Belg. 66: 275. TAIWAN.
 Holotype ♀: Formosa, Sauter/ Daitorinsho, 1909 VII./
Xenolophosia diversipes Type ♀, Villen./
Lophosia excisa Tothill, det. R.W. Crosskey 1975.

Xenolophosia hamulata Villeneuve, 1926c, Bull. Annls Soc. ent.
 Belg. 66: 274. TAIWAN.
 Holotype ♂: Formosa, Sauter/ Taihorin, 1911 VII./
Xenolophosia hamulata Type ♂, Villen./ *Lophosia*
hamulata Vill., det. R.W. Crosskey 1975.

Xiphomyia rivalis Reinhard, 1953b, J. Kans. ent. Soc. 26: 101.
 MEXICO.
 Holotype ♀: Acatlipa, Mor. VII/14/50
 Coll.W.G.Downs/ HOLOTYPE *Xiphomyia rivalis*
 Reinhard.

Xystomima umbrinervis Villeneuve, 1925, Konowia 4: 50. SOUTH
 AFRICA.

Syntypes: 2♀♀.
 ♀: Durban Marley 18.7.16/ *Xystomima*
umbrinervis n.sp. Villen./ Therobia umbrinervis
 Vill. L.P. Mesnil det., 1970/ TYPE/ Syntype/ CNC
 Syntype *Xystomima umbrinervis* Villeneuve Label
 affixed 1994.
 ♀: Marley 4 1915 Durban/ Therobia umbrinervis
 Villen L.P. Mesnil det., 1970/ CNC Syntype
Xystomima umbrinervis Villeneuve Label affixed
 1994/ EX L.-P. MESNIL COLLECTION 1970.

Zambesa claripalpis Villeneuve, 1926c, Bull. Annls Soc. ent. Belg.
 66: 272. TAIWAN.

Syntype: 2♀♀.
 ♀: Formosa Sauter/ Koshun 909.III./ *Zambesa*
claripalpis Typ. Villen./ *Zambesa claripalpis* Vill.
 L.P. Mesnil det., 1970/ TYPE/ *Zambesa*
formosensis T.T. L.P. Mesnil det., 1970/ EX
 2L.-P. MESNIL COLLECTION 1970/ CNC
 Syntype *Zambesa claripalpis* Villeneuve Label
 affixed 1994.
 ♀: Formosa Sauter/ Koshun 909.III./ *Zambesa*
formosensis Town L.P. Mesnil det., 1970/ CNC
 Syntype *Zambesa claripalpis* Villeneuve Label
 affixed 1994/ EX L.-P. MESNIL COLLECTION
 1970.

Zambesomima hirsuta Mesnil, 1967, Mushi 41: 45. JAPAN.
 Holotype ♂: TYPE/ Maruyama, Sapporo, 21.VII. 1964,
 S.Takano/ 381./ *Zambesomima hirsuta* Mesnil,
 L.P. Mesnil det. 1967.

Zenillia adamsoni Thompson, 1963b, Studia ent. 6: 293.
 TRINIDAD.

Holotype ♀: HOLOTYPE Zenillia adamsoni Thomp.
CNC No. 9788/ Ex pupa of Monarch (Danaus sp.)/
St. Augustine, Trinidad, BWI. May, 1944/ A. M.
Adamson Collector 4078/ Zenillia sp. det.
James'45/ Zenillia adamsoni n.sp.

Remarks: Date of collection of holotype was published
as 19 April 1944.

Zenillia mirabilis Mesnil, 1970, Mushi 44: 108. JAPAN.
Holotype ♀: Ichihata Shimane 12.VI.1964 S.Takano/
383/ Zenillia mirabilis Mesnil L.P. Mesnil det.,
1969/ TYPE/ EX L.-P. MESNIL COLLECTION
1970.
Remarks: Shimane is spelled Schimana in the original
description.

Zenillia (Sericozenillia) albipila Mesnil, 1957, Mém. Soc. r. ent.
Belg. 28: 18. JAPAN.
Holotype ♀: 24/X 1949 Mt. Takao N.FUKUHARA/
167/ Sericozenillia albipila Mesnil L.P. Mesnil
det., 1969/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

Zeuxia rubrapex Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24):
56. ALGERIA.
Holotype ♂: MUSEUM PARIS ALGÉRIE COLL. H.
LUCAS 78-49/ Zeuxia rubrapex Mesnil L.P.
Mesnil det., 1967/ TYPE/ EX L.-P. MESNIL
COLLECTION 1985.

Zeuxia zernyi Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg. 39(24):
56. SPAIN.
Holotype ♂: Andalusia Algeciras 1.10.V.'25.Zerny/
Zeuxia zernyi Mesnil L.P. Mesnil det., 1976/
TYPE/ EX L.-P. MESNIL COLLECTION 1985.

Zizyphomyia arguta Reinhard, 1967, J. Kans. ent. Soc. 40: 109.
BRAZIL.
Holotype ♂: Nova Teutonia 27°11'S, 52°23'W Brazil,
300-500m 1. 1965 Fritz Plaumann/ HOLOTYPE
Zizyphomyia arguta Reinhard/ *Zizyphomyia*
arguta R Rnh./ *Zizyphomyia* ? Det.D.M.Wood
1966.

Zophomyia nitens Mesnil, 1963a, Bull. Inst. r. Sci. nat. Belg.
39(24): 47. JAPAN.
Holotype ♂: Obihiro S.Takano [underside of label]
19/VII'53/ 72/ Zophomyia nitens Mesnil L.P.
Mesnil det., 1970/ TYPE/ EX L.-P. MESNIL
COLLECTION 1970.

LITERATURE CITED/ BIBLIOGRAPHIE

- Aldrich, J.M. 1916. Two new Canadian Diptera. Can. Ent. 48: 20–22.
- 1926. North American two-winged flies of the genus *Cylindromyia* Meigen (*Ocyptera* of authors). Proc. U.S. natn. Mus. 68: 1–27, figs. 1–32.
- Aldrich, J.M., & R.T. Webber. 1924. The North American species of parasitic two-winged flies belonging to the genus *Phorocera* and allied genera. Proc. U.S. natn. Mus. 63: 1–90, 1 fig.
- Arnaud, P.H. 1963. Two new genera of the Germariini from the western Nearctic, with notes on related genera (Diptera, Tachinidae). Am. Mus. Novit. 2157: 1–20, figs. 1–22.
- Barracough, D.A. The systematics of the Australasian Dexiini (Diptera: Tachinidae: Dexiinae) with revisions of endemic genera. Invert. Taxon. 6: 1127–1371, figs. 1–591.
- Becker, T. 1908. Dipteren der Insel Madeira. Mitt. zool. Mus. Berl. 4: 181–206.
- 1910. Dipteren aus Südarabien und von der Insel Sokótra. Denkschr. Akad. Wiss., Wien 71: 131–160, figs. 1–5.
- Bischof, J. 1904. Beitrag zur Kenntnis der *Muscaria schizometopa*. Verh. zool.-bot. Ges. Wien 54: 79–101.
- Brooks, A.R. 1942. *Clistomorpha*, *Psalidopteryx* and allies (Diptera, Tachinidae). Can. Ent. 74: 140–150, figs. 1–4.
- 1943. A review of the Canadian species of *Ernestia* sens. lat. (Tachinidae, Diptera). Can. Ent. 75: 66–78, figs. 1–22.
- 1944a. A review of the North American species of *Gonia* sens. lat. (Diptera, Tachinidae). Can. Ent. 75 (1943): 219–236, figs. 1–13.
- 1944b. A review of the North American species of *Linnaemya* sens. lat. (Diptera, Tachinidae). Can. Ent. 76: 193–206, figs. 1–11.
- 1945a. A revision of the North American species of the *Phasia* complex (Diptera, Tachinidae). Scient. Agric. 25: 647–679, figs. 1–28.
- 1945b. New Canadian Diptera (Tachinidae). Can. Ent. 77: 78–96.
- 1946. A revision of the North American species of the *Rhodogyne* complex (Diptera, Larvaevoridae). Can. Ent. 77 (1945): 218–230, figs. 1–21.
- 1947. A revision of the North American species of *Leschenaultia* sens. lat. (Diptera, Larvaevoridae). Can. Ent. 78 (1946): 169–182, figs. 1–23.
- 1949. New North American larvaevorine flies (Diptera, Larvaevoridae). Can. Ent. 81: 21–24.
- Cortés, R. 1975. Taquinidos Chilenos nuevos o poco conocidos, IV-(Diptera, Tachinidae). Boln téc Estac. exp. agron. Fac. Agronomía Univ. Chile. 40: 3–14.
- Crosskey, R.W. 1966. The putative fossil genus *Palexorista* Townsend and its identity with *Prosturmia* Townsend (Diptera: Tachinidae). Proc. R. ent. Soc. Lond. (B). 35: 133–137.
- 1976. A taxonomic conspectus of the Tachinidae (Diptera) of the Oriental Region. Bull. Br. Mus. nat. Hist. (Ent.) Suppl. 26: 1–357, figs. 1–150.
- Curran, C.H. 1923. New North American Diptera. Can. Ent. 55: 245–246.
- 1924a. New species of *Ernestia* and *Mericia* (Dipt.: Tachinidae). Ent. News 35: 245–250.
- 1924b. Four apparently undescribed Diptera from Canada. Can. Ent. 56: 250–253.
- 1924c. New Canadian Diptera, with synopsis of the genus *Cynorrhina*. Can. Ent. 56: 193–196.
- 1924d. On the genus *Arctophyto* Townsend in North America (Tachinidae, Diptera). Can. Ent. 56: 302–303.
- 1925a. Some apparently new Nearctic Tachinidae (Diptera). Can. Ent. 57: 281–286.
- 1925b. New Tachinidae in the Canadian National Collection (Diptera). Can. Ent. 57: 150–156.
- 1925c. The American species of the tachinid genus *Peleteria* Desv. (Diptera). Trans. R. Soc. Can., Ser. 3, 19: 225–257, 2 pls.
- 1925d. Three new Nearctic Tachinidae (Dipt.). Ent. News 36: 13–18.
- 1926a. The Canadian species of the tachinid genera *Cryptomeigenia* B.B. and *Tachinomyia* Tns. (Dipt.). Trans. R. Soc. Can., Ser. 3, 20: 155–171, 2 pls.
- 1926b. *Grisdalemyia*, a new genus of Tachinidae (Diptera). Can. Ent. 58: 133–135, figs. 1–2.
- 1926c. Descriptions of new Canadian Diptera. Can. Ent. 58: 211–218.
- 1926d. New Nearctic Diptera mostly from Canada. Can. Ent. 58: 81–89.
- 1927a. Some new American Tachinidae (Diptera). Bull. Brooklyn ent. Soc. 22: 144–154.
- 1927b. Some new North American Diptera. Can. Ent. 59: 290–303, figs. 1–10.
- 1927c. Studies in Canadian Diptera-III. The species of the tachinid genera related to *Lydella*, as represented in the Canadian National Collection. Can. Ent. 59: 11–24.

- 1928a. Two new species of *Wagneria* (Tachinidae, Dipt.). Can. Ent. 60: 48-49.
- 1928b. Revision of the American species of *Archytas* (Tachinidae, Diptera). Can. Ent. 60: 218-226; 249-256; 275-282, pls. 17-19.
- 1929. Some new Nearctic Diptera. Can. Ent. 61: 30-34, 1 fig.
- 1930a. Three new Diptera from Canada. Jl N.Y. ent. Soc. 38: 73-76.
- 1930b. A new tachinid parasitic on a sawfly. Can. Ent. 62: 246-247.
- 1933. The North American species of *Actia* in the American Museum of Natural History. Am. Mus. Novit. 614: 1-7.
- Guimarães, J.H. 1963. Primeira contribuição ao conhecimento da tribo Cuphoceratini (Diptera, Tachinidae). Mem. Inst. Oswaldo Cruz 61: 41-72, figs. 1-121.
- 1972. A revision of the genus *Winthemia* Robineau-Desvoidy in America north of Mexico (Diptera, Tachinidae). Archos Zool. S Paulo 22: 27-112, figs. 1-93.
- 1976. A revision of the genus *Cylindromyia* Meigen in the Americas south of the United States (Diptera, Tachinidae). Archos Zool. S Paulo 27: 1-50, figs. 1-85.
- Hermann, F. & J. Villeneuve. 1909. Pp. 147-159. Diptera. Aus der Sinaihalbinsel In Kneucker, A., Zoologische Ergebnisse zweier in den Jahren 1902 und 1904 durch die Sinaihalbinsel unternommener botanischer Studienreisen nebst zoologischen Beobachtungen aus Ägypten, Palästina und Syrien. Verh. naturw. Ver. Karlsruhe 21.
- Herting, B. 1966. Beiträge zur Kenntnis der europäischen Raupenfliegen (Dipt. Tachinidae). IX. Stuttg. Beitr. Naturk. 146: 1-12.
- 1967. Beiträge zur Kenntnis der europäischen Raupenfliegen (Dipt. Tachinidae). Stuttg. Beitr. Naturk. 173: 1-11, figs. 1-9.
- 1975. Neue paläarktische Tachiniden (Diptera). Stuttg. Beitr. Naturk. (A) 287: 1-7, figs. 1-2.
- 1977. Beiträge zur Kenntnis der europäischen Raupenfliegen (Dipt. Tachinidae). Stuttg. Beitr. Naturk. (A) 295: 1-16, figs. 1-14.
- 1979. Beschreibungen neuer Raupenfliegen (Dipt. Tachinidae) und Revision der *Besseria anthophila*-Gruppe. Stuttg. Beitr. Naturk. (A) 323: 1-10, figs. 1-4.
- 1983. Phasiinae. Fliegen palaearkt. Reg. 64c: 1-88, figs. 1-83.
- 1984. Catalogue of Palearctic Tachinidae (Diptera). Stuttg. Beitr. Naturk. (A) 369: 1-228.
- Jacentkovski, I.D. 1944. Kuklice (Tachinoidea, Diptera) Zdánského lesa. Raupenfliegen (Tachinoidea, Diptera) des Steinitzer Waldes. Sb. ent. Odd. nár. Mus. Praze 21-22: 380-395.
- Malloch, J.R. 1919. The Diptera collected by the Canadian Expedition, 1913-1918 (excluding the Tipulidae and Culicidae). Pp. 34-90, pls. 7-10. In Anderson, R.M., ed., Rep. Can. arct. Exped. 1913-18 [q.v.]. Vol. 3: Insects, Pt. C: Diptera, 90 pp., 2 figs., 10 pls. Ottawa, Ont.
- Mesnil, L.P. 1939a. Essai sur les Tachinaires (Larvaevoridae). Monogr Sta. Lab. Rech. agron. 7: 1-67, 2 pls.
- 1939b. Descriptions d'espèces nouvelles de Tachinaires (Dipt. Larvaevoridae). Bull. Annls Soc. ent. Belg. 79: 209-212.
- 1941a. Espèces nouvelles du genre *Exorista* Meig. (Dipt. Larvaevoridae). Bull. Soc. ent. Fr. 46: 20-22.
- 1941b. Notes synonymiques. Bull. Soc. ent. Fr. 46: 98.
- 1944a. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 1-48.
- 1944b. Nouveaux Larvaevoridae exotiques du Muséum de Paris (Diptera). Revue fr. Ent. 11: 10-17.
- 1946. Revision des Phorocerini de l'ancien monde (Larvaevoridae). Encycl. ent. (B) II Dipt. 10: 37-80, figs 1-4.
- 1949a. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 49-104.
- 1949b. Essai de révision des espèces du genre *Drino* Robineau-Desvoidy Sturmia a oeufs macrotypes. Bull. Inst. r. Sci. nat. Belg. 25(42): 1-38.
- 1950a. Notes sur les Carceliina (Dipt. Tachinidae) et révision des espèces d'Afrique. Revue Zool. Bot. afr. 43: 1-24.
- 1950b. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 105-160.
- 1950c. Critiques et suggestions à propos de récents travaux concernant les Protachinides d'Afrique. Bull. Annls Soc. ent. Belg. 86: 104-117.
- 1951. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 161-208.
- 1952a. Seconde note préliminaire sur les Tachinaires du Congo Belge (suite). Bull. Inst. r. Sci. nat. Belg. 28(23): 1-18.
- 1952b. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 209-256.
- 1952c. Notes détachées sur quelques Tachinaires paléarctiques. Bull. Annls Soc. ent. Belg. 88: 149-158.
- 1953a. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 257-304.
- 1953b. Nouveaux Tachinaires d'Orient (1re et 2e partie). Bull. Annls Soc. ent. Belg. 89: 85-114, 146-178.
- 1954a. Larvaevoridae (Tachininae). Fliegen palaearkt. Reg. 64g: 305-368.

- 1954b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 369-416.
- 1954c. Revision du genre *Thelairosoma* Villeneuve (Diptera Tachinidae). Annls Mus. r. Congo belge Sér. 4to (Zool.) 1: 469-474.
- 1955a. Contributions à l'étude de la faune entomologique du Ruanda-Urundi (Mission P. Basilewsky 1953). LXXI. Diptera Tachinidae. Annls Mus. r. Congo belge Sér. 8vo (Zool.) 40: 359-367.
- 1955b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 417-464.
- 1956a. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 465-512.
- 1956b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 513-554.
- 1956c. Trois nouveaux Tachinaires d'Afrique (Dipt. Tachinidae). Entomophaga 1: 76-80.
- 1957. Nouveaux Tachinaires d'Orient (Deuxième série). Mém. Soc. r. ent. Belg. 28: 1-80.
- 1959. Tachinidae d'Afrique orientale (Dipt.). Stuttg. Beitr. Naturk. 23: 1-31.
- 1960a. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 561-608.
- 1960b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 609-656.
- 1960c. Note préliminaire sur les *Siphona* Meig. (Dipt. Tachinidae) d'Europe et du bassin méditerranéen. Bull. Annls Soc. r. ent. Belg. 96: 187-192.
- 1961a. Deux nouvelles *Siphona* Meigen (Dipt. Tachinidae) d'Europe. Bull. Annls Soc. r. ent. Belg. 97: 201-204.
- 1961b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 657-704.
- 1962. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 753-800.
- 1963a. Nouveaux Tachinaires de la Region Palearctique principalement de l'URSS et du Japan. Bull. Inst. r. Sci. nat. Belg. 39(24): 1-56.
- 1963b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 801-848.
- 1964. Description d'une nouvelle espèce d'Ormiini récemment découverte dans le Sud de la France (Dipt. Tachinidae). Bull. Soc. ent. Fr. 69: 261-264.
- 1966. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 881-928.
- 1967. Tachinaires Paléarctiques inédits (Diptera). Mushi 41: 37-57, Figs 1-3.
- 1968a. Nouveaux Tachinaires d'Orient (Troisième série). Bull. Annls Soc. r. ent. Belg. 104: 173-188.
- 1968b. Quelques espèces inédites de Tachinaires africains (Dipt. Tachinidae). Stuttg. Beitr. Naturk. 187: 1-12.
- 1969. Quelques Tachinaires nouveaux de Mélanésie (Dipt. Tachinidae). Entomophaga 13: 203-208.
- 1970. Description de nouveaux Tachinaires de l'ancien monde, et notes synonymiques (Diptera, Tachinidae). Mushi 44: 89-123, figs. 1-6.
- 1971a. Quelques nouveaux Tachinaires (Dipt. Tachinidae) de l'ancien monde. Entomophaga 16: 67-73.
- 1971b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 977-1024.
- 1972. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 1065-1112.
- 1973a. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 1113-1168.
- 1973b. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 1169-1232.
- 1974. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 1233-1304.
- 1975a. Deux espèces de Tachinaires (Dipt. Tachinidae) nouvelles et intéressantes. Stuttg. Beitr. Naturk. (A) 277: 1-5.
- 1975b. Larvaevorinae (Tachinidae). Fliegen palaearkt. Reg. 64g: 1305-1384.
- 1975c. Larvaevorinae (Tachininae). Fliegen palaearkt. Reg. 64g: 1385-1435.
- 1977. Nouveaux Tachinaires de Madagascar (Dipt. Tachinidae) 6e partie. Bull. Soc. ent. Suisse 50: 321-329.
- 1980. Dexiinae. Fliegen palaearkt. Reg. 64f: 1-52.
- Mesnil, L.P. & H. Pschorn-Walcher. 1968. A preliminary list of Tachinidae (Diptera) from Japan. Mushi 41: 149-174.
- Morrison, F.O. 1940. A revision of the American species of *Gonia* Meigen (Diptera: Tachinidae). Can. J. Res. 18: 336-362, figs. 1-22a.
- O'Hara, J.E. 1983. Classification, phylogeny and zoogeography of the North American species of *Siphona* Meigen (Diptera: Tachinidae). Quaest. Ent. 18 (1982): 261-380, figs. 1-108.
- 1984. *Baeomyia* n.g. (Diptera: Tachinidae): descriptions and notes about phylogenetic and zoogeographic relationships. Can. J. Zool. 62: 1387-1396, figs. 1-15.
- 1989. Systematics of the genus group taxa of the Siphonini (Diptera: Tachinidae). Quaest. Ent. 25: 1-229, figs. 1-171.
- 1991. Revision of Nearctic species of *Actia* Robineau-Desvoidy (Diptera: Tachinidae). Can. Ent. 123: 745-776, figs. 1-30.

- 1993. Revision of the species of *Frontiniella* Townsend (Diptera: Tachinidae). Can. Ent. 125: 11–45, figs. 1–45.
- 1994. Revision of Nearctic species of *Ceromyia* Robineau-Desvoidy (Diptera: Tachinidae). Can. Ent. 126: 775–806, figs. 1–29.
- In press. The tachinid taxa of Louis P. Mesnil, with notes on nomenclature (Insecta: Diptera). Can. Ent.
- Pandellé, L. 1896. Études sur les Muscides de France. II partie (suite). Revue Ent. 15: 1–230.
- Reinhard, H.J. 1922. Some new species of Texas Tachinidae (Diptera). Ann. ent. Soc. Am. 14 (1921): 329–335, figs. 1–10.
- 1934a. North American parasitic flies of the genus *Spathidexia* with descriptions of two new species. Bull. Brooklyn ent. Soc. 29: 150–154.
- 1934b. New North American Tachinidae. Bull. Brooklyn ent. Soc. 29: 186–195.
- 1934c. Two new species of the tachinid genus *Siphosturmopsis* with key and notes (Diptera). Ent. News 45: 15–19.
- 1935a. North American two-winged flies of the genus *Doryphorophaga*, (Tachinidae, Diptera). Jl N.Y. ent. Soc. 43: 387–394.
- 1935b. Notes on the tachinid genus *Pseudotachinomyia* with descriptions of two new species (Diptera). Ent. News 46: 132–135.
- 1935c. New genera and species of American muscoid flies (Tachinidae: Diptera). Ann. ent. Soc. Am. 28: 160–173.
- 1937. New North American muscoid Diptera. Bull. Brooklyn ent. Soc. 32: 62–74.
- 1938. Four new Nearctic species of *Fabriciella*, (Tachinidae, Diptera). Can. Ent. 70: 8–11.
- 1939. New genera and species of muscoid Diptera. Bull. Brooklyn ent. Soc. 34: 61–74.
- 1942a. A new species of *Oedematocera* with notes and key (Tachinidae, Diptera). Ent. News 53: 106–108.
- 1942b. New North American Tachinidae belonging to the genera *Microchaetina* and *Hypenomyia* with key to the known species (Diptera). Can. Ent. 74: 88–91.
- 1942c. Notes on *Fabriciella* with descriptions of five new species (Tachinidae, Diptera). Bull. Brooklyn ent. Soc. 37: 24–30.
- 1943a. New genera of North American muscoid Diptera. Can. Ent. 75: 163–169.
- 1943b. New North American Muscoidea (Tachinidae, Diptera). J. Kans. ent. Soc. 16: 14–23.
- 1943c. New Tachinidae from Northeastern United States (Diptera). Bull. Brooklyn ent. Soc. 38: 78–90.
- 1944a. New North American Tachinidae belonging to the genus *Muscopteryx* (Diptera). Ann. ent. Soc. Am. 37: 352–358.
- 1944b. New muscoid Diptera from the United States. J. Kans. ent. Soc. 17: 57–72.
- 1945. New genera and species of North American Tachinidae (Diptera). Can. Ent. 77: 28–36.
- 1946a. A review of the tachinid genera *Siphophyto* and *Coronimyia* (Diptera). Proc. ent. Soc. Wash. 48: 79–92.
- 1946b. The tachinid genera *Pseudochaeta* and *Phaenopsis* in North America, (Diptera). Can. Ent. 78: 111–121.
- 1946c. The genus *Minthozelia* in the United States (Diptera, Tachinidae). J. Kans. ent. Soc. 19: 52–59.
- 1947. New genera and species of muscoid Diptera. J. Kans. ent. Soc. 20: 15–24.
- 1951. New American muscoid Diptera. Bull. Brooklyn ent. Soc. 46: 1–9.
- 1952a. Muscoid flies of the genus *Chaetophlepsis* (Diptera). J. Kans. ent. Soc. 25: 13–21.
- 1952b. New genera and species of muscoid Diptera. Bull. Brooklyn ent. Soc. 47: 1–12.
- 1953a. New species of Tachinidae from Mexico (Diptera). Bull. Brooklyn ent. Soc. 48: 89–96.
- 1953b. New Mexican Tachinidae (Diptera). J. Kans. ent. Soc. 26: 95–102.
- 1953c. New muscoid Diptera from the western United States. Pan-Pacif. Ent. 29: 49–59.
- 1953d. Notes on muscoid synonymy with descriptions of three new species (Diptera). Proc. ent. Soc. Wash. 55: 243–247.
- 1955a. New genera and species of North American Tachinidae (Diptera). J. Kans. ent. Soc. 28: 123–130.
- 1955b. North American tachinid flies of the genus *Wagneria* (Diptera). J. Kans. ent. Soc. 28: 49–59.
- 1955c. New Nearctic Sarcophagidae and Tachinidae (Diptera). Bull. Brooklyn ent. Soc. 50: 128–133.
- 1955d. North American Muscoidea (Diptera: Tachinidae). Ent. News 66: 233–238.
- 1956a. A synopsis of the tachinid genus *Leucostoma* (Diptera). J. Kans. ent. Soc. 29: 155–168.
- 1956b. New muscoid Diptera mainly from California. Pan-Pacif. Ent. 32: 103–110.
- 1956c. New Tachinidae (Diptera). Ent. News 67: 121–129.
- 1957. New American muscoid Diptera (Sarcophagidae, Tachinidae) Ent. News 68: 99–111.

- 1958a. Notes on *Spathimeigenia* with descriptions of four new species (Diptera, Tachinidae). Proc. ent. Soc. Wash. 60: 207–212.
- 1958b. New genera and species of North American Tachinidae (Diptera). J. Kans. ent. Soc. 31: 225–232.
- 1958c. Parasitic flies of the genus *Mochlosoma* (Tachinidae, Diptera). Can. Ent. 90: 98–110.
- 1958d. North American Tachinidae (Diptera). J. Kans. ent. Soc. 31: 277–284.
- 1958e. New American Tachinidae (Diptera). Ent. News 69: 233–242.
- 1959a. New North American Tachinidae (Diptera). Ent. News 70: 225–234.
- 1959b. New Nearctic Tachinidae (Diptera). Pan-Pacif. Ent. 35: 157–163.
- 1961. New American Tachinidae and Sarcophagidae (Diptera). J. Kans. ent. Soc. 34: 204–213.
- 1962a. North American muscoid Diptera. Ent. News 73: 169–178.
- 1962b. New North American Tachinidae (Diptera). Pan-Pacif. Ent. 38: 215–224.
- 1964a. North American Tachinidae (Diptera). J. Kans. ent. Soc. 37: 35–51.
- 1964b. Mexican Tachinidae (Diptera). Acta zool. mex. 7: 1–21.
- 1964c. Parasitic flies of the genera *Helioptrosopa* and *Euhelioptrosopa* (Diptera : Tachinidae). Pan-Pacif. Ent. 40: 117–124.
- 1967. New Nearctic and Neotropical muscoidean Diptera (Sarcophagidae and Tachinidae). J. Kans. ent. Soc. 40: 94–110.
- 1968a. A review of the genus *Macromya* (Tachinidae: Diptera). Can. Ent. 100: 1280–1287.
- 1968b. New world species of *Erynnia* (Tachinidae, Diptera). J. Kans. ent. Soc. 41: 501–508.
- 1974. New genera and species of American Tachinidae (Diptera). Can. Ent. 106: 1155–1170.
- Roubaud, E. & J. Villeneuve. 1914. Contribution à l'étude des espèces du genre *Anacamptomyia* Bischof (Dipt.). Revue zool. afr. 4: 121–128.
- Rowe, J.A. 1931. A revision of the males of the Nearctic species in the genus *Fabriciella* (Tachinidae). Ann. ent. Soc. Am. 24: 643–678, figs. 1–25.
- Sabrosky, C.W. 1981. A partial revision of the genus *Eucelatoria* (Diptera, Tachinidae), including important parasites of *Heliothis*. Tech. Bull. U.S. Dep. Agric. 1635: 1–18, figs. 1–14.
- Say, T. 1829. Descriptions of North American dipterous insects. J. Acad. nat. Sci. Philad. 6: 149–178.
- Strickland, E.H. 1941. A new genus of the family Tachinidae from Alberta. Can. Ent. 73: 64–66, figs. 1–2.
- Thompson, W.R. 1963a. The tachinids of Trinidad II. Echinomyiines, Dexiines, and allies. Can. J. Zool. 41: 335–576, figs. 1–174.
- 1963b. The tachinids of Trinidad III. The Goniines with microtype eggs (Dipt. Tachinidae). Studia ent. 6: 257–404, figs. 1–67.
- 1963c. The tachinids of Trinidad IV. Winthemiines. Can. Ent. 95: 953–995, figs. 1–38.
- 1963d. The tachinids of Trinidad V. Siphosturmiiines and Masiphyiines. Can. Ent. 95: 1292–1320, figs. 1–29.
- 1964. The tachinids of Trinidad. VI (Dipt. Tachinidae). Studia ent. 7: 97–151, figs. 1–46.
- 1966. The tachinids of Trinidad VII. The larviparous Goniines with broad cheeks (Dipt. Tachinidae). Studia ent. 8 (1965): 353–434, figs. 1–43.
- 1968. The tachinids of Trinidad VIII. Phorocerines. Mem. ent. Soc. Can. 56: 1–207, figs. 1–140.
- Tothill, J.D. 1912. Systematic notes on North American Tachinidae. Part I. Can. Ent. 44: 1–5.
- 1921. A revision of the Nearctic species of the tachinid genus *Ernestia* R.D. (Diptera). Can. Ent. 53: 247–252, 270–274.
- 1922. The natural control of the fall webworm (*Hypantria cunea* Drury) in Canada together with an account of its several parasites. Bull. Can. Dep. Agric. (n. ser.) 3: 1–107, figs. 1–99, 6 pls.
- 1924a. A revision of the Nearctic species of the genus *Gonia* (Diptera, Tachinidae). Can. Ent. 56: 196–200, 206–212.
- 1924b. A revision of the Nearctic species in the genus *Fabriciella* (Tachinidae). Can. Ent. 56: 257–269.
- Townsend, C.H.T. 1892a. Notes on North American Tachinidae sens. str. with descriptions of new genera and species. Paper III. Trans. Am. ent. Soc. 19: 88–132.
- 1892b. Notes on North American Tachinidae, with descriptions of new genera and species. – Paper V. Can. Ent. 24: 64–70.
- Villeneuve, J. 1908a. Travaux diptérologiques. Wien. ent. Ztg. 27: 281–288.
- 1908b. Contribution au catalogue des Diptères de France. Feuille jeun. Nat. 38: 114–118.
- 1910a. Diptères nouveaux. Wien. ent. Ztg. 29: 86–92.

- 1910b. Description de nouvelles espèces de Tachinaires provenant de l'Afrique occidentale. Wien. ent. Ztg 29: 249–254.
- 1911. Notes diptérologiques. Wien. ent. Ztg 30: 84–87.
- 1912a. Diptères nouveaux du Nord Africain. Bull. Mus. natn. Hist. nat., Paris 18: 415–417, 1 fig.
- 1912b. Diptères nouveaux du Nord Africain (Deuxième note). Bull. Mus. natn. Hist. nat., Paris 18: 505–511.
- 1913. Myodaires supérieurs de l'Afrique tropical (I^e liste). Revue zool. afr. 3: 24–46.
- 1914. Sur quatre formes nouvelles se rapportant aux 'Oestridae dubiosae B. B.'. Annls hist.-nat. Mus. natn. hung. 12: 435–442, figs. 1–4.
- 1915a. Myodaires supérieurs recueillis à Madagascar. Revue zool. afr. 4: 191–209.
- 1915b. Nouveaux Myodaires supérieurs de Formose. Annls hist.-nat. Mus. natn. hung. 13: 90–94, figs. 1–3.
- 1916a. Espèces africaines nouvelles du genre *Nemoraea* R.-D. (Dipt.). Annls Soc. ent. Fr. 85: 197–202.
- 1916b. A new species of Tachino-oestrid from South Africa (Diptera). Ann. S. Afr. Mus. 15: 465–468, figs. 1–3.
- 1916c. A contribution to the study of the South African higher Myodarii (Diptera Calyptratae) based mostly on the material in the South African Museum. Ann. S. Afr. Mus. 15: 469–515, figs. 1–8.
- 1919. Descriptions de deux Tachinaires nouveaux (Dipt.). Bull. Soc. ent. Fr. 1919: 264–266.
- 1920a. Étude de quelques Myodaires supérieurs (recueillis par le Dr Brauns, à Willowmore, Cap). Revue zool. afr. 8: 151–162.
- 1920b. Diptères paléarctiques nouveaux ou peu connus. Annls Soc. ent. Belg. 60: 114–120.
- 1920c. Diptères inédits. Annls Soc. ent. Belg. 60: 199–205.
- 1921. Descriptions d'espèces nouvelles du genre *Actia* Rob. Desv. Annls Soc. ent. Belg. 61: 45–47.
- 1922a. Description de deux espèces nouvelles de Tachinaires nord-africains. Bull. Mus. natn. Hist. nat., Paris 28: 291–294, figs. 1–2.
- 1922b. Myodaires supérieurs du Soudan communiqués par M. le Prof. Rich. Ebner, de Vienne. Revue zool. afr. 10: 62–65.
- 1922c. Myodaires supérieurs paléarctiques nouveaux. Annls Sci. nat. Zool. (10)5: 337–342.
- 1923a. Myodaires supérieurs d'Egypte inédits (Dipt.). Bull. Soc. r. ent. Égypte 7 (1922): 88–94.
- 1923b. Descriptions de Phasiinae nouveaux (Dipt.). Revue zool. afr. 11: 78–81.
- 1925. Descriptions de nouveaux Tachino-oestrides (Dipt.). Konowia 4: 48–52.
- 1926a. Myodaires supérieurs de l'Afrique nouveaux ou peu connus. Revue zool. afr. 14: 64–69.
- 1926b. Sur *Masicera casta* Rond. et espèces affines. Revue zool. afr. 14: 242–247.
- 1926c. Descriptions de Myodaires supérieurs nouveaux. Bull. Annls Soc. ent. Belg. 66: 269–275.
- 1926d. Descriptions de nouveaux Tachinidae (Dipt.) de l'Europe méridionale. Bull. Annls Soc. ent. Belg. 66: 197–200.
- 1927. Description d'un nouveau Tachino-oestride africain (Dipt.) et autres descriptions. Revue zool. afr. 15: 118–122.
- 1928. Myodaires supérieurs nouveaux (Dipt.). Bull. Annls Soc. ent. Belg. 68: 47–52.
- 1929a. Propos diptérologiques (suite). Bull. Annls Soc. ent. Belg. 69: 181–187.
- 1929b. Diagnoses de Myodaires supérieurs inédits. Bull. Annls Soc. ent. Belg. 69: 99–102.
- 1930a. Description de deux Myodaires supérieurs sud-africains. Bull. Annls Soc. ent. Belg. 69 (1929): 352–353.
- 1930b. Propos diptérologiques. Bull. Annls Soc. ent. Belg. 70: 41–45.
- 1930c. Diptères inédits. Bull. Annls Soc. ent. Belg. 70: 98–104.
- 1930d. Description d'un *Sturmia* congolais nouveau (Dipt.). Revue Zool. Bot. afr. 20: 59–60.
- 1931. Aperçus critiques sur le mémoire de P. Stein: "Die verbreitetsten Tachiniden Mitteleuropas". Konowia 10: 47–74.
- 1932a. Descriptions de nouveaux Myodaires supérieurs paléarctiques (Dipt.). Bull. Annls Soc. ent. Belg. 71 (1931): 241–245.
- 1932b. Descriptions de Myodaires supérieurs (Larvaevoridae) nouveaux de Formose. Bull. Soc. ent. Fr. 37: 268–271.
- 1933. Myodaires supérieurs africains nouveaux. Revue Zool. Bot. afr. 23: 278–280.
- 1934a. Myodaires supérieurs inédits. Revue Zool. Bot. afr. 25: 408–411.
- 1934b. Myodaires supérieurs inédits d'Afrique. Revue Zool. Bot. afr. 26: 68–72, fig. 1.
- 1935a. Sur le genre *Mormonomya* Br.-Berg. (Dipt.). Bull. Soc. ent. Fr. 40: 251–253.

- 1935b. Myodaires supérieurs africains inédits. *Revue Zool. Bot. afr.* 27: 136–143.
- 1936a. Descriptions de Larvaevoridae africains (Dipt.). *Bull. Mus. r. Hist. nat. Belg.* 12(4): 1–10.
- 1936b. Notes sur quelques Larvaevoridae africains. *Bull. Mus. r. Hist. nat. Belg.* 12(12): 1–5.
- 1936c. Myodaires supérieurs nouveaux du continent africain. *Bull. Mus. r. Hist. nat. Belg.* 12(41): 1–3.
- 1936d. Myodaires supérieurs de Chine. *Bull. Mus. r. Hist. nat. Belg.* 12(42): 1–7.
- 1936e. Schwedisch-chenesische wissenschaftliche Expedition nach den nordwestlichen Provinzen chinas, unter Leitung von Dr. Sven Hedin und Prof. Sü Ping-chang. Insekten gesammelt vom schwedischen Arzt der Expedition Dr. David Hummel 1927–1930. 52. Diptera. 16. Muscidae. *Ark. Zool.* 27A: 1–13.
- 1936f. Myodaires supérieurs africains récoltés à Kampala (Uganda) par M.H. Hargreaves. *Bull. Annls Soc. ent. Belg.* 76: 415–419, 1 fig.
- 1937a. Descriptions de Myodaires supérieurs. *Revue Zool. Bot. afr.* 29: 205–212, figs. 1–4.
- 1937b. Myodaires supérieurs inédits de la région Paléarctique. *Bull. Mus. r. Hist. nat. Belg.* 13(13): 1–4.
- 1937c. Myodaires supérieurs de Chine. *Bull. Mus. r. Hist. nat. Belg.* 13(34): 1–16, figs. 1–5.
- 1937d. Myodaires supérieurs africains. *Bull. Mus. r. Hist. nat. Belg.* 13(35): 1–4.
- 1937e. Notes diptérologiques. *Bull. Annls Soc. ent. Belg.* 77: 404–408.
- 1938a. *Anurophylla* nov. subg. des Myodaires supérieurs. *Bull. Annls Soc. ent. Belg.* 78: 413–414, figs. 1–2.
- 1938b. Myodaires supérieurs inédits. *Bull. Mus. r. Hist. nat. Belg.* 14(4): 1–5, figs. 1–2.
- 1938c. Myodaires africains (notes et espèces inédites). *Bull. Mus. r. Hist. nat. Belg.* 14(38): 1–16.
- 1939a. Présentation de quelques Myodaires supérieurs inédits. *Bull. Annls Soc. ent. Belg.* 79: 347–354, figs. 1–3.
- 1939b. Myodaires supérieurs africains (descriptions et observations). *Bull. Mus. r. Hist. nat. Belg.* 15(48): 1–10.
- 1941a. De quelques espèces africaines inédites du genre *Linnaemyia* R. D. (Dipt. Tachinidae). *Bull. Soc. ent. Fr.* 46: 107–110.
- 1941b. Myodaires supérieurs nouveaux (Dipt.). *Bull. Soc. ent. Fr.* 46: 122–126.
- 1942a. Descriptions de Myodaires supérieurs nouveaux (Dipt. Tachinidae). *Bull. Soc. ent. Fr.* 47: 50–55.
- 1942b. Espèces inédites de la famille des Larvaevoridae (Dipt.). *Bull. Soc. ent. Fr.* 47: 133–135.
- 1943a. Myodaires supérieurs inédits (Dipt.). *Bull. Soc. ent. Fr.* 48: 36–40.
- 1943b. Sur les Dexiaires africains (Dipt.). *Revue Zool. Bot. afr.* 37: 93–96.
- 1944. Myodaires supérieurs nouveaux (Dipt.). *Bull. Soc. ent. Fr.* 48 (1943): 144–145.
- Villeneuve, J. & L. Mesnil. 1936. Au sujet du Genre *Salia* R. D. (Dipt. Tachinidae). *Bull. Soc. ent. Fr.* 41: 270–276.
- Wilder, D.D. 1979. Systematics of the Nearctic *Ptilodexia* Brauer and Bergenstamm (Diptera: Tachinidae). *Proc. Calif. Acad. Sci.* 42: 1–55, figs. 1–113.
- Williston, S.W. 1886. Dipterological notes and descriptions. *Trans. Am. ent. Soc.* 13: 287–307.
- Wood, D.M. 1972. A revision of the new world Exoristini (Diptera: Tachinidae) I. *Phorocera* subgenus *Pseudotachinomyia*. *Can. Ent.* 104: 471–503, figs. 1–30.
- 1974a. A new species of *Masistylum* in the Nearctic region (Diptera: Tachinidae). *Can. Ent.* 106: 175–178, figs. 1a–4c.
- 1974b. Notes on *Allophorocera* with a description of a new species (Diptera: Tachinidae) from Finland. *Can. Ent.* 106: 667–671, figs. 1a–4b.
- 1985. A taxonomic conspectus of the Blondeliini of North and Central America and the West Indies (Diptera: Tachinidae). *Mem. ent. Soc. Can.* 132: 1–130, figs. 1–56.

INDEX

<i>abdominalis</i> Mesnil, <i>Medina</i>	49
<i>aberrans</i> Thompson, <i>Pseudosiphosturmia</i>	66
<i>aberrans</i> Villeneuve, <i>Ocyptera</i>	55
<i>abrelicta</i> Mesnil, <i>Carcelia</i>	21
<i>acanthina</i> Mesnil, <i>Picconia incurva</i>	63
<i>aclista</i> Reinhard, <i>Peleteria</i>	59
<i>actinosa</i> Reinhard, <i>Fabriciella</i>	37
<i>adamsoni</i> Thompson, <i>Zenillia</i>	78
<i>aemulus</i> Reinhard, <i>Sitellitergus</i>	70
<i>aequalis</i> Reinhard, <i>Pseudotachinomyia</i>	67
<i>aequalis</i> Villeneuve, <i>Carcelia</i>	21
<i>aethiopica</i> Mesnil, <i>Ctenophorocera</i>	26
<i>africana</i> Villeneuve, <i>Paraprosena marmorata</i>	58
<i>africanum</i> Mesnil, <i>Dolichocolon</i>	29
<i>africanum</i> Villeneuve, <i>Leucostoma</i>	45
<i>agilis</i> Reinhard, <i>Ptilodexia</i>	67
<i>agrianiomei</i> Mesnil, <i>Billaea</i>	17
<i>akidomyia</i> O'Hara, <i>Siphona</i>	69
<i>alactaga</i> Reinhard, <i>Doliolomyia</i>	29
<i>albagena</i> Morrison, <i>Gonia</i>	39
<i>alberta</i> Curran, <i>Clistomorpha</i>	25
<i>alberta</i> Curran, <i>Hypostenia</i>	43
<i>alberta</i> Curran, <i>Mericia</i>	49
<i>alberta</i> Curran, <i>Peleteria</i>	59
<i>albertana</i> Reinhard, <i>Myioclonia</i>	53
<i>albescens</i> Pandellé, <i>Pachystylum</i>	57
<i>albiceps</i> Reinhard, <i>Meledonus</i>	49
<i>albidopilosa</i> Mesnil, <i>Winthemia</i>	78
<i>albifacies</i> Villeneuve, <i>Plesiooestrus</i>	63
<i>albifrons</i> Villeneuve, <i>Atylomyia</i>	16
<i>albipennis</i> Brooks, <i>Paraphasia</i>	58
<i>albipennis</i> Villeneuve, <i>Heraultia</i>	42
<i>albipila</i> Mesnil, <i>Zenillia</i>	79
<i>albisquama</i> Villeneuve, <i>Litophasia</i>	47
<i>albocincta</i> Villeneuve, <i>Bucentes</i>	19
<i>aldrichi</i> Curran, <i>Erycia</i>	33
<i>aldrichi</i> Curran, <i>Hesperophasia</i>	42
<i>aldrichi</i> Curran, <i>Paralispe</i>	58
<i>aldrichi</i> Tohill, <i>Gonia</i>	39
<i>aldrichi</i> Townsend, <i>Hystricia</i>	44
<i>alearis</i> Reinhard, <i>Masiphyomyia</i>	48
<i>algens</i> Curran, <i>Arctophyto</i>	15
<i>alipes</i> Villeneuve, <i>Actia</i>	11
<i>alpina</i> Villeneuve, <i>Wagneria</i>	77
<i>ambulatoria</i> Villeneuve, <i>Ocyptera</i>	55
<i>americana</i> Brooks, <i>Thelaira</i>	74
<i>americana</i> Curran, <i>Eipogona</i>	32
<i>americana</i> Reinhard, <i>Minella</i>	51
<i>amicta</i> Reinhard, <i>Clairvillia</i>	24
<i>amicula</i> Mesnil, <i>Drino</i>	30
<i>amoena</i> Mesnil, <i>Elfriedella</i>	32
<i>amoena</i> Mesnil, <i>Erycilla</i>	33
<i>ampliforceps</i> Rowe, <i>Fabriciella</i>	37
<i>amplifrons</i> Brooks, <i>Pallasia</i>	57
<i>analabella</i> Thompson, <i>Winthemia</i>	77
<i>anassa</i> Reinhard, <i>Phorocera</i>	61
<i>anatolica</i> Mesnil, <i>Nilea</i>	55
<i>ancylostomiae</i> Thompson, <i>Carmodymyia</i>	22
<i>ancyrensis</i> Villeneuve, <i>Rhinotachina</i>	67
<i>andersoni</i> Guimarães, <i>Winthemia</i>	77
<i>angulata</i> Curran, <i>Peleteria</i>	59
<i>angulicornis</i> Villeneuve, <i>Carcelia</i>	21
<i>angulinea</i> Reinhard, <i>Politomyia</i>	64

<i>angustifacies</i> Mesnil, <i>Drino</i>	30
<i>angustifrons</i> Mesnil, <i>Crossocosmia jacobseni</i>	25
<i>angustifrons</i> Mesnil, <i>Ctenophorocera</i>	26
<i>angustipennis</i> Reinhard, <i>Aplomyiopsis</i>	14
<i>anorbitalis</i> Brooks, <i>Aubaeanaetia</i>	16
<i>anorbitalis</i> Mesnil, <i>Lydellina</i>	48
<i>antennalis</i> Mesnil, <i>Crocuta</i>	25
<i>antennalis</i> Thompson, <i>Maracasimyia</i>	48
<i>antennula</i> Villeneuve, <i>Voria</i>	76
<i>aperta</i> Mesnil, <i>Diaprochaeta</i>	28
<i>apicalis</i> Mesnil, <i>Lomatacantha</i>	47
<i>apicalis</i> Villeneuve, <i>Gonioclea</i>	40
<i>apicata</i> Curran, <i>Tachinomyia</i>	73
<i>apivorus</i> Villeneuve, <i>Rondaniooestrus</i>	68
<i>aplomyiodes</i> Villeneuve, <i>Exorista</i>	36
<i>aprifica</i> Villeneuve, <i>Phryno</i>	62
<i>arctica</i> Malloch, <i>Peleteria</i>	59
<i>argentata</i> Reinhard, <i>Actinodorria</i>	12
<i>argentea</i> Rowe, <i>Fabriciella</i>	37
<i>argentea</i> Thompson, <i>Actinodorria</i>	13
<i>argenteus</i> Thompson, <i>Heliodexodes</i>	41
<i>argentifrons</i> Brooks, <i>Alophorellopsis</i>	13
<i>argentosa</i> Reinhard, <i>Minthozelia</i>	51
<i>arguta</i> Reinhard, <i>Zizophomyia</i>	79
<i>arida</i> Reinhard, <i>Psilopleura</i>	67
<i>aristalis</i> Villeneuve, <i>Phytomyzoneura</i>	63
<i>aristatum</i> Mesnil, <i>Dexiosoma</i>	28
<i>arnaudi</i> Reinhard, <i>Phorocera</i>	62
<i>arrugans</i> Reinhard, <i>Echinopyrrhosia</i>	31
<i>ascita</i> Reinhard, <i>Microphthalmia</i>	50
<i>ater</i> Mesnil, <i>Gastropilops</i>	39
<i>atixcoensis</i> Reinhard, <i>Eutrapelus</i>	35
<i>atoma</i> Reinhard, <i>Aphantorhapha</i>	14
<i>atricans</i> Villeneuve, <i>Eutachina</i>	35
<i>atricans</i> Villeneuve, <i>Gonioclea</i>	41
<i>atricauda</i> Mesnil, <i>Argyrophylax</i>	16
<i>atricauda</i> Mesnil, <i>Blepharella</i>	18
<i>atriceps</i> Reinhard, <i>Chaetophlepsis</i>	24
<i>atrotincta</i> Reinhard, <i>Cistogaster</i>	24
<i>aturgida</i> Brooks, <i>Gonia</i>	39
<i>aurantiaca</i> Mesnil, <i>Istoglossa</i>	44
<i>auranticauda</i> Arnaud, <i>Metamyia</i>	50
<i>aurata</i> Mesnil, <i>Rhacodinella</i>	67
<i>aurea</i> Thompson, <i>Dexodioidea</i>	28
<i>aurea</i> Thompson, <i>Dexodiopsis</i>	28
<i>aurea</i> Thompson, <i>Ignotomyia</i>	44
<i>aurea</i> Thompson, <i>Trochilogglossa</i>	76
<i>aureoaustralis</i> Thompson, <i>Drino</i>	29
<i>aureocephala</i> Thompson, <i>Calocarcelia</i>	20
<i>aureola</i> Mesnil, <i>Drino</i>	31
<i>aureola</i> Mesnil, <i>Solieria</i>	70
<i>aureonigra</i> Thompson, <i>Wintheimia</i>	77
<i>auricaudata</i> Brooks, <i>Paraphorantha</i>	58
<i>auriceps</i> Reinhard, <i>Cyosoprocta</i>	27
<i>aurifacies</i> Guimarães, <i>Neocuphocera</i>	55
<i>aurifera</i> Villeneuve, <i>Sturmia</i>	72
<i>aurifrons</i> Curran, <i>Spathimeigenia</i>	71
<i>aurifrons</i> Thompson, <i>Anoxynops</i>	14
<i>aurora</i> Herting, <i>Cylindromyia</i>	27
<i>aurora</i> Mesnil, <i>Drino</i>	30
<i>australis</i> Reinhard, <i>Doryphorophaga</i>	29
<i>badiceps</i> Reinhard, <i>Admontia</i>	13
<i>bali</i> O'Hara, <i>Ceromyia</i>	23
<i>barbara</i> Mesnil, <i>Germaria</i>	39
<i>barbata</i> Mesnil, <i>Lypha</i>	48
<i>barbata</i> Villeneuve, <i>Calyptromyia</i>	20

<i>bellina</i> Mesnil, <i>Meigenia</i>	49
<i>bennetti</i> Thompson, <i>Archytas</i>	15
<i>benoisti</i> Mesnil, <i>Myothryia</i>	53
<i>bequaerti</i> Villeneuve, <i>Bogosia</i>	19
<i>bequaerti</i> Villeneuve, <i>Lissoglossa</i>	46
<i>bicolor</i> Curran, <i>Ginglimyia</i>	39
<i>bicoloripes</i> Mesnil, <i>Oswaldia</i>	56
<i>bicornis</i> Mesnil, <i>Lophosiosoma</i>	47
<i>bigelowi</i> Curran, <i>Grisdalemyia</i>	41
<i>bigelowi</i> Curran, <i>Gymnodexia</i>	41
<i>bisetosa</i> Roubaud & Villeneuve, <i>Roubaudia</i>	68
<i>bisetosa</i> Thompson, <i>Argyrophylax</i>	16
<i>blandoides</i> Thompson, <i>Eusisyropa</i>	35
<i>bogotensis</i> Reinhard, <i>Lindigepalpus</i>	45
<i>bohemica</i> Mesnil, <i>Drino</i>	30
<i>bonasus</i> Reinhard, <i>Ergolabus</i>	32
<i>borbonensis</i> Villeneuve, <i>Cadurcia</i>	19
<i>borealis</i> Brooks, <i>Ceromasia</i>	23
<i>boreata</i> Mesnil, <i>Siphona</i>	69
<i>boreotis</i> Reinhard, <i>Solieropsis</i>	70
<i>brevicornis</i> Brooks, <i>Metaplagia</i>	50
<i>brevicornis</i> Mesnil, <i>Thyellina</i>	75
<i>breviforceps</i> Tothill, <i>Gonia</i>	39
<i>brevipulvilli</i> Tothill, <i>Gonia</i>	39
<i>brevirostris</i> Tothill, <i>Fabriciella</i>	37
<i>brevisetata</i> Villeneuve, <i>Linnaemyia</i>	45
<i>brimleyi</i> Curran, <i>Homalactia</i>	42
<i>brimleyi</i> Reinhard, <i>Cryptomeigenia</i>	26
<i>brunnea</i> Brooks, <i>Psalidopteryx</i>	64
<i>brunnea</i> O'Hara, <i>Siphona</i>	69
<i>brunnescens</i> Villeneuve, <i>Actia</i>	11
<i>brunnicosa</i> Villeneuve, <i>Mormonomyia</i>	52
<i>brunnipalpis</i> Villeneuve, <i>Actia</i>	11
<i>bryanti</i> Curran, <i>Peleteria</i>	59
<i>bryanti</i> Curran, <i>Thelaira</i>	74
<i>buckelli</i> Curran, <i>Hylotomomyia</i>	43
<i>caesar</i> Aldrich, <i>Exorista</i>	36
<i>caffer</i> Villeneuve, <i>Myxarchiclops</i>	54
<i>caledoniae</i> Mesnil, <i>Winthemia</i>	77
<i>callani</i> Thompson, <i>Calodexia</i>	20
<i>callithrix</i> Reinhard, <i>Trichosaundersia</i>	75
<i>callitrix</i> Reinhard, <i>Jalapamyia</i>	44
<i>campestre</i> Curran, <i>Peleteria</i>	59
<i>campestris</i> Curran, <i>Mericia</i>	49
<i>canadensis</i> Brooks, <i>Pseudochaeta</i>	65
<i>canadensis</i> Brooks, <i>Rhodogyne</i>	67
<i>canadensis</i> Reinhard, <i>Myiopharus</i>	53
<i>canadensis</i> Tothill, <i>Fabriciella</i>	37
<i>cantans</i> Mesnil, <i>Exorista</i>	37
<i>carbonata</i> Mesnil, <i>Blepharella setigera</i>	18
<i>carbonata</i> Mesnil, <i>Crossocosmia</i>	25
<i>carnata</i> Reinhard, <i>Peleteria</i>	59
<i>caroniensis</i> Thompson, <i>Archytas</i>	15
<i>caudata</i> Villeneuve, <i>Akosempomyia</i>	13
<i>cepelaki</i> Mesnil, <i>Trichoparia</i>	75
<i>cerussata</i> Reinhard, <i>Spathidexia</i>	71
<i>chaetosa</i> Thompson, <i>Masphyoidea</i>	49
<i>chaoi</i> Mesnil, <i>Tachina</i>	73
<i>cib dela</i> Villeneuve, <i>Actia</i>	11
<i>cidaris</i> Reinhard, <i>Stolatosoma</i>	71
<i>cincta</i> Villeneuve, <i>Amesioclea</i>	13
<i>cinctella</i> Mesnil, <i>Lomatacantha</i>	47
<i>cinctella</i> Mesnil, <i>Phorinia</i>	61
<i>cinereola</i> Reinhard, <i>Technamyla</i>	74
<i>ciniscula</i> Reinhard, <i>Macromya</i>	48
<i>clara</i> Curran, <i>Peleteria</i>	59
<i>claripalpis</i> Thompson, <i>Neoxynopsoidae</i>	55
<i>claripalpis</i> Thompson, <i>Pseudoprosphrysa</i>	66
<i>claripalpis</i> Thompson, <i>Stenochaeta</i>	71
<i>claripalpis</i> Villeneuve, <i>Zambesa</i>	78
<i>claripennis</i> Mesnil, <i>Blepharella</i>	18
<i>claripennis</i> Reinhard, <i>Histochaeta</i>	42
<i>claripennis</i> Thompson, <i>Santacruziopsis</i>	68
<i>claripennis</i> Villeneuve, <i>Mormonomyia</i>	52
<i>claripennis</i> Villeneuve, <i>Proxystomima</i>	64
<i>clausa</i> Villeneuve, <i>Ceratochaeta</i>	23
<i>clavipalpis</i> Thompson, <i>Lespesia</i>	45
<i>clavipalpis</i> Thompson, <i>Pseudosturmia</i>	67
<i>clunalis</i> Reinhard, <i>Phorocera</i>	62
<i>clurina</i> Reinhard, <i>Pseudochaeta</i>	65
<i>cobala</i> Reinhard, <i>Mericia</i>	49
<i>coeruleonigra</i> Mesnil, <i>Ctenophorocera coerulea</i>	27
<i>coerulescens</i> Mesnil, <i>Thelairosova</i>	75
<i>colludens</i> Reinhard, <i>Spallanzania</i>	70
<i>columbia</i> Curran, <i>Hypostena</i>	43
<i>comitata</i> Villeneuve, <i>Actia</i>	11
<i>communis</i> Thompson, <i>Winthemia</i>	77
<i>comosa</i> Reinhard, <i>Masiphyomyia</i>	48
<i>compascua</i> Reinhard, <i>Pseudotachinomyia</i>	67
<i>compressa</i> Aldrich, <i>Cylindromyia</i>	27
<i>compressa</i> Mesnil, <i>Aphelogaster</i>	14
<i>condebensis</i> Reinhard, <i>Erynnia</i>	33
<i>confusa</i> Curran, <i>Peleteria</i>	60
<i>confusa</i> Mesnil, <i>Blepharella setigera</i>	18
<i>confusa</i> Mesnil, <i>Siphona</i>	69
<i>conica</i> Reinhard, <i>Dexodes</i>	28
<i>conjuncta</i> Curran, <i>Peleteria</i>	60
<i>connecta</i> Curran, <i>Lydella</i>	47
<i>consobrina</i> Villeneuve, <i>Linnaemyia</i>	46
<i>conspicendum</i> Cortés, <i>Spanipalpus</i>	71
<i>contristans</i> Villeneuve, <i>Pales</i>	57
<i>contumax</i> Brooks, <i>Gonia</i>	39
<i>convexa</i> Wood, <i>Phorocera</i>	62
<i>coracina</i> Reinhard, <i>Erynnia</i>	33
<i>cordiforceps</i> Rowe, <i>Fabriciella</i>	37
<i>cornigera</i> Curran, <i>Peleteria</i>	60
<i>cornualis</i> Reinhard, <i>Crocinosoma</i>	25
<i>cornuta</i> Curran, <i>Peleteria</i>	60
<i>cornuta</i> Curran, <i>Wagneria</i>	76
<i>cornuticaudata</i> Curran, <i>Peleteria</i>	60
<i>corsicana</i> Villeneuve, <i>Echinomyia magnicornis</i>	31
<i>crassipalpis</i> Reinhard, <i>Cryptomeigenia</i>	26
<i>crassipalpis</i> Villeneuve, <i>Phorocera</i>	62
<i>crassiseta</i> Villeneuve, <i>Plagiocoma</i>	63
<i>crassispinosa</i> Wood, <i>Policheta</i>	64
<i>crecentis</i> Reinhard, <i>Sturmia</i>	72
<i>cribrata</i> Villeneuve, <i>Ocyptera</i>	55
<i>cristata</i> Villeneuve, <i>Erigone</i>	32
<i>crocata</i> Reinhard, <i>Macromya</i>	48
<i>crossi</i> Villeneuve, <i>Stictodexia</i>	71
<i>cumutoensis</i> Thompson, <i>Euphorocera</i>	34
<i>curepei</i> Thompson, <i>Drino</i>	29
<i>curepei</i> Thompson, <i>Pseudochaeta</i>	65
<i>curepeiensis</i> Thompson, <i>Paranoxynops</i>	58
<i>curialis</i> Reinhard, <i>Clairvillia</i>	24
<i>curulis</i> Reinhard, <i>Pantagathus</i>	57
<i>cussilirix</i> Reinhard, <i>Euhystricia</i>	34
<i>dasymyia</i> O'Hara, <i>Actia</i>	11
<i>dasyops</i> Mesnil, <i>Bactromyia</i>	16
<i>dasyops</i> Villeneuve, <i>Sturmia</i>	72
<i>deceptor</i> Curran, <i>Clistomorpha</i>	25
<i>decorata</i> Villeneuve, <i>Hemilinnaemyia</i>	42

<i>decumata</i> Reinhard, <i>Hemyda</i>	42	<i>excisoides</i> Mesnil, <i>Calocarcelia</i>	20
<i>defecta</i> Reinhard, <i>Elassomyia</i>	32	<i>exigua</i> Wood, <i>Phorocera</i>	62
<i>delecta</i> Curran, <i>Erycia</i>	33	<i>exsecta</i> Villeneuve, <i>Actia</i>	11
<i>delicatula</i> Mesnil, <i>Bactromyia</i>	17	<i>faceta</i> Reinhard, <i>Plagiomima</i>	63
<i>delicatula</i> Mesnil, <i>Carcelia</i>	22	<i>facialis</i> Reinhard, <i>Metavoria</i>	50
<i>delusa</i> Reinhard, <i>Mastigiomyia</i>	49	<i>fallaciosa</i> Mesnil, <i>Blepharella</i>	18
<i>demissum</i> Reinhard, <i>Mochlosoma</i>	52	<i>fallax</i> Mesnil, <i>Lixophaga</i>	47
<i>depressa</i> Villeneuve, <i>Caducria</i>	20	<i>fallax</i> Reinhard, <i>Leucostoma</i>	45
<i>derisus</i> Reinhard, <i>Gremlinotrophus</i>	41	<i>fasciata</i> Curran, <i>Ernestia</i>	32
<i>deserta</i> Villeneuve, <i>Ocyptera</i>	56	<i>fasciata</i> Villeneuve, <i>Tachinomima multisetosa</i>	73
<i>dextella</i> Reinhard, <i>Lasioneura</i>	44	<i>fascipes</i> Villeneuve, <i>Sturmia</i>	72
<i>diaphana</i> Villeneuve, <i>Eupododexia</i>	34	<i>fasciventris</i> Curran, <i>Mericia</i>	49
<i>diffidens</i> Curran, <i>Actia</i>	11	<i>fastuosus</i> Villeneuve, <i>Xanthoestrus</i>	78
<i>dilaticornis</i> Mesnil, <i>Carcelia</i>	21	<i>fatigantis</i> Reinhard, <i>Metopomuscopteryx</i>	50
<i>dimorpha</i> O'Hara, <i>Actia</i>	11	<i>favilla</i> Reinhard, <i>Acemya</i>	11
<i>discalis</i> Brooks, <i>Catalinovoria</i>	22	<i>fenestrata</i> Mesnil, <i>Hypotachina</i>	43
<i>discalis</i> Mesnil, <i>Thelyconychia</i>	75	<i>festiva</i> Villeneuve, <i>Eupododexia</i>	34
<i>discalis</i> Morrison, <i>Gonia</i>	39	<i>fidelis</i> Reinhard, <i>Punaclista</i>	67
<i>discalis</i> Thompson, <i>Dexodimyia</i>	28	<i>filialis</i> Reinhard, <i>Cloacina</i>	25
<i>discolor</i> Villeneuve, <i>Helocera</i>	42	<i>fimbriata</i> Bischof, <i>Antistasea</i>	14
<i>dispar</i> Villeneuve, <i>Amphicestonia</i>	13	<i>finalis</i> Reinhard, <i>Pseudochaeta</i>	65
<i>dispar</i> Villeneuve, <i>Periprosena</i>	60	<i>finitina</i> Reinhard, <i>Mauromyia</i>	49
<i>disparis</i> Reinhard, <i>Thelymyia</i>	75	<i>fissiforceps</i> Tothill, <i>Gonia</i>	40
<i>dissimiloides</i> Thompson, <i>Archytas</i>	15	<i>flava</i> O'Hara, <i>Ceromya</i>	23
<i>dissita</i> Reinhard, <i>Paraphasmophaga</i>	58	<i>flavipalpis</i> Mesnil, <i>Compsiluroides</i>	25
<i>dissitus</i> Reinhard, <i>Eurylochus</i>	34	<i>flavipalpis</i> Thompson, <i>Pseudochaeta</i>	65
<i>distant</i> Villeneuve, <i>Prozenillia</i>	64	<i>flavipes</i> Reinhard, <i>Lispideosoma</i>	46
<i>diversipes</i> Villeneuve, <i>Xenolophosia</i>	78	<i>flavipes</i> Thompson, <i>Amazonoughia</i>	13
<i>dolopis</i> Reinhard, <i>Spathimeigenia</i>	71	<i>flavipes</i> Thompson, <i>Pelecotheca</i>	59
<i>dolosa</i> Reinhard, <i>Spathimeigenia</i>	71	<i>flavipes</i> Thompson, <i>Synorbitalia</i>	73
<i>downsi</i> Reinhard, <i>Colurus</i>	25	<i>flaviseta</i> Villeneuve, <i>Actia</i>	11
<i>dubia</i> Curran, <i>Admontia</i>	13	<i>flavogrisea</i> Villeneuve, <i>Subfischeria</i>	73
<i>dubia</i> Curran, <i>Cryptomeigenia</i>	26	<i>florentis</i> Reinhard, <i>Eubrachymera</i>	33
<i>dubiosa</i> Thompson, <i>Santacruzia</i>	68	<i>floridensis</i> Guimarães, <i>Winthemia</i>	77
<i>dubiosa</i> Thompson, <i>Winthemia</i>	77	<i>foliacea</i> Mesnil, <i>Blepharomyia</i>	19
<i>dumalis</i> Reinhard, <i>Phobetromyia</i>	61	<i>formosana</i> Villeneuve, <i>Hermyia</i>	42
<i>dumonti</i> Mesnil, <i>Linnaemyia</i>	46	<i>formosensis</i> Villeneuve, <i>Linnaemyia</i>	46
<i>duplare</i> Reinhard, <i>Mochlosoma</i>	52	<i>formosus</i> Reinhard, <i>Diotrephe</i>	28
<i>duplaria</i> Villeneuve, <i>Tachina</i>	73	<i>fracida</i> Reinhard, <i>Ginglymia</i>	39
<i>duplaris</i> Reinhard, <i>Nearchus</i>	54	<i>fracida</i> Reinhard, <i>Mactomyia</i>	48
<i>ebneri</i> Villeneuve, <i>Tachina</i>	73	<i>frontalis</i> Brooks, <i>Gymnocheta</i>	41
<i>eburneola</i> Villeneuve, <i>Linnaemyia</i>	46	<i>frontalis</i> Brooks, <i>Lypha</i>	48
<i>effetus</i> Reinhard, <i>Mimologus</i>	51	<i>frontalis</i> Mesnil, <i>Lydellina</i>	48
<i>efflatouni</i> Mesnil, <i>Siphona</i>	69	<i>frontalis</i> Reinhard, <i>Pseudochaeta</i>	65
<i>efflatouni</i> Villeneuve, <i>Alloprosopaea</i>	13	<i>frontosa</i> Say, <i>Gonia</i>	40
<i>effrenatum</i> Reinhard, <i>Leucostoma</i>	45	<i>fucosa</i> Mesnil, <i>Eurythia</i>	35
<i>egula</i> Reinhard, <i>Fabriciella</i>	37	<i>fulgens</i> Reinhard, <i>Tarpessita</i>	74
<i>electilis</i> Reinhard, <i>Helioptrosopa</i>	41	<i>fumator</i> Reinhard, <i>Canelomyia</i>	20
<i>elegans</i> Reinhard, <i>Minthodexiopsis</i>	51	<i>fumiferanae</i> Tothill, <i>Winthemia</i>	77
<i>elegans</i> Villeneuve, <i>Laufferiella</i>	44	<i>fumipennis</i> Brooks, <i>Lypha</i>	48
<i>elegantula</i> Mesnil, <i>Nemoraea</i>	54	<i>fumipennis</i> Thompson, <i>Cataphoriniopsis</i>	22
<i>emdeni</i> Mesnil, <i>Eurythia</i>	35	<i>fumipennis</i> Thompson, <i>Santacruziopsis</i>	68
<i>emdeni</i> Mesnil, <i>Rhodogyne</i>	67	<i>fumosa</i> Villeneuve, <i>Mormonomyia</i>	52
<i>emulatus</i> Reinhard, <i>Mimologus</i>	51	<i>fumosum</i> Villeneuve, <i>Thelairosoma</i>	75
<i>erebiae</i> Mesnil, <i>Blepharella</i>	18	<i>funebris</i> Reinhard, <i>Cyosoprocta</i>	27
<i>erilis</i> Reinhard, <i>Nimiocauda</i>	55	<i>funebris</i> Reinhard, <i>Gigamyiopsis</i>	39
<i>eronis</i> Curran, <i>Peleteria</i>	60	<i>furacis</i> Reinhard, <i>Dolichocodia</i>	28
<i>erratilis</i> Reinhard, <i>Dolichocodia</i>	28	<i>fuscicosta</i> Curran, <i>Phyllomya</i>	63
<i>erronis</i> Reinhard, <i>Spathimeigenia</i>	71	<i>fuscipennis</i> Mesnil, <i>Blepharella</i>	18
<i>erythrocera</i> Reinhard, <i>Myioclonia</i>	53	<i>fuscisquamis</i> Brooks, <i>Psalidopteryx</i>	64
<i>ethiae</i> Brooks, <i>Eufrontina</i>	34	<i>galbae</i> Thompson, <i>Cataphoriniopsis</i>	22
<i>euthes</i> Reinhard, <i>Plagiomima</i>	63	<i>geminia</i> Herting, <i>Trafoia</i>	75
<i>eurekana</i> Reinhard, <i>Fabriciella</i>	37	<i>genibarbis</i> Mesnil, <i>Platymyia nemestrina</i>	63
<i>evanida</i> Reinhard, <i>Fabriciella</i>	37	<i>genibarbis</i> Villeneuve, <i>Cavalieria</i>	22
<i>evexus</i> Reinhard, <i>Tarassophorus</i>	74	<i>genurufa</i> Villeneuve, <i>Echinomyia</i>	31
<i>excavata</i> Herting, <i>Besseria</i>	17	<i>ghanii</i> Mesnil, <i>Exorista</i>	36

<i>ghanii</i> Mesnil, <i>Sisyropa</i>	70	<i>iterata</i> Mesnil, <i>Drino</i>	30
<i>ghanii</i> Mesnil, <i>Tachinophytopsis</i>	74	<i>itoi</i> Mesnil, <i>Hyalurgus</i>	43
<i>ghanii</i> Mesnil, <i>Thelaira</i>	75	<i>jacentkowskyi</i> Mesnil, <i>Winthemia</i>	78
<i>ghesquierei</i> Villeneuve, <i>Syneplaca</i>	73	<i>japonica</i> Mesnil, <i>Actinochaetopteryx</i>	12
<i>gilpiniae</i> Mesnil, <i>Drino</i>	30	<i>japonica</i> Mesnil, <i>Bothria</i>	19
<i>glauca</i> Brooks, <i>Bonellimyia</i>	19	<i>japonica</i> Mesnil, <i>Ceranthia</i>	22
<i>globosa</i> Reinhard, <i>Nicetria</i>	55	<i>japonica</i> Mesnil, <i>Drino chatterjeeana</i>	30
<i>grandicornis</i> Mesnil, <i>Myxexoristops</i>	54	<i>japonica</i> Mesnil, <i>Nemosturmia</i>	54
<i>grandigena</i> Villeneuve, <i>Echinomyia</i>	31	<i>japonica</i> Mesnil, <i>Promedina</i>	64
<i>grandipulvilli</i> Morrison, <i>Gonia</i>	40	<i>jepsoni</i> Villeneuve, <i>Stroblomyia</i>	72
<i>grandis</i> Mesnil, <i>Eloceria delecta</i>	32	<i>jocosa</i> Villeneuve, <i>Actia</i>	12
<i>griphus</i> Reinhard, <i>Hegesinus</i>	41	<i>jocularis</i> Mesnil, <i>Actia</i>	12
<i>grisea</i> Mesnil, <i>Ctenophorinia</i>	26	<i>jorgensenii</i> O'Hara, <i>Frontiniella</i>	38
<i>grisea</i> Thompson, <i>Anoxynopsoidea</i>	14	<i>juniperi</i> O'Hara, <i>Baeomyia</i>	17
<i>grisea</i> Thompson, <i>Vivianoidea</i>	76	<i>katoi</i> Mesnil, <i>Phryno</i>	62
<i>griseola</i> Mesnil, <i>Siphona maculata</i>	69	<i>kingi</i> Brooks, <i>Dolichotarsus</i>	29
<i>griseus</i> Brooks, <i>Dolichotarsus</i>	29	<i>klapperichi</i> Mesnil, <i>Aphria</i>	14
<i>gymnops</i> Herting, <i>Clemelis</i>	25	<i>klapperichi</i> Mesnil, <i>Chaetoxorista</i>	23
<i>haematodes</i> Villeneuve, <i>Gaedioxenis</i>	39	<i>klapperichi</i> Mesnil, <i>Dolichocolon</i>	29
<i>halisidota</i> Brooks, <i>Leschenaultia</i>	44	<i>knowltoni</i> Reinhard, <i>Athanatus</i>	16
<i>hamilla</i> Reinhard, <i>Mericia</i>	50	<i>kolomyetzi</i> Mesnil, <i>Billaea</i>	17
<i>hamulata</i> Villeneuve, <i>Xenolophosia</i>	78	<i>kugleri</i> Mesnil, <i>Exorista</i>	36
<i>hariola</i> Reinhard, <i>Uramya</i>	76	<i>labeosus</i> Reinhard, <i>Parepalpus</i>	59
<i>harpeza</i> Reinhard, <i>Saundersiops</i>	69	<i>labradorensis</i> Brooks, <i>Petinops</i>	61
<i>harpi</i> Reinhard, <i>Myiophasia</i>	53	<i>lacteipennis</i> Mesnil, <i>Exorista</i>	36
<i>harrisi</i> Reinhard, <i>Exoristoides</i>	37	<i>lacteipennis</i> Villeneuve, <i>Phytomyptera</i>	63
<i>hedini</i> Villeneuve, <i>Echinomyia</i>	31	<i>laeta</i> Mesnil, <i>Nemosturmia</i>	54
<i>heringi</i> Mesnil, <i>Myxexoristops</i>	54	<i>laetabilis</i> Villeneuve, <i>Sturmia</i>	72
<i>hiemalis</i> Reinhard, <i>Muscopteryx</i>	53	<i>lampuris</i> Reinhard, <i>Juriniopsis</i>	44
<i>hiemalis</i> Thompson, <i>Archytas</i>	15	<i>languidula</i> Villeneuve, <i>Actia</i>	12
<i>hilaris</i> Reinhard, <i>Muscopteryx</i>	53	<i>lapponica</i> Wood, <i>Allophorocera</i>	13
<i>hilaris</i> Villeneuve, <i>Hypovoria</i>	44	<i>lasia</i> Reinhard, <i>Myiophasia</i>	53
<i>hirsuta</i> Mesnil, <i>Oswaldia</i>	56	<i>laterale</i> Curran, <i>Lydella</i>	47
<i>hirsuta</i> Mesnil, <i>Zambesomima</i>	78	<i>lathami</i> Curran, <i>Lydella</i>	47
<i>hirsutus</i> Reinhard, <i>Pacidianus</i>	57	<i>latianulum</i> Tothill, <i>Fabriciella</i>	37
<i>hirtipleura</i> Mesnil, <i>Graphogaster buccata</i>	41	<i>latifascia</i> Villeneuve, <i>Bogosia</i>	19
<i>hispanica</i> Mesnil, <i>Gaeda</i>	38	<i>latiforceps</i> Tothill, <i>Fabriciella</i>	37
<i>hispanica</i> Mesnil, <i>Germaria</i>	39	<i>latifrons</i> Jacentkovski, <i>Aporotachina</i>	15
<i>hispanica</i> Mesnil, <i>Lomacantha parra</i>	47	<i>latifrons</i> Mesnil, <i>Thecocarcelia</i>	74
<i>hispida</i> Tothill, <i>Fabriciella</i>	37	<i>latifrons</i> Reinhard, <i>Metavoria</i>	50
<i>hokkaidensis</i> Mesnil, <i>Siphona</i>	69	<i>latifrons</i> Tothill, <i>Fabriciella</i>	37
<i>hosпита</i> Reinhard, <i>Leschenaultia</i>	44	<i>latigena</i> Mesnil, <i>Crossocosmia</i>	26
<i>hoysi</i> Mesnil, <i>Hyleorus</i>	43	<i>latigena</i> Tothill, <i>Fabriciella</i>	38
<i>hyphantriae</i> Tothill, <i>Lydella</i>	47	<i>latigena</i> Villeneuve, <i>Cestonionerva</i>	23
<i>hystricoides</i> Williston, <i>Jurinia</i>	44	<i>latigena</i> Wood, <i>Ligeria</i>	45
<i>hystrix</i> Mesnil, <i>Macquartia</i>	48	<i>latimana</i> Villeneuve, <i>Aplomyia</i>	14
<i>illocale</i> Reinhard, <i>Mochlosoma</i>	52	<i>latimargo</i> Villeneuve, <i>Billaea</i>	17
<i>illustris</i> Mesnil, <i>Diaprochaeta</i>	28	<i>latipennis</i> Curran, <i>Hemyda</i>	42
<i>imitator</i> Curran, <i>Hypostena</i>	43	<i>latitarsus</i> Thompson, <i>Pseudochaeta</i>	65
<i>immissa</i> Reinhard, <i>Lydella</i>	47	<i>laudatum</i> Reinhard, <i>Mochlosoma</i>	52
<i>impatiens</i> Curran, <i>Hypostena</i>	43	<i>laxifrons</i> Reinhard, <i>Eutrixia</i>	35
<i>incanus</i> Reinhard, <i>Isidotus</i>	44	<i>lenis</i> Reinhard, <i>Tesseracephalus</i>	74
<i>incarcerata</i> O'Hara, <i>Frontiniella</i>	38	<i>leonidei</i> Mesnil, <i>Plesiooestrus</i>	64
<i>incompleta</i> Curran, <i>Besseria</i>	17	<i>lesnei</i> Villeneuve, <i>Stictodexia</i>	71
<i>incurva</i> Reinhard, <i>Metopomuscopteryx</i>	50	<i>leucodes</i> Villeneuve, <i>Mormonomyia</i>	52
<i>indica</i> Mesnil, <i>Acemyia</i>	11	<i>levata</i> Reinhard, <i>Erycia</i>	33
<i>indutile</i> Reinhard, <i>Mochlosoma</i>	52	<i>liciata</i> Reinhard, <i>Helioptrosopa</i>	42
<i>infida</i> Reinhard, <i>Evidomyia</i>	35	<i>lindensis</i> Reinhard, <i>Meledonus</i>	49
<i>infoederata</i> Villeneuve, <i>Nemoraea</i>	54	<i>livescens</i> Reinhard, <i>Dolichotarsus</i>	29
<i>ingrami</i> Mesnil, <i>Thelairosome</i>	75	<i>lodosi</i> Mesnil, <i>Blepharella</i>	18
<i>instabilis</i> Curran, <i>Archytas</i>	15	<i>longicornis</i> Mesnil, <i>Paratryphera</i>	58
<i>intermedia</i> Brooks, <i>Lypha</i>	48	<i>longicornis</i> Reinhard, <i>Masiphyomyia</i>	48
<i>inusta</i> Mesnil, <i>Carcelia</i>	21	<i>longiforceps</i> Brooks, <i>Pseudotachinomyia</i>	67
<i>inusta</i> Mesnil, <i>Drino inconspicua</i>	30	<i>longiforceps</i> Tothill, <i>Gonia</i>	40
<i>inusta</i> Mesnil, <i>Fausta</i>	38	<i>longipes</i> Thompson, <i>Cryptomeigenioidea</i>	26
<i>invulnerata</i> Reinhard, <i>Aulicomysia</i>	16	<i>longipes</i> Thompson, <i>Microleskia</i>	50

<i>longipulvilli</i> Tothill, <i>Gonia</i>	40	<i>minor</i> Curran, <i>Lydella</i>	47
<i>longirostris</i> Villeneuve, <i>Myobia</i>	53	<i>minor</i> Thompson, <i>Eumacrohoughia</i>	34
<i>longirostris</i> Villeneuve, <i>Phorocera</i>	62	<i>minor</i> Thompson, <i>Floradalia</i>	38
<i>longiseta</i> Reinhard, <i>Muscopteryx</i>	53	<i>minor</i> Thompson, <i>Palpexorista</i>	57
<i>longissima</i> O'Hara, <i>Siphona</i>	69	<i>minor</i> Villeneuve, <i>Epineura</i>	32
<i>longiunguis</i> Tothill, <i>Fabriciella</i>	38	<i>mira</i> Reinhard, <i>Minthozelia</i>	51
<i>loriola</i> Reinhard, <i>Siphokeskia</i>	69	<i>mirabilis</i> Mesnil, <i>Zenillia</i>	79
<i>loxostegeae</i> Reinhard, <i>Exorista</i>	36	<i>mirabilis</i> Thompson, <i>Mystacomyoidea</i>	54
<i>lucentis</i> Reinhard, <i>Xanthocera</i>	78	<i>miranda</i> Mesnil, <i>Leskia</i>	45
<i>lucidula</i> Villeneuve, <i>Carcelia</i>	21	<i>misella</i> Villeneuve, <i>Wagneria</i>	76
<i>luciflua</i> Villeneuve, <i>Ocyptera</i>	56	<i>monnula</i> Reinhard, <i>Gaediophana</i>	38
<i>lurida</i> Reinhard, <i>Siphona</i>	69	<i>monochaeta</i> Mesnil, <i>Phebellia</i>	61
<i>luteifacies</i> Mesnil, <i>Chloropales</i>	24	<i>montanum</i> Cortés, <i>Philocorus</i>	61
<i>lutescens</i> Villeneuve, <i>Oestrocharis</i>	56	<i>monticola</i> Mesnil, <i>Eurythia</i>	35
<i>luxatura</i> Reinhard, <i>Adoryphorophaga</i>	13	<i>monticola</i> Villeneuve, <i>Dexia</i>	28
<i>macdunnoughi</i> Brooks, <i>Psalidopteryx</i>	64	<i>montium</i> Villeneuve, <i>Actia</i>	12
<i>macella</i> Reinhard, <i>Doryphorophaga</i>	29	<i>multifaria</i> O'Hara, <i>Siphona</i>	70
<i>macrocephala</i> Mesnil, <i>Ctenophorocera</i>	27	<i>multipunctata</i> Villeneuve, <i>Pararondania</i>	58
<i>macrocera</i> Reinhard, <i>Helioptrosopa</i>	42	<i>multispinosa</i> Thompson, <i>Peletieria</i>	60
<i>macronymchia</i> Mesnil, <i>Ceromyia</i>	23	<i>murina</i> Mesnil, <i>Pales</i>	57
<i>maculiguama</i> Mesnil, <i>Ctenophorocera coerulea</i>	27	<i>muscoidea</i> Curran, <i>Cryptomeigenia</i>	26
<i>maculosus</i> Villeneuve, <i>Myxarchiclops</i>	54	<i>mutans</i> Mesnil, <i>Antistasea</i>	14
<i>madagascariensis</i> Villeneuve, <i>Pseudogonia</i>	66	<i>nana</i> Becker, <i>Gonia</i>	40
<i>madecassa</i> Mesnil, <i>Pseudoperichaeta</i>	66	<i>neglecta</i> Mesnil, <i>Blepharella</i>	18
<i>mageritensis</i> Mesnil, <i>Archiclops carthaginiensis</i>	15	<i>neotexensis</i> Brooks, <i>Peletieria</i>	60
<i>mageritensis</i> Villeneuve & Mesnil, <i>Salia</i>	68	<i>neptis</i> Reinhard, <i>Euceromasia</i>	33
<i>magna</i> Brooks, <i>Onychogonria</i>	56	<i>nexa</i> Reinhard, <i>Spathidexia</i>	71
<i>magnicornis</i> Mesnil, <i>Smidtia</i>	70	<i>nigella</i> Reinhard, <i>Fabriciella</i>	38
<i>major</i> Thompson, <i>Floradalia</i>	38	<i>nigeriensis</i> Villeneuve, <i>Microphthalmia europaea</i>	51
<i>major</i> Thompson, <i>Opsoleskia</i>	56	<i>nigra</i> Brooks, <i>Phylacteropoda</i>	63
<i>major</i> Villeneuve, <i>Myxarchiclops</i>	54	<i>nigra</i> Brooks, <i>Rhedia</i>	67
<i>majuscula</i> Mesnil, <i>Clemelis</i>	25	<i>nigra</i> Curran, <i>Eulasiona</i>	34
<i>malacosomae</i> Curran, <i>Frontina</i>	38	<i>nigra</i> Mesnil, <i>Blepharella</i>	18
<i>malitana</i> Reinhard, <i>Siphosturmia</i>	70	<i>nigrapex</i> Mesnil, <i>Carcelia</i>	21
<i>manitoba</i> Brooks, <i>Mericia</i>	50	<i>nigrescens</i> Curran, <i>Linnaemyia</i>	46
<i>maracasi</i> Thompson, <i>Celatoria</i>	22	<i>nigrescens</i> Mesnil, <i>Charitella</i>	24
<i>maracasi</i> Thompson, <i>Metaphorocera</i>	50	<i>nigrescens</i> Mesnil, <i>Nanoplagia</i>	54
<i>maracasi</i> Thompson, <i>Thelairodoriopsis</i>	75	<i>nigricans</i> Villeneuve, <i>Bucentes</i>	19
<i>marginalis</i> Curran, <i>Arctophyto</i>	16	<i>nigricauda</i> Mesnil, <i>Phebellia</i>	61
<i>marginalis</i> Reinhard, <i>Pseudochaeta</i>	65	<i>nigricauda</i> Thompson, <i>Drino</i>	30
<i>maroccana</i> Mesnil, <i>Drino latigena</i>	30	<i>nigricornis</i> Reinhard, <i>Myioclonia</i>	53
<i>martini</i> Reinhard, <i>Eleodiphaga</i>	32	<i>nigricornis</i> Reinhard, <i>Opsotheresia</i>	56
<i>mayeri</i> Mesnil, <i>Lubutana</i>	47	<i>nigricosta</i> Curran, <i>Gibsonomyia</i>	39
<i>mayneana</i> Villeneuve, <i>Sturmia</i>	72	<i>nigrifrons</i> Thompson, <i>Gopaulia</i>	41
<i>medeola</i> Reinhard, <i>Thematheca</i>	75	<i>nigrina</i> Mesnil, <i>Crossocosmia</i>	26
<i>medialis</i> O'Hara, <i>Siphona</i>	69	<i>nigripalpis</i> Reinhard, <i>Houghia</i>	42
<i>medicinensis</i> Strickland, <i>Evanalia</i>	35	<i>nigripalpis</i> Thompson, <i>Drino</i>	30
<i>melancholica</i> Mesnil, <i>Chaetovoria</i>	24	<i>nigripalpis</i> Thompson, <i>Eucelatarioidea</i>	33
<i>melancholica</i> Mesnil, <i>Drino</i>	31	<i>nigripalpis</i> Thompson, <i>Patelloa</i>	59
<i>melancholica</i> Mesnil, <i>Wardarina</i>	77	<i>nigripes</i> Curran, <i>Cryptomeigenia</i>	26
<i>mentitum</i> Reinhard, <i>Mochlosoma</i>	52	<i>nigripes</i> Mesnil, <i>Hygiella</i>	43
<i>meraculus</i> Reinhard, <i>Melanopalpus</i>	49	<i>nigripes</i> Thompson, <i>Calodexia</i>	20
<i>meralis</i> Reinhard, <i>Clastoneuriopsis</i>	25	<i>nigripilosa</i> Curran, <i>Cryptomeigenia</i>	26
<i>meridionalis</i> Thompson, <i>Palpoxorista</i>	57	<i>nigrita</i> Thompson, <i>Admontia</i>	13
<i>merista</i> Reinhard, <i>Paradidyma</i>	58	<i>nigrofimbriata</i> Villeneuve, <i>Mormonomomyia</i>	52
<i>mesnilii</i> Herting, <i>Paratryphera</i>	59	<i>nigrohalterata</i> Villeneuve, <i>Actia</i>	12
<i>mesnilii</i> Villeneuve, <i>Cestonoptera</i>	23	<i>nigropalpis</i> Tothill, <i>Ernestia</i>	32
<i>metallicus</i> Reinhard, <i>Minthozelia</i>	51	<i>nigroscutellata</i> Mesnil, <i>Sturmia</i>	72
<i>metallifica</i> Villeneuve, <i>Stomatomyia</i>	72	<i>ningulus</i> Reinhard, <i>Nearchus</i>	54
<i>micronychia</i> Mesnil, <i>Oswaldia</i>	57	<i>nipigonensis</i> Curran, <i>Billaea</i>	17
<i>micronychia</i> Mesnil, <i>Wagneria</i>	77	<i>nishijimai</i> Mesnil, <i>Strongygaster</i>	72
<i>micronyx</i> Mesnil, <i>Chaetoria</i>	24	<i>nitens</i> Mesnil, <i>Zophomyia</i>	79
<i>microps</i> Mesnil, <i>Baumhaueria</i>	17	<i>nitens</i> Thompson, <i>Pseudochaeta</i>	65
<i>mimela</i> Reinhard, <i>Epidexia</i>	32	<i>nitida</i> Curran, <i>Cyrtophloeba</i>	27
<i>mimica</i> Villeneuve, <i>Peleteria</i>	60	<i>nivalis</i> Curran, <i>Archytas</i>	15
<i>minima</i> Thompson, <i>Calocarcelia</i>	20	<i>nobilis</i> Mesnil, <i>Phyllomyia</i>	63

<i>noera</i> Reinhard, <i>Phorocera</i>	62	<i>pammelas</i> Reinhard, <i>Charasoma</i>	24
<i>notialis</i> Reinhard, <i>Chesippus</i>	24	<i>papuana</i> Mesnil, <i>Winthemia</i>	77
<i>noumeensis</i> Mesnil, <i>Carcelia</i>	21	<i>paralis</i> Reinhard, <i>Masiophyomyia</i>	49
<i>nova</i> Mesnil, <i>Bithia</i>	17	<i>parilis</i> Reinhard, <i>Muscopteryx</i>	53
<i>nova</i> Mesnil, <i>Drino</i>	31	<i>parva</i> Brooks, <i>Lypha</i>	48
<i>nova</i> Mesnil, <i>Dufouria</i>	31	<i>patelliforceps</i> Mesnil, <i>Exorista</i>	37
<i>novella</i> Mesnil, <i>Argyrophylax nova</i>	16	<i>patellipalpis</i> Mesnil, <i>Nemoraea</i>	54
<i>novemmaculata</i> Wood, <i>Steleoneura</i>	71	<i>patrita</i> Reinhard, <i>Doryphorophaga</i>	29
<i>nubecula</i> Reinhard, <i>Cephaloplagia</i>	22	<i>patruelis</i> Mesnil, <i>Drino</i>	31
<i>nuda</i> Brooks, <i>Psalidopteryx</i>	64	<i>pauciseta</i> Mesnil, <i>Ctenophorocera</i>	27
<i>nuda</i> Thompson, <i>Leschenaultia</i>	45	<i>pauciseta</i> Mesnil, <i>Murana</i>	53
<i>nudicauda</i> Mesnil, <i>Eucarcelia</i>	33	<i>pavonacea</i> Reinhard, <i>Siphoclytia</i>	69
<i>nudicosta</i> Mesnil, <i>Arrhinomyia</i>	16	<i>peccator</i> Reinhard, <i>Leucostoma</i>	45
<i>nudigena</i> Mesnil, <i>Loewia</i>	47	<i>pechumani</i> Reinhard, <i>Vibrissotheresia</i>	76
<i>nudigena</i> Mesnil, <i>Macquartia</i>	48	<i>pedalis</i> Reinhard, <i>Microphthalmia</i>	51
<i>nudigena</i> Villeneuve, <i>Ceratochaeta</i>	23	<i>pedia</i> Reinhard, <i>Viviania</i>	76
<i>nudinerva</i> Mesnil, <i>Suensonomyia</i>	73	<i>pellacis</i> Reinhard, <i>Echinopyrrhosia</i>	32
<i>nudioculata</i> Villeneuve, <i>Palesisa</i>	57	<i>pellecta</i> Reinhard, <i>Phorocera</i>	62
<i>numidica</i> Mesnil, <i>Exorista aberrans</i>	36	<i>pellucida</i> Mesnil, <i>Blepharella</i>	18
<i>numidica</i> Mesnil, <i>Microphthalmia europaea</i>	51	<i>peponis</i> Reinhard, <i>Nicetria</i>	55
<i>nuperus</i> Reinhard, <i>Meleterus</i>	49	<i>peraequalis</i> Mesnil, <i>Carcelia</i>	21
<i>oblectanea</i> Mesnil, <i>Carcelia</i>	21	<i>perdecora</i> Reinhard, <i>Pseudochaeta</i>	66
<i>oblimata</i> Mesnil, <i>Bessa</i>	17	<i>perdives</i> Villeneuve, <i>Exorista</i>	36
<i>oblimata</i> Mesnil, <i>Carcelia</i>	21	<i>perfida</i> Mesnil, <i>Blepharella</i>	18
<i>oblimata</i> Mesnil, <i>Clemelis pullata</i>	25	<i>peringueyi</i> Villeneuve, <i>Pleropeleteria</i>	63
<i>obliterata</i> Mesnil, <i>Carcelia</i>	21	<i>perrunar</i> Reinhard, <i>Leucostoma</i>	45
<i>obscura</i> Curran, <i>Lydella</i>	48	<i>persica</i> Mesnil, <i>Rhodogyne</i>	67
<i>obscura</i> Thompson, <i>Braziliomyia</i>	19	<i>persimilis</i> Reinhard, <i>Pacidianus</i>	57
<i>obscura</i> Thompson, <i>Sturmioptoidae</i>	73	<i>petentis</i> Reinhard, <i>Muscopteryx</i>	53
<i>obscurella</i> Mesnil, <i>Anacamptomyia pallida</i>	13	<i>petiolata</i> Thompson, <i>Pseudomasiphya</i>	66
<i>obscurior</i> Villeneuve, <i>Linnaemyia</i>	46	<i>phairi</i> Curran, <i>Peleteria</i>	60
<i>obsoleta</i> Curran, <i>Peleteria</i>	60	<i>phoenix</i> Villeneuve, <i>Compsoptesis</i>	25
<i>occidentale</i> Curran, <i>Gymnosoma</i>	41	<i>phyllophagae</i> Curran, <i>Microphthalmia</i>	51
<i>occidentalis</i> Brooks, <i>Gonia</i>	40	<i>physontae</i> Thompson, <i>Eucelatarioidea</i>	33
<i>occidentalis</i> Brooks, <i>Mericia</i>	50	<i>piarconensis</i> Thompson, <i>Archytas</i>	15
<i>occidentalis</i> Curran, <i>Tachinomyia</i>	74	<i>picta</i> Mesnil, <i>Blepharella</i>	18
<i>occidentalis</i> O'Hara, <i>Ceromya</i>	23	<i>picta</i> Villeneuve, <i>Pexomyia</i>	61
<i>ocellaris</i> Reinhard, <i>Wagneria</i>	76	<i>picticornis</i> Reinhard, <i>Paradmontia</i>	58
<i>ochreigaster</i> Curran, <i>Cryptomeigenia</i>	26	<i>pictilis</i> Reinhard, <i>Chaetona</i>	23
<i>oldi</i> Mesnil, <i>Blepharella</i>	18	<i>pictipennis</i> Reinhard, <i>Paraphasmophaga</i>	58
<i>oligomyia</i> O'Hara, <i>Siphona</i>	70	<i>pilabilis</i> Villeneuve, <i>Voria</i>	76
<i>oncoperae</i> Mesnil, <i>Platomyia</i>	63	<i>piliceps</i> Mesnil, <i>Loewia</i>	47
<i>ontario</i> Curran, <i>Actia</i>	12	<i>piligena</i> Mesnil, <i>Nemoraea</i>	54
<i>ontario</i> Curran, <i>Cryptomeigenia</i>	26	<i>piligena</i> Mesnil, <i>Pseudalsomyia</i>	65
<i>ontario</i> Curran, <i>Peteina</i>	61	<i>pilosa</i> Brooks, <i>Gonia</i>	40
<i>opaca</i> Reinhard, <i>Lixophaga</i>	47	<i>pilosa</i> Mesnil, <i>Pexopsis</i>	61
<i>optata</i> Reinhard, <i>Oedematocera</i>	56	<i>pilosa</i> Tothill, <i>Fabriciella</i>	38
<i>orbitalis</i> Reinhard, <i>Fabriciella</i>	38	<i>pilosa</i> Villeneuve, <i>Pseudoperichaeta</i>	66
<i>orbitalis</i> Reinhard, <i>Orphanotrophus</i>	56	<i>pisinnia</i> O'Hara, <i>Siphona</i>	70
<i>orbitalis</i> Villeneuve, <i>Winthemia</i>	77	<i>plagioderae</i> Mesnil, <i>Anthomyiopsis</i>	14
<i>oriens</i> O'Hara, <i>Ceromya</i>	23	<i>planicosta</i> Reinhard, <i>Nimioglossa</i>	55
<i>orientalis</i> Villeneuve, <i>Prodemoticus</i>	64	<i>plauasilis</i> Reinhard, <i>Gueriniopsis</i>	41
<i>orilla</i> Curran, <i>Pelatachina</i>	59	<i>politia</i> Brooks, <i>Chaetophlepis</i>	24
<i>ornata</i> Reinhard, <i>Collatia</i>	25	<i>politia</i> Brooks, <i>Eutrigocena</i>	35
<i>otiorynchi</i> Villeneuve, <i>Pandelleia</i>	57	<i>polittifrons</i> Reinhard, <i>Leucostoma</i>	45
<i>ouelleti</i> Curran, <i>Hypostena</i>	43	<i>politura</i> Reinhard, <i>Siphophyto</i>	70
<i>ovipara</i> Thompson, <i>Montserratia</i>	52	<i>pollinosa</i> Brooks, <i>Psalidopteryx</i>	64
<i>pacata</i> Reinhard, <i>Wagneria</i>	76	<i>pollinosa</i> Curran, <i>Admontia</i>	13
<i>pachychaeta</i> Villeneuve, <i>Ceratochaeta</i>	23	<i>pollinosa</i> Mesnil, <i>Carcelia</i>	21
<i>paligera</i> Mesnil, <i>Exorista</i>	36	<i>pollinosa</i> Thompson, <i>Lespisia</i>	45
<i>pallens</i> Reinhard, <i>Trypheromyia</i>	76	<i>pollinosa</i> Thompson, <i>Talparomyia</i>	74
<i>pallida</i> Roubaud & Villeneuve, <i>Roubaudia rufescens</i>	68	<i>pollinosa</i> Thompson, <i>Winthemia</i>	78
<i>pallida</i> Villeneuve, <i>Paratamiclea</i>	58	<i>pollinosa</i> Villeneuve, <i>Dexia</i>	28
<i>pallidula</i> Mesnil, <i>Ctenophorocera</i>	27	<i>portentosa</i> Mesnil, <i>Paragonia</i>	58
<i>pallipes</i> Reinhard, <i>Panacemyia</i>	57	<i>pruinosa</i> Roubaud & Villeneuve, <i>Roubaudia</i>	68
<i>palpalis</i> Reinhard, <i>Oberomyia</i>	55	<i>pschorni</i> Mesnil, <i>Pandelleia</i>	57

<i>pseudofallax</i> Villeneuve, <i>Tachina fallax</i>	73	<i>rutilans</i> Mesnil, <i>Exorista</i>	36
<i>pseudonuda</i> Brooks, <i>Psalidopteryx</i>	65	<i>rutilans</i> Villeneuve, <i>Phorostoma</i>	62
<i>psilocorsiphaga</i> Brooks, <i>Psalidopteryx</i>	65	<i>sagax</i> Townsend, <i>Gonia</i>	40
<i>pubiseta</i> Mesnil, <i>Dexiotrix</i>	28	<i>saginata</i> Reinhard, <i>Angustiopsis</i>	14
<i>pulchella</i> Villeneuve, <i>Linnaemyia</i>	46	<i>samarensis</i> Villeneuve, <i>Actia</i>	12
<i>pulchricornis</i> Villeneuve, <i>Histochaeta</i>	42	<i>sanctaecrucis</i> Thompson, <i>Archytas</i>	15
<i>pulla</i> Reinhard, <i>Aphantorhapha</i>	14	<i>sanctaecrucis</i> Thompson, <i>Leskiopalpus</i>	45
<i>pumilis</i> Reinhard, <i>Trismegistus</i>	75	<i>santacruzi</i> Thompson, <i>Pseudodrosophrysa</i>	66
<i>pupilla</i> Villeneuve, <i>Sarrorrhina</i>	68	<i>sartura</i> Reinhard, <i>Dexodes</i>	28
<i>purpurea</i> Thompson, <i>Bombyliomyia</i>	19	<i>scapularis</i> Villeneuve, <i>Ocyptera</i>	56
<i>purseglovei</i> Thompson, <i>Archytas</i>	15	<i>schineri</i> Mesnil, <i>Blepharipoda</i>	18
<i>pyrenaica</i> Mesnil, <i>Periscepsia</i>	61	<i>schistacea</i> Mesnil, <i>Chlorolydella</i>	24
<i>pyrenaica</i> Villeneuve, <i>Trixia</i>	75	<i>schnabli</i> Villeneuve, <i>Hemimasicera</i>	42
<i>radialis</i> O'Hara, <i>Actia</i>	12	<i>scintilla</i> Reinhard, <i>Lixophaga</i>	47
<i>raoi</i> Mesnil, <i>Hypotachina</i>	44	<i>scissilis</i> Reinhard, <i>Pseudolomyia</i>	66
<i>raoi</i> Mesnil, <i>Spoggosia</i>	71	<i>scordalus</i> Reinhard, <i>Milonius</i>	51
<i>rasa</i> Mesnil, <i>Pexopsis</i>	61	<i>scutellata</i> Mesnil, <i>Takanomyia</i>	74
<i>rasilis</i> Reinhard, <i>Spathidexia</i>	71	<i>sectilis</i> Reinhard, <i>Erycia</i>	33
<i>ravida</i> Reinhard, <i>Nimioglossa</i>	55	<i>securis</i> Reinhard, <i>Myiopharus</i>	53
<i>recincta</i> Reinhard, <i>Paradidyma</i>	58	<i>secutoris</i> Reinhard, <i>Adoryphorophaga</i>	13
<i>regalis</i> Villeneuve, <i>Deuteroclara</i>	28	<i>sedula</i> Reinhard, <i>Doryphorophaga</i>	29
<i>reinhardi</i> Brooks, <i>Gonia</i>	40	<i>segestris</i> Reinhard, <i>Eomeigenielloides</i>	32
<i>remora</i> Reinhard, <i>Lixophaga</i>	47	<i>sellersi</i> Thompson, <i>Myxexoristops</i>	54
<i>residis</i> Reinhard, <i>Charasoma</i>	24	<i>semiothisae</i> Brooks, <i>Chaetophlepsis</i>	24
<i>retroflexa</i> Villeneuve, <i>Ocyptera</i>	56	<i>semitestacea</i> Villeneuve, <i>Sturmia</i>	72
<i>rhodesiana</i> Villeneuve, <i>Minthodes</i>	51	<i>septentrionalis</i> Curran, <i>Meriania</i>	49
<i>riederi</i> Reinhard, <i>Cacozelus</i>	19	<i>serotina</i> Reinhard, <i>Stenoneura</i>	71
<i>ringdahli</i> Villeneuve, <i>Billaea</i>	17	<i>serotoninum</i> Reinhard, <i>Mochlosoma</i>	52
<i>rivalis</i> Reinhard, <i>Xiphomyia</i>	78	<i>sesamiae</i> Mesnil, <i>Descampsina</i>	27
<i>robusta</i> Brooks, <i>Dichocera</i>	28	<i>sesamiae</i> Mesnil, <i>Metagonistylum</i>	50
<i>robusta</i> Brooks, <i>Gonia</i>	40	<i>seticincta</i> Mesnil, <i>Dexia</i>	28
<i>robusta</i> Brooks, <i>Hyalomyiopsis</i>	43	<i>setifacies</i> Brooks, <i>Euthera</i>	35
<i>robusta</i> Mesnil, <i>Thecocarcelia</i>	74	<i>setifacies</i> Brooks, <i>Rhedia</i>	67
<i>robusta</i> Thompson, <i>Pseudocelatoria</i>	65	<i>setifacies</i> Reinhard, <i>Microchaetina</i>	50
<i>roderi</i> Villeneuve, <i>Peyritschia</i>	61	<i>setifrons</i> Brooks, <i>Okanagania</i>	56
<i>rogalis</i> Reinhard, <i>Pseudapinops</i>	65	<i>setifrons</i> Villeneuve, <i>Gaedionexis</i>	39
<i>rondaniana</i> Villeneuve, <i>Salia</i>	68	<i>setinerva</i> Mesnil, <i>Anaeogmena</i>	14
<i>rostrata</i> Tothill, <i>Fabriciella</i>	38	<i>setosa</i> Villeneuve, <i>Anurophylla</i>	14
<i>rubens</i> Villeneuve, <i>Epineura</i>	32	<i>sexualis</i> Curran, <i>Laximasicera</i>	44
<i>rubentis</i> Reinhard, <i>Eugaediopsis</i>	34	<i>shewelli</i> Cortés, <i>Enchomyia</i>	32
<i>rubida</i> Mesnil, <i>Aphria</i>	14	<i>sigilla</i> Reinhard, <i>Myiophasia</i>	53
<i>rubidiapex</i> Reinhard, <i>Hypenomyia</i>	43	<i>silvarum</i> Herting, <i>Siphona</i>	70
<i>rubiginans</i> Villeneuve, <i>Weberia</i>	77	<i>similis</i> Thompson, <i>Pseudodrosophrysa</i>	66
<i>rubrapex</i> Mesnil, <i>Zeuxia</i>	79	<i>similis</i> Villeneuve, <i>Bogosia</i>	19
<i>rubrapex</i> Villeneuve, <i>Echinomyia</i>	31	<i>siminina</i> Reinhard, <i>Pseudochaeta</i>	66
<i>rubricornis</i> Thompson, <i>Pseudorrhinactia</i>	66	<i>simiolus</i> Reinhard, <i>Sitellitergus</i>	70
<i>rubricose</i> Villeneuve, <i>Sturmia</i>	72	<i>simmondsi</i> Thompson, <i>Eugaediopsis</i>	34
<i>rubripes</i> Thompson, <i>Calodexia</i>	20	<i>simonyi</i> Villeneuve, <i>Goniophthalmus</i>	41
<i>rufescens</i> Villeneuve, <i>Phytomyptera</i>	63	<i>simplex</i> Curran, <i>Cryptomeigenia</i>	26
<i>rufescens</i> Villeneuve, <i>Roubaudia</i>	68	<i>sinaica</i> Villeneuve, <i>Plagia hilfii</i>	63
<i>ruficauda</i> Mesnil, <i>Blepharella setigera</i>	18	<i>siphonina</i> Villeneuve, <i>Myiobia</i>	53
<i>ruficauda</i> Reinhard, <i>Minthozelia</i>	51	<i>smithi</i> Brooks, <i>Gonia</i>	40
<i>ruficauda</i> Thompson, <i>Drino</i>	30	<i>sobrina</i> Reinhard, <i>Euceromasia</i>	33
<i>rufimana</i> Villeneuve, <i>Ocyptera</i>	56	<i>solata</i> Reinhard, <i>Euceromasia</i>	34
<i>rufipalpis</i> Brooks, <i>Gymnocheta</i>	41	<i>solitaris</i> Thompson, <i>Paradrino</i>	58
<i>rufipalpis</i> Villeneuve, <i>Cadurciella</i>	20	<i>sonorensis</i> O'Hara, <i>Baeomyia</i>	17
<i>rufipes</i> Brooks, <i>Argentoepalpus</i>	16	<i>soporis</i> Reinhard, <i>Celotrophus</i>	22
<i>rufipes</i> Villeneuve, <i>Alsomyia</i>	13	<i>soror</i> Mesnil, <i>Chaetogena</i>	23
<i>rufiventris</i> Curran, <i>Archytas</i>	15	<i>sororcula</i> Reinhard, <i>Catemophrys</i>	22
<i>rufiventris</i> Mesnil, <i>Dexiotrix</i>	28	<i>sororcula</i> Villeneuve, <i>Linnaemyia</i>	46
<i>rufiventris</i> Reinhard, <i>Siphosturmiosis</i>	70	<i>spathulatus</i> Mesnil, <i>Hyalurgus</i>	43
<i>rufochaeta</i> Curran, <i>Admontia</i>	13	<i>speciosissima</i> Mesnil, <i>Linnaemyia</i>	46
<i>rufula</i> Villeneuve, <i>Compsoptesis</i>	25	<i>speciosissima</i> Mesnil, <i>Winthemia</i>	78
<i>rungsi</i> Mesnil, <i>Atylomyia</i>	16	<i>spectabilis</i> Aldrich, <i>Frontina</i>	38
<i>rungsi</i> Mesnil, <i>Rhodogyne</i>	68	<i>spineiventer</i> Tothill, <i>Fabriciella</i>	38
<i>russulum</i> Reinhard, <i>Mochlosoma</i>	52	<i>spinosa</i> Tothill, <i>Fabriciella</i>	38

<i>splendidula</i> Villeneuve, <i>Echinomyia</i>	31
<i>starkei</i> Mesnil, <i>Actia</i>	12
<i>stenommatum</i> Wood, <i>Masistylum</i>	49
<i>sternalis</i> O'Hara, <i>Actia</i>	12
<i>sternalis</i> Reinhard, <i>Cylindromyia</i>	27
<i>stilla</i> Reinhard, <i>Achaetoneura</i>	11
<i>stipatus</i> Reinhard, <i>Dolichotarsus</i>	29
<i>stolidia</i> Reinhard, <i>Phorocera</i>	62
<i>subdistincta</i> Villeneuve, <i>Catagonia</i>	22
<i>subita</i> Reinhard, <i>Hypertophomma</i>	43
<i>subnitens</i> Reinhard, <i>Hypenomyia</i>	43
<i>subolis</i> Reinhard, <i>Charasoma</i>	24
<i>subpolita</i> Brooks, <i>Bonellimyia</i>	19
<i>sufferta</i> Villeneuve, <i>Actia</i>	12
<i>sulfurea</i> Mesnil, <i>Ceranthia</i>	22
<i>summatis</i> Reinhard, <i>Calolydella</i>	20
<i>syngamiae</i> Thompson, <i>Pseudochaeta</i>	66
<i>syriaca</i> Mesnil, <i>Exorista aberrans</i>	36
<i>tabida</i> Reinhard, <i>Imaguncula</i>	44
<i>taeniata</i> Villeneuve, <i>Lissoglossa</i>	46
<i>takanoi</i> Mesnil, <i>Calocarcelia</i>	20
<i>takanoi</i> Mesnil, <i>Dolichopodomintho</i>	29
<i>takanoi</i> Mesnil, <i>Linnaemyia</i>	46
<i>takanoi</i> Mesnil, <i>Paratrixa</i>	58
<i>takanoi</i> Mesnil, <i>Parerigone</i>	59
<i>takanoi</i> Mesnil, <i>Phorostoma</i>	62
<i>takanoi</i> Mesnil, <i>Phyllomyia</i>	63
<i>takanoi</i> Mesnil, <i>Scaphimyia</i>	69
<i>takanoi</i> Mesnil, <i>Steinomyia</i>	71
<i>takaoi</i> Mesnil, <i>Phorocera agilis</i>	62
<i>teliosis</i> Reinhard, <i>Chaetophlepsis</i>	24
<i>tenuiforceps</i> Morrison, <i>Gonia</i>	40
<i>tenuiforceps</i> Reinhard, <i>Gymnocarcelia</i>	41
<i>tenuipalpis</i> Villeneuve, <i>Actia</i>	12
<i>tenuirostris</i> Reinhard, <i>Philocalia</i>	61
<i>tessellans</i> Mesnil, <i>Ctenophorocera</i>	27
<i>tessellata</i> Brooks, <i>Linnaemyia</i>	46
<i>teutonia</i> Sabrosky, <i>Eucelatoria</i>	33
<i>texensis</i> Curran, <i>Peleteria</i>	60
<i>thessa</i> Reinhard, <i>Doliolomyia</i>	29
<i>thomasi</i> Mesnil, <i>Exorista</i>	36
<i>thompsoni</i> Guimarães, <i>Cylindromyia</i>	27
<i>tibialis</i> Curran, <i>Eulasiona</i>	34
<i>torta</i> Reinhard, <i>Peleteria</i>	60
<i>tothilli</i> Curran, <i>Hypostena</i>	43
<i>transcaspica</i> Villeneuve, <i>Hyperectinea</i>	43
<i>triangularis</i> Curran, <i>Cryptomeigenia</i>	26
<i>triangularis</i> Curran, <i>Mericia</i>	50
<i>triangularis</i> Thompson, <i>Phasiopsis</i>	61
<i>triangulifera</i> Thompson, <i>Argyrophylax</i>	16
<i>trichops</i> Herting, <i>Thecocarcelia</i>	74
<i>tricincta</i> Mesnil, <i>Elodimyia</i>	32
<i>tricincta</i> Reinhard, <i>Erynnia</i>	33
<i>triconis</i> Reinhard, <i>Messioniomyia</i>	50
<i>trigonata</i> Villeneuve, <i>Echinomyia</i>	31
<i>trinidadensis</i> Thompson, <i>Drino</i>	30
<i>trinidadensis</i> Thompson, <i>Neominthoidea</i>	55
<i>trinidadensis</i> Thompson, <i>Pelecotheca</i>	59
<i>trinitatis</i> Thompson, <i>Antillicola</i>	14
<i>trinitatis</i> Thompson, <i>Archytas</i>	15
<i>trinitatis</i> Thompson, <i>Calocarcelia</i>	20
<i>trinitatis</i> Thompson, <i>Cubaemyiopsis</i>	27
<i>trinitatis</i> Thompson, <i>Cuphocera</i>	27
<i>trinitatis</i> Thompson, <i>Eutelothryia</i>	35
<i>trinitatis</i> Thompson, <i>Neotrafoia</i>	55
<i>trinitatis</i> Thompson, <i>Panacemyia</i>	57
<i>trinitatis</i> Thompson, <i>Paraphasiopsis</i>	58
<i>trinitatis</i> Thompson, <i>Peleteria</i>	60
<i>trinitatis</i> Thompson, <i>Phasiopsis</i>	61
<i>trinitatis</i> Thompson, <i>Plagiprospherysa</i>	63
<i>trinitatis</i> Thompson, <i>Pseudochaeta</i>	66
<i>trinitatis</i> Thompson, <i>Sturmellina</i>	73
<i>trinitatis</i> Thompson, <i>Tamanamyia</i>	74
<i>trinitatis</i> Thompson, <i>Triodontopyga</i>	75
<i>trinitatis</i> Thompson, <i>Uromacquartia</i>	76
<i>trinitatis</i> Thompson, <i>Winthemia</i>	78
<i>triserilla</i> Villeneuve, <i>Exorista</i>	36
<i>triseta</i> Brooks, <i>Rileymyia</i>	68
<i>trispina</i> Villeneuve, <i>Hemiphania</i>	42
<i>triste</i> Reinhard, <i>Lispideosoma</i>	46
<i>trivittata</i> Curran, <i>Eutheresia</i>	35
<i>trivittatus</i> Reinhard, <i>Xanthomelanodes</i>	78
<i>trochanterata</i> Mesnil, <i>Billaea</i>	17
<i>trossulus</i> Reinhard, <i>Telemus</i>	74
<i>trudis</i> Reinhard, <i>Guerinia</i>	41
<i>tuchucheensis</i> Thompson, <i>Chaetona</i>	24
<i>umbrinervis</i> Mesnil, <i>Campylochaeta</i>	20
<i>umbrinervis</i> Mesnil, <i>Eriothrix</i>	32
<i>umbrinervis</i> Villeneuve, <i>Wagneria</i>	76
<i>umbrinervis</i> Villeneuve, <i>Xystomima</i>	78
<i>umbripennis</i> Mesnil, <i>Lydellina</i>	48
<i>umbrosa</i> Villeneuve, <i>Mormonomomyia</i>	52
<i>uncinata</i> Mesnil, <i>Meigenia</i>	49
<i>unicolor</i> Brooks, <i>Procistogaster</i>	64
<i>valida</i> Curran, <i>Dexodes</i>	28
<i>valida</i> Curran, <i>Peleteria</i>	60
<i>vallicola</i> Reinhard, <i>Orasturmia</i>	56
<i>vapulare</i> Reinhard, <i>Leucostoma</i>	45
<i>varia</i> Curran, <i>Linnaemyia</i>	46
<i>varians</i> Villeneuve, <i>Stomina caliendrata</i>	72
<i>variata</i> Curran, <i>Tachinomyia</i>	74
<i>varicornis</i> Curran, <i>Uclesia</i>	76
<i>varifrons</i> Curran, <i>Erycia</i>	33
<i>veleda</i> Reinhard, <i>Helioptrosopa</i>	42
<i>veniseta</i> Mesnil, <i>Eurysthaea</i>	35
<i>venusta</i> Reinhard, <i>Phaenopsis</i>	61
<i>venustoides</i> Mesnil, <i>Winthemia</i>	78
<i>verbeki</i> Mesnil, <i>Rhodogyne</i>	68
<i>versatilis</i> Villeneuve, <i>Sturmia</i>	72
<i>verticalis</i> Reinhard, <i>Exoristoides</i>	37
<i>verticalis</i> Reinhard, <i>Panacemyia</i>	57
<i>vetula</i> Reinhard, <i>Acroglossa</i>	11
<i>vicina</i> Mesnil, <i>Campogaster</i>	20
<i>vicinalis</i> Reinhard, <i>Euptilopareia</i>	34
<i>vicinum</i> Mesnil, <i>Dolichocolon</i>	29
<i>villeneuvei</i> Mesnil, <i>Eucarcelia</i>	33
<i>vinnula</i> Reinhard, <i>Eugaediopsis</i>	34
<i>viridis</i> Reinhard, <i>Elfia</i>	32
<i>vitripennis</i> Mesnil, <i>Billaea</i>	17
<i>vittata</i> Thompson, <i>Atactosturmia</i>	16
<i>vulgaris</i> Curran, <i>Archytas</i>	15
<i>walleyi</i> Brooks, <i>Phytomyptera</i>	63
<i>westi</i> Wilder, <i>Ptilodexia</i>	67
<i>wheeleri</i> Reinhard, <i>Cowania</i>	25
<i>xanthaspis</i> Mesnil, <i>Blepharella</i>	18
<i>xanthogaster</i> O'Hara, <i>Baeomyia</i>	17
<i>xylota</i> Curran, <i>Dexodes</i>	28
<i>yukonensis</i> Tothill, <i>Gonia</i>	40
<i>zernyi</i> Mesnil, <i>Zeuxia</i>	79
<i>zonalis</i> Curran, <i>Uclesia</i>	76