

OPTIMAL CONTROL OF A NUTRIENT-PHYTOPLANKTON-ZOOPLANKTON-FISH SYSTEM*

MARCUS R. GARVIE[†] AND CATALIN TRENCHEA[‡]

Abstract. We consider the mathematical formulation, analysis, and numerical solution of an optimal control problem for a nonlinear “nutrient-phytoplankton-zooplankton-fish” reaction-diffusion system. We study the existence of optimal solutions, derive an optimality system, and determine optimal solutions. In the original spatially homogeneous formulation [M. Scheffer, *Oikos*, 62 (1991), pp. 271–282] the dynamics of plankton were investigated as a function of parameters for nutrient levels and fish predation rate on zooplankton. In our paper the model is spatially extended and the parameter for fish predation treated as a multiplicative control variable. The model has implications for the biomanipulation of food-webs in eutrophic lakes to help improve water quality. In order to illustrate the control of irregular spatiotemporal dynamics of plankton in the model we implement a semi-implicit (in time) finite element method with “mass lumping” and present the results of numerical experiments in two space dimensions.

Key words. chaos, optimal control, biomanipulation, predator-prey interaction, finite element method

AMS subject classifications. 49K20, 92D25, 35K57

DOI. 10.1137/050645415

1. Introduction.

1.1. Model equations. In this paper we study the following nutrient-phytoplankton-zooplankton-fish reaction-diffusion system:

$$(1.1) \quad \begin{cases} \frac{\partial A}{\partial \tau} = D_A \Delta A + \hat{r} \frac{n}{n + h_n} A - cA^2 - pZ \frac{A}{A + h_a}, \\ \frac{\partial Z}{\partial \tau} = D_Z \Delta Z + peZ \frac{A}{A + h_a} - \hat{m}Z - F \frac{Z^2}{Z^2 + h_z^2}, \end{cases}$$

where A is phytoplankton biomass, Z is zooplankton biomass, D_a and D_z are the diffusion coefficients of phytoplankton and zooplankton, respectively, and n is the nutrient level of the system. F is the rate of zooplankton biomass consumed by fish per unit volume of water per day (average predation rate times the density of fish). It is important to note that in our formulation $F \equiv F(x, t)$, i.e., F is defined at every point in the lake and at every point in time. For definitions of the positive parameters c , e , \hat{m} , \hat{r} , p , h_a , h_z , and h_n , see [39]. The symbol τ denotes time in days, and $\Delta = \sum_{i=1}^d \partial^2 / \partial X_i^2$ is the usual Laplacian operator in $d = 2$ or 3 space dimensions. The grazing rate of zooplankton on phytoplankton is of a type II functional response, while the predation rate of fish on zooplankton is of type III [17]. In the absence of zooplankton the phytoplankton are assumed to grow logistically.

*Received by the editors November 16, 2005; accepted for publication (in revised form) November 6, 2006; published electronically May 29, 2007.

<http://www.siam.org/journals/sicon/46-3/64541.html>

[†]Department of Mathematics and Statistics, University of Guelph, Guelph, ON N1G 2W1, Canada (mgarvie@uoguelph.ca).

[‡]Department of Mathematics, University of Pittsburgh, Pittsburgh, PA 15260 (trenchea@pitt.edu).




FIG. 1.1. Interactions incorporated into a nutrient-plankton-fish model. Arrows indicate positive effects, circles indicate negative effects (redrawn from [39]).

The nutrient-phytoplankton-zooplankton-fish model was originally formulated as a system of ordinary differential equations (ODEs) by Scheffer [39] and has since been spatially extended (see, for example, [32, 29, 30, 33, 34, 45]).

The reaction-diffusion system models a simple food-web in lakes where planktivorous fish feed on zooplankton, and the zooplankton feed on phytoplankton (algae). The basic interactions in the model are illustrated in Figure 1.1. The model is “minimal” in the sense that only a few important interactions are taken into account. For example, the positive effect that fish have on the nutrients of the system is omitted in this model (indicated by the dotted arrow in Figure 1.1). Nevertheless, such minimal predator-prey systems display a wide range of ecologically relevant behavior, for example, spiral waves [34], target waves [41], diffusion-induced instability [30], and chaos [36, 34]. See [31] for a historical overview of modeling plankton dynamics and pattern formation mechanisms.

It is simpler to work with equations that have been scaled to nondimensional form; thus after letting $N := n/(n + h_n)$ in (1.1) we define dimensionless phytoplankton densities, zooplankton densities, spatial coordinates, and time via

$$u = \frac{cA}{\widehat{r}N}, \quad v = \frac{cZ}{\widehat{r}eN}, \quad x_i = \frac{kX_i}{L}, \quad t = R_0\tau$$

(cf. [33] and [34]), where R_0 is the characteristic (or typical) growth rate of phytoplankton, k is a factor related to the scale of expected patchy patterns, and L is the maximum diameter of the lake in the coordinate direction x_2 , or x_3 . Note that in the case of a square domain, L is the side length. We also rescale the parameters via

$$\begin{aligned} r &= \frac{R}{R_0}, & a &= \frac{C_1K}{C_2R_0}, & b &= \frac{K}{C_2}, & m &= \frac{M}{R_0}, \\ f &= \frac{F}{C_3R_0}, & g &= \frac{K}{C_3A}, & d_1 &= \frac{k^2D_A}{L^2R_0}, & d_2 &= \frac{k^2D_Z}{L^2R_0}, \end{aligned}$$




FIG. 1.2. Numerical solution v of (1.2) at time $t = 1000$ with $d_1 = d_2 = 0.05$, $f = 0$, $g = 10$, solved on a space-time grid of $101 \times 101 \times 12000$. A semi-implicit (in time) Galerkin finite element method with piecewise linear continuous basis functions was employed with homogeneous Neumann boundary conditions. In both cases initial data of the form $u_0 = A(x/100) + B$, $v_0 = C(100 - y)/100 + D$ was employed with the following parameter values: (a) $A = 0.02$, $B = 0.19$, $C = 0.02$, $D = 0.31$, $r = 1$, $a = b = 5$, $m = 0.5$; (b) $A = 0.2$, $B = 0.1$, $C = 0.2$, $D = 0.1$, $r = 1$, $a = b = 20$, $m = 0.8$.

which leads to the dimensionless system

$$(1.2) \quad \begin{cases} \frac{\partial u}{\partial t} = d_1 \Delta u + ru(1 - u) - \frac{auv}{1 + bu}, \\ \frac{\partial v}{\partial t} = d_2 \Delta v + \frac{auv}{1 + bu} - mv - f \frac{gv^2}{1 + g^2 v^2}. \end{cases}$$

Any 2-component reaction-diffusion system with reaction kinetics close to a super critical Hopf bifurcation, with equal diffusion coefficients, can be transformed into a generic reaction-diffusion system of “ λ - ω ” form [40]. Thus as spiral wave solutions of λ - ω reaction-diffusion systems have been proved to exist [9], for appropriate parameter values and initial data we also expect spiral wave solutions to exist for system (1.1). In Figure 1.2 we present snapshots at $t = 1000$ for the uncontrolled system (1.2) representing spiral wave solutions that persist indefinitely (Figure 1.2(a)) or, after initialization, rapidly break up into irregular patterns (Figure 1.2(b)). In both cases we checked that this behavior persists up to $t = 10,000$. This behavior is important to our study as we apply controls that drive the system from the unstable regime to form regular patterns.

In this paper we consider the above model in the context of eutrophication of lakes. Eutrophication is the process where excessive input of nutrients in lakes leads to high levels of phytoplankton (algae) and hence degraded water quality (see [6, 18] and the references therein). The most common approach to improving water quality in this situation is to either reduce the external nutrient loading or enhance zooplankton by reducing planktivorous fish, thereby reducing algal biomass. We focus on the latter approach (“top-down” control), where the nutrient level in the system is determined by a parameter. In practice, planktivorous fish can be reduced by fish removal, or by piscivore¹ stocking. This manipulation of the food-web is called biomanipulation and

¹Species that feed on fish.

is an important approach for improving water quality in eutrophic lakes (e.g., [23, 25, 28, 38, 44, 50]). Evidence from the limnological literature supports the hypothesis that the most effective biomanipulation strategy for improving water quality is the partial removal of fish, and that there may be an optimum harvesting rate of planktivorous fish [11, 21, 48, 49]. This observation helped motivate the work in this paper, where we consider the optimal control of phytoplankton u and zooplankton v densities, where f is the distributed control. We aim to minimize the quadratic cost functional

$$(1.3) \quad J(u, v, f) = \frac{1}{2} \int_Q (|u - \bar{u}|^2 + |v - \bar{v}|^2) dxdt + \frac{\alpha}{2} \int_Q \left| \frac{\partial f}{\partial t} \right|^2 dxdt,$$

where (\bar{u}, \bar{v}) are the desired phytoplankton-zooplankton densities, and $Q = \Omega \times (0, T)$ is the space-time domain of interest. The first two terms in (1.3) measure, with respect to the L^2 -norm in space and time, the difference between the given target densities (\bar{u}, \bar{v}) and the state densities (u, v) . The last term in (1.3) reflects the fact that we want to avoid changing the control (adding or removing fish) too often. The constant α can be chosen to adjust the relative importance of this cost. We assume the state equations (1.2) are augmented with appropriate initial and boundary conditions. Note that as there is no forcing in the first equation of (1.2), the control of phytoplankton must result indirectly through the coupling with the second equation. Furthermore, the forcing in (1.2) enters not merely as an additive inhomogeneous source term, but rather in a multiplicative manner. Additionally, as the control f is a *rate*, it is strictly nonnegative.

We emphasize that although this work has implications for the control of eutrophication in lakes, the main focus of this work is on the rigorous mathematical analysis of the optimal control problem. We do not provide practical implementational details for the improvement of water quality in lakes, but we do provide a theoretical basis on which such a task would be based. We also remark that the mathematical problem and results from numerical simulations may provide insights into the field situation.

It is important to distinguish between the mathematical (optimal control) problem and the practical problem in the field. Mathematically, we assume that f can be manipulated at every point in space and time. However, from a practical point of view, we only have direct control of the net density of fish in the lake at any instant. For example, fish released into a lake may distribute themselves uniformly throughout the lake, or move in schools. Nevertheless, there is significant overlap between the mathematical problem and the field situation, for example, in the case where we wish to reduce the (net) algal growth. In addition to improving water quality we are also interested in the more fundamental aim of maintaining a stable equilibrium between the plankton, thus avoiding the extinction of one or more species.

The aim of this paper is to undertake the mathematical analysis of the optimal control problem introduced above, namely, to minimize (1.3) subject to (1.2). We also provide some numerical results that illustrate the theoretical results. We remark that there are few optimal control studies in the literature for interacting species involving space and time (e.g., [1, 2, 3, 8, 13, 24, 26]) or for reaction-diffusion equations applied in other contexts (e.g., [5, 7, 12, 19, 43]).

2. Mathematical preliminaries.

2.1. Local analysis. We present some details of the local dynamics of the reaction-diffusion system for the state equations (1.2). This is important for deriving necessary conditions on the system parameters for the kinetics to possess biologically




FIG. 2.1. Nullclines for the local kinetics of (1.2) with $r = 1$, $a = b = 20$, $m = 0.8$, $g = 10$, for fish predation rates $f_0 = 0$ (limit cycle, no fish), $f_1 = 0.1$ (limit cycle), $f_2 = 0.25$ (limit cycle), and $f_3 = 0.4$ (phytoplankton dominance).

meaningful equilibria. Furthermore, this acts as a guide in the appropriate choice of parameters for the numerical simulation of optimal solutions.

The local dynamics can be analyzed by considering the nullclines (“zero-isoclines”) of this system, which are the solution curves for

$$(2.1) \quad \begin{aligned} v &= \psi(u) := \frac{r}{a}(1-u)(1+bu), \\ u &= \phi(v) := \frac{g^2mv^2 + fgv + m}{(a-bm)g^2v^2 - bfgv + a - bm}, \end{aligned}$$

corresponding to the first and second equations of (1.2). It is easy to show that there are saddle points at $(0,0)$ and $(0,1)$, and a stationary point (u_s, v_s) (stable or unstable) corresponding to the coexistence of phytoplankton and zooplankton. Note that for positive u_s and v_s we must have

$$(2.2) \quad m < \frac{a}{1+b},$$

which follows from the restriction $0 < u_s < 1$. The nullclines are illustrated in Figure 2.1 for a specific parameter set and increasing predation rate. With no fish present, or a low fish predation rate, there is a limit cycle in the reaction kinetics surrounding the unstable stationary point, while at higher predation rates the system is dominated by a phytoplankton-only state [33, 39]. In general there are no closed

form expressions for u_s and v_s and consequently the analysis must be done numerically in each case. However, in the special case with no fish present, the analysis is straightforward and is briefly outlined below.

With $f = 0$ the intersection of the nullclines $v = \psi(u)$ and $u = \phi(v)$ in (2.1) intersect at (u_s, v_s) , where

$$(2.3) \quad u_s = \frac{m}{a - bm}, \quad v_s = \frac{r}{a}(1 - u_s)(1 + bu_s).$$

To find conditions that guarantee limit cycle kinetics in the positive quadrant of phase space we first prove the existence of a positively invariant region in the sense of [47, Definition 1.1.4] that contains (u_s, v_s) , and then apply standard theory of dynamical systems. We claim that the trapezoidal region $\Sigma \in [0, \infty)^2$ defined by

$$(2.4) \quad \Sigma := \left\{ (u, v) : u \leq l, \quad v - \frac{l}{m}(r + m) + u \leq 0, \quad u \geq 0, \quad v \geq 0 \right\}, \quad l > 1,$$

is positively invariant. To see this, first observe that the reaction kinetics do not point out of Σ along $u = 0$, $v = 0$, and $u = l$. To show that this is also true along the line $v = \frac{l}{m}(r + m) - u$, set $G(u, v) := v - \frac{l}{m}(r + m) + u$ and denote the outward normal to Σ along this line by $\nabla G := (\partial G / \partial u, \partial G / \partial v)^T = (1, 1)^T$. Then denoting the vector of reaction kinetics by $\hat{\mathbf{f}} = (\hat{f}, \hat{g})^T$, observe that

$$\nabla G \cdot \hat{\mathbf{f}}|_{v = \frac{l}{m}(r + m) - u} \leq (r + m)(u - l) \leq 0,$$

which proves the assertion. We derive a simple condition that ensures the critical point given by (2.3) is either an unstable node or an unstable focus, and thus by the Poincaré–Bendixson theorem [47, Theorem 1.1.19] there exists a limit cycle solution surrounding this point. Define

$$A := \begin{pmatrix} \hat{f}_u & \hat{f}_v \\ \hat{g}_u & \hat{g}_v \end{pmatrix} \Big|_{(u_s, v_s)};$$

direct calculation then leads to

$$\text{tr } A := mr \left[\frac{(b + 1)}{a} - \frac{2}{(a - bm)} \right], \quad |A| = \frac{mr}{a} [a - (b + 1)m].$$

Now from (2.2) we have $|A| > 0$, and with the condition

$$(2.5) \quad m < \frac{a(b - 1)}{b(b + 1)},$$

it also follows that $\text{tr } A > 0$. Thus if (2.5) is satisfied, then the critical point (u_s, v_s) is an unstable node or focus (e.g., [46, p. 107]), as required. Thus, to summarize, in the special case with $f = 0$, if the system parameters satisfy (2.5), then there exists a limit cycle solution in the positive quadrant of phase space surrounding the unstable stationary state given by (2.3) (note that condition (2.5) implies condition (2.2)). It is easy to check that (2.5) is satisfied for the parameter sets in Figure 1.2, and also for the numerical results of the uncontrolled problem in section 4.

2.2. Well-posedness of the state equations. We use results from semigroup theory and the abstract theoretical setup of Morgan [35], which is based on the kinetics satisfying a Lyapunov-type condition, to infer the global existence and uniqueness of classical solutions of the state equations (1.2).

Before proving well-posedness of the equations we need to establish the formal setting and restate the fish-plankton system with appropriate initial and boundary data. Let Ω be a bounded and open subset of \mathbb{R}^d , $d \leq 3$, with a boundary $\partial\Omega$ of class C^{2+s} , $s > 0$, i.e., $\partial\Omega$ is a $(d-1)$ -dimensional $C^{2+\nu}$ manifold on which Ω lies locally on one side. The model problem is formulated as follows:

Find the phytoplankton $u(\mathbf{x}, t)$ and zooplankton $v(\mathbf{x}, t)$ densities such that

$$(2.6a) \quad \frac{\partial u}{\partial t} = d_1 \Delta u + ru(1-u) - \frac{auv}{1+bu} \quad \text{in } Q := \Omega \times (0, T),$$

$$(2.6b) \quad \frac{\partial v}{\partial t} = d_2 \Delta v + \frac{auv}{1+bu} - mv - f \frac{gv^2}{1+g^2v^2} \quad \text{in } Q,$$

$$(2.6c) \quad u(\mathbf{x}, 0) = u_0(\mathbf{x}), \quad v(\mathbf{x}, 0) = v_0(\mathbf{x}), \quad \mathbf{x} \in \Omega$$

$$(2.6d) \quad \frac{\partial u}{\partial \boldsymbol{\nu}} = \frac{\partial v}{\partial \boldsymbol{\nu}} = 0 \quad \text{on } \partial\Omega \times (0, T),$$

where the parameters a , b , m , and g are real and strictly positive, and $\boldsymbol{\nu}$ denotes the outward normal to $\partial\Omega$. We assume that the control $f \equiv f(\mathbf{x}, t)$ is a Lipschitz continuous function on Q , which we denote by $f \in \text{Lip}(Q)$, and that the initial data is bounded, i.e., $u_0(\mathbf{x}), v_0(\mathbf{x}) \in L^\infty(\Omega)$. It will be convenient to denote the vector of reaction kinetics by $\hat{\mathbf{f}}(\mathbf{u}) := (\hat{f}(\mathbf{u}), \hat{g}(\mathbf{u}))^T$, where $\mathbf{u} := (u, v)^T$.

THEOREM 2.1. *Let $f \in \text{Lip}(Q)$ and $u_0(\mathbf{x}), v_0(\mathbf{x}) \in L^\infty(\Omega)$. Then there exists a unique nonnegative classical solution of the fish-plankton system (2.6a)–(2.6d) for all $(\mathbf{x}, t) \in \Omega \times [0, \infty)$. Furthermore, if $d_1 = d_2$ and the initial data is chosen in the positively invariant region $\Sigma \in [0, \infty)^2$ given by (2.4), then $(u, v) \in \Sigma$ for all $(\mathbf{x}, t) \in \Omega \times [0, \infty)$.*

Proof. Local existence of solutions is based on well-known semigroup theory (see, for example, Pazy [37] or Henry [20]). From Proposition 1 in [22] it follows immediately that (2.6a)–(2.6d) has a unique noncontinuable classical solution (u, v) for $(\mathbf{x}, t) \in \Omega \times [0, T_{max})$. Moreover, if $T_{max} < \infty$, then

$$(2.7) \quad \lim_{t \uparrow T_{max}} \sup_{\mathbf{x} \in \Omega} \{|u(\mathbf{x}, t)| + |v(\mathbf{x}, t)|\} = \infty.$$

To prove the nonnegativity of solutions observe that the reaction kinetics satisfy

$$\hat{f}(0, v), \hat{g}(u, 0) \geq 0 \quad \forall u, v \geq 0,$$

and the initial data $(u_0(\mathbf{x}), v_0(\mathbf{x}))$ is in $[0, \infty)^2$ for all (or almost every) $\mathbf{x} \in \Omega$. Thus by a maximum principle (see, e.g., [42, Lemma 14.20]) the solution $(u(\mathbf{x}, t), v(\mathbf{x}, t))$ lies in $[0, \infty)^2$ for all $\mathbf{x} \in \Omega$ and for all $t > 0$ for which the solution of (2.6a)–(2.6d) exists. In other words $[0, \infty)^2$ is positively invariant for the system. Proving global existence of solutions from local existence in the equal diffusion case is straightforward. The invariant region yields an L^∞ a priori bound [42] which contradicts nonglobal existence as solutions either exist for all time or blow up in the sup-norm in finite time (see (2.7)) [4]. The proof of global existence results in the distinct diffusion coefficient case requires additional theory as the only admissible invariant regions are products of intervals [42]. We apply the theoretical framework of Morgan [35] to prove global

existence and uniqueness, which involves verifying “intermediate sum” conditions and polynomial growth conditions on the kinetics.

We first define a so-called Lyapunov-type function given by

$$H(\mathbf{u}) := h_1(u) + h_2(v), \quad \text{where } h_1(u) = u, \quad h_2(v) = v.$$

Then with $a_{11} = a_{22} = a_{21} = \mu = q = 1$, $K_2 = K_4 = K_6 = 0$, $K_1 = K_5 = r$, $K_3 = \max\{r, a/b\}$ the following conditions are easily verified for all $\mathbf{u} \in [0, \infty)^2$, corresponding to conditions (H1), (H3), (H4)(i), (H5), and (H6) in [35], respectively:

$$\begin{aligned} a_{11}h_1'(u)\widehat{f}(\mathbf{u}) &\leq K_1(H(\mathbf{u}))^\mu + K_2, \\ a_{21}h_1'(u)\widehat{f}(\mathbf{u}) + a_{22}h_2'(v)\widehat{g}(\mathbf{u}) &\leq K_1(H(\mathbf{u}))^\mu + K_2, \\ h_1'(u)\widehat{f}(\mathbf{u}), h_2'(v)\widehat{g}(\mathbf{u}) &\leq K_3(H(\mathbf{u}))^q + K_4, \\ \nabla H(\mathbf{u}) \cdot \widehat{\mathbf{f}}(\mathbf{u}) &\leq K_5H(\mathbf{u}) + K_6. \end{aligned}$$

Thus Theorems 3.2 and 2.2 in [35] hold, which implies $T_{max} = \infty$, i.e., we have global existence of nonnegative, classical solutions. \square

3. The optimal control problem. Let Ω be a bounded, open subset of \mathbb{R}^2 with smooth boundary $\partial\Omega$ and let the set of all possible target densities $L^2_{loc}(0, T; L^2(\Omega))$ be denoted by \mathcal{T}_{ad} . There are no particular requirements on the target densities (\bar{u}, \bar{v}) other than the fact that the cost functional (1.3) must be bounded. The target densities need not to be solutions of (2.6a)–(2.6d).

Let \mathcal{U}_{ad} be the set of admissible controls

$$\mathcal{U}_{ad} = \{f \in \text{Lip}(Q); 0 \leq f(x, t) \forall (x, t) \in Q\}.$$

Given $T > 0$, $u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega)$, and $(\bar{u}, \bar{v}) \in \mathcal{T}_{ad}$, then (u, v, f) is said to be an admissible element if $u, v \in L^2(0, T; H^1(\Omega))$, $f \in \mathcal{U}_{ad}$, the functional $J(u, v, f)$ is bounded, and (u, v, f) satisfies (2.6a)–(2.6d). Let \mathcal{A}_{ad} be the set of admissible states and controls. With this notation, the formulation of the optimal control problem is given by the following:

$$\begin{aligned} \text{(P)} \quad &\text{Given } T > 0, u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega), \text{ and } (\bar{u}, \bar{v}) \in \mathcal{T}_{ad}, \\ &\text{find } (u^*, v^*, f^*) \in \mathcal{A}_{ad} \text{ such that } J(u^*, v^*, f^*) \leq J(u, v, f) \\ &\forall (u, v, f) \in \mathcal{A}_{ad}. \end{aligned}$$

THEOREM 3.1. *Given $u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega)$ and $(\bar{u}, \bar{v}) \in \mathcal{T}_{ad}$, there exists a solution (u^*, v^*, f^*) of the optimal control problem (P).*

Proof. The admissible set is bounded and nonempty, e.g., the system (2.6a)–(2.6d) has a solution for $f = 0$ (see [14]). Let $\{(u_n, v_n, f_n)\}_n$ be a minimizing sequence in \mathcal{A}_{ad} . The sequence $\{\frac{\partial f_n}{\partial t}\}$ is bounded in $L^2(0, T; L^2(\Omega))$, and therefore with a subsequence, again indexed by n , we have

$$f_n \rightarrow f^* \quad \text{weakly in } H^1([0, T]; L^2(\Omega)).$$

On the other hand, from (2.6a)–(2.6b) we have

$$\|u_n(t)\|_{L^2(\Omega)}^2 + \|v_n(t)\|_{L^2(\Omega)}^2 + \int_0^t \left(d_1 \|\nabla u_n(t)\|_{L^2(\Omega)}^2 + d_2 \|\nabla v_n(t)\|_{L^2(\Omega)}^2 \right) dt \leq C$$

for all $t \in [0, T]$. Here and in what follows we denote by C a positive constant independent of u, v, f , and n . Next we multiply system (2.6a)–(2.6b) by $(-\Delta u_n, -\Delta v_n)$ and integrate over $(0, t)$ to get, after some calculation involving Gronwall’s lemma,

$$\begin{aligned} & \|\nabla u_n(t)\|_{L^2(\Omega)}^2 + \|\nabla v_n(t)\|_{L^2(\Omega)}^2 + \int_0^t \left(d_1 \|\Delta u_n(s)\|_{L^2(\Omega)}^2 + d_2 \|\Delta v_n(s)\|_{L^2(\Omega)}^2 \right) dt \leq C, \\ & \left\| \frac{d}{dt} u_n(t) \right\|_{L^2(\Omega)}^2 + \left\| \frac{d}{dt} v_n(t) \right\|_{L^2(\Omega)}^2 \leq C \quad \forall t \in [0, T]. \end{aligned}$$

Using the classical Lions–Aubin compactness lemma [27] and Ascoli’s theorem [10], and noting that the injection of $H^1(\Omega)$ into $L^2(\Omega)$ is compact, we infer that $\{u_n\}, \{v_n\}$ are compact in $L^2(0, T; H^1(\Omega)) \cap C([0, T]; L^2(\Omega))$. Thus selecting further subsequences, if necessary, we have

$$\begin{aligned} u_n &\rightarrow u^*, \quad v_n \rightarrow v^* && \text{strongly in } L^2(0, T; H^1(\Omega)) \cap C([0, T]; L^2(\Omega)), \\ u_n &\rightarrow u^*, \quad v_n \rightarrow v^* && \text{weakly in } L^2(0, T; H^2(\Omega)), \\ \frac{d}{dt} u_n &\rightarrow \frac{d}{dt} u^*, \quad \frac{d}{dt} v_n \rightarrow \frac{d}{dt} v^* && \text{weakly in } L^2(0, T; L^2(\Omega)). \end{aligned}$$

Moreover we have that

$$\begin{aligned} u_n(1 - u_n) &\rightarrow u^*(1 - u^*) && \text{strongly in } L^2(0, T; L^2(\Omega)), \\ \frac{u_n v_n}{1 + bu_n} &\rightarrow \frac{u^* v^*}{1 + bu^*} && \text{strongly in } L^2(0, T; L^2(\Omega)), \\ f_n \frac{v_n^2}{1 + g^2 v_n^2} &\rightarrow f^* \frac{v^{*2}}{1 + g^2 v^{*2}} && \text{weakly in } L^2(0, T; L^2(\Omega)). \end{aligned}$$

Letting n go to ∞ we see that (u^*, v^*, f^*) satisfies the system (2.6a)–(2.6d) and $J(u^*, v^*, f^*) = \inf_{\mathcal{A}_{ad}} J$. \square

3.1. First-order necessary conditions. We show that the optimal solution must satisfy the first-order necessary condition associated with the optimal control problem (P).

We introduce the tangential (contingent) cone to \mathcal{A}_{ad} at $(u, v, f) \in \mathcal{A}_{ad}$:

$$\begin{aligned} \text{Tan } \mathcal{A}_{ad}(u, v, f) &= \left\{ (y, z, h) \mid y, z \in L^2(Q), h \in \text{Tan } \mathcal{U}_{ad}(f) \quad \text{and} \right. \\ & \frac{\partial y}{\partial t} = d_1 \Delta y + ry(1 - 2u) - \frac{av}{(1 + bu)^2} y - \frac{au}{1 + bu} z \text{ in } Q, \\ & \frac{\partial z}{\partial t} = d_2 \Delta z + \frac{av}{(1 + bu)^2} y + \left(\frac{au}{1 + bu} - m - \frac{2gvf}{(1 + g^2 v^2)^2} \right) z - \frac{gv^2 h}{1 + g^2 v^2} \text{ in } Q, \\ (3.1) \quad & \frac{\partial y}{\partial \nu} = \frac{\partial z}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T), \\ & \left. y(x, 0) = z(x, 0) = 0 \quad \text{in } \Omega \right\}. \end{aligned}$$

Recall that if

$$J(u^*, v^*, f^*) = \inf_{(u, v, f) \in \mathcal{A}_{ad}} J(u, v, f)$$

and the functional $J(u, v, f)$ is Gâteaux differentiable, then necessarily

$$(3.2) \quad \partial J(u^*, v^*, f^*)(y, z, h) \geq 0 \quad \forall (y, z, h) \in \text{Tan } \mathcal{A}_{ad}(u^*, v^*, f^*),$$

where $\partial J(u^*, v^*, f^*)$ denotes the Gâteaux derivative of J at $(u^*, v^*, f^*) \in \mathcal{A}_{ad}$. Applying the optimum principle given by (3.2), it follows that

$$(3.3) \quad \int_Q \left[(u^* - \bar{u})y + (v^* - \bar{v})z + \alpha \frac{\partial f^*}{\partial t} \frac{\partial h}{\partial t} \right] dxdt \geq 0 \quad \forall (y, z, h) \in \text{Tan } \mathcal{A}_{ad}(u^*, v^*, f^*).$$

THEOREM 3.2. *Let $u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega)$. The mapping $(u, v) = (u(f), v(f))$ from \mathcal{U}_{ad} to $L^2(0, T; H^1(\Omega))$, defined as the solution of the system (2.6a)–(2.6d), has a Gâteaux derivative $(D(u, v)/Df) \cdot h$ in every direction $h \in \text{Tan } \mathcal{U}_{ad}(f)$. Furthermore, $(y(h), z(h)) = (D(u, v)/Df) \cdot h$ is the classical solution of problem (3.1).*

Proof. Let $h \in \text{Lip}(Q)$ be such that $f + \lambda h \in \mathcal{U}_{ad}$ for all $\lambda \in \mathbb{R}$ sufficiently small and let (u_f, v_f) and $(u_{f+\lambda h}, v_{f+\lambda h})$ denote the solutions of (2.6a)–(2.6d) with fish predation rates f and $f + \lambda h$, respectively. To simplify the notation in this proof we use $\|\cdot\|$ to denote $\|\cdot\|_{L^2(\Omega)}$.

We need to prove that

$$\lim_{\lambda \rightarrow 0} \frac{1}{\lambda} \left\{ \|u_{f+\lambda h} - u_f - \lambda y(h)\|_{L^2(0, T; H^1(\Omega))} + \|v_{f+\lambda h} - v_f - \lambda z(h)\|_{L^2(0, T; H^1(\Omega))} \right\} = 0.$$

We set $\tilde{u} = u_{f+\lambda h} - u_f - \lambda y(h)$, $\tilde{v} = v_{f+\lambda h} - v_f - \lambda z(h)$ so that (\tilde{u}, \tilde{v}) satisfies the system

$$(3.4a) \quad \frac{\partial \tilde{u}}{\partial t} = d_1 \Delta \tilde{u} + r\tilde{u} - r\Lambda_1 - a\Lambda_2 \quad \text{in } Q,$$

$$(3.4b) \quad \frac{\partial \tilde{v}}{\partial t} = d_2 \Delta \tilde{v} + a\Lambda_2 - m\tilde{v} - g\Lambda_3 \quad \text{in } Q,$$

$$(3.4c) \quad \frac{\partial \tilde{u}}{\partial \nu} = \frac{\partial \tilde{v}}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T),$$

$$(3.4d) \quad \tilde{u}(x, 0) = \tilde{v}(x, 0) = 0 \quad \text{in } \Omega,$$

where

$$\Lambda_1 = u_{f+\lambda h}^2 - u_f^2 - 2\lambda y u_f, \quad \Lambda_2 = \frac{u_{f+\lambda h} v_{f+\lambda h}}{1 + b u_{f+\lambda h}} - \frac{u_f v_f}{1 + b u_f} - \lambda \frac{v_f y}{(1 + b u_f)^2} - \lambda \frac{u_f z}{1 + b u_f},$$

$$\Lambda_3 = (f + \lambda h) \frac{v_{f+\lambda h}^2}{1 + g^2 v_{f+\lambda h}^2} - f \frac{v_f^2}{1 + g^2 v_f^2} - \lambda \frac{2v_f f}{(1 + g^2 v_f^2)^2} z - \lambda \frac{v_f^2 h}{1 + g^2 v_f^2}.$$

Now we multiply (3.4a)–(3.4b) by \tilde{u} and \tilde{v} , respectively, and get

$$\frac{d}{dt} (\|\tilde{u}\|^2 + \|\tilde{v}\|^2) + (\|\nabla \tilde{u}\|^2 + \|\nabla \tilde{v}\|^2) \leq C (\|\tilde{u}\|^2 + \|\tilde{v}\|^2 + \|\Lambda_1\|^2 + \|\Lambda_2\|^2 + \|\Lambda_3\|^2).$$

We denote $\hat{u} = u_{f+\lambda h} - u_f$, $\hat{v} = v_{f+\lambda h} - v_f$ and use the fact that (u_f, v_f) is a classical solution to (2.6a)–(2.6d), yielding

$$\begin{aligned} \int_\Omega \Lambda_1^2 dx &= \int_\Omega (\hat{u}^2 + 2u_f \tilde{u})^2 dx \leq 2\|\hat{u}\|_{L^4(\Omega)}^4 + C\|\tilde{u}\|^2, \\ \int_\Omega \Lambda_2^2 dx &\leq C \int_\Omega (\hat{u}^4 + \hat{v}^4 + \tilde{u}^2 + \tilde{v}^2) dx, \\ \int_\Omega \Lambda_3^2 dx &\leq C \int_\Omega (\hat{v}^4 + \hat{v}^6 + \tilde{v}^2 + \lambda^2 h^2 \tilde{v}^2) dx \leq C \int_\Omega (\hat{v}^4 + \tilde{v}^2 + \lambda^4) dx. \end{aligned}$$

Hence from the above estimates and the Gronwall inequality, this leads to

$$\begin{aligned} & \|\tilde{u}(t)\|^2 + \|\tilde{v}(t)\|^2 + \int_0^T (\|\nabla\tilde{u}(s)\|^2 + \|\nabla\tilde{v}(s)\|^2) ds \\ & \leq C \left(\lambda^4 + \int_0^T (\|\hat{u}\|^2 \|\hat{u}\|_{H^1(\Omega)}^2 + \|\hat{v}\|^2 \|\hat{v}\|_{H^1(\Omega)}^2) dt \right) \quad \forall t \in [0, T]. \end{aligned}$$

To estimate the norm of \hat{u}, \hat{v} in $L^2(0, T; H^1(\Omega))$ we note that

$$\begin{aligned} \frac{\partial \hat{u}}{\partial t} &= d_1 \Delta \hat{u} + r \hat{u} - r \hat{u}(\hat{u} + 2u_f) - a \left(\frac{\hat{v}(\hat{u} + u_f)}{1 + b(\hat{u} + u_f)} - \frac{\hat{u}v_f}{(1 + bu_f)[1 + b(\hat{u} + u_f)]} \right), \\ \frac{\partial \hat{v}}{\partial t} &= d_2 \Delta \hat{v} + a \left(\frac{\hat{v}(\hat{u} + u_f)}{1 + b(\hat{u} + u_f)} - \frac{\hat{u}v_f}{(1 + bu_f)[1 + b(\hat{u} + u_f)]} \right) - m \hat{v} \\ & \quad + g \left(f \frac{\hat{v}(\hat{v} + 2v_f)}{(1 + g^2v_f^2)(1 + g^2(\hat{v} + v_f)^2)} + \lambda h \frac{(\hat{v} + v_f)^2}{1 + g^2(\hat{v} + v_f)^2} \right), \\ \frac{\partial \hat{u}}{\partial \nu} &= \frac{\partial \hat{v}}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T), \\ \hat{u}(x, 0) &= \hat{v}(x, 0) = 0 \quad \text{in } \Omega. \end{aligned}$$

We then easily obtain

$$\|\hat{u}(t)\|^2 + \|\hat{v}(t)\|^2 + \int_0^t (\|\nabla\hat{u}(s)\|^2 + \|\nabla\hat{v}(s)\|^2) ds \leq C\lambda^2 \int_0^t \|h(s)\|^2 ds \quad \forall t \in [0, T].$$

From the last two results we obtain the estimate

$$\int_0^T (\|\tilde{u}\|_{H^1(\Omega)}^2 + \|\tilde{v}\|_{H^1(\Omega)}^2) dt \leq C\lambda^4,$$

from which our claim follows. The proof of the regularity of y and z is similar to the proof of Theorem 2.1. \square

The Gâteaux derivative gives useful information about the sensitivity of the system (2.6a)–(2.6d) at a particular point (u, v) in a particular direction h , but complete information requires the solution of (3.1) for every direction h . Fortunately, in order to minimize the functional we need only an integral over all these directions, which can be obtained by solving a single adjoint equation.

THEOREM 3.3. *Let $u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega)$ and $(\bar{u}, \bar{v}) \in \mathcal{T}_{ad}$ be given. If (u^*, v^*, f^*) is an optimal solution for (P), then we have*

$$(3.5) \quad \int_Q \left(\frac{gv^{*2}k}{1 + g^2v^{*2}}(f - f^*) + \alpha \frac{\partial f^*}{\partial t} \left(\frac{\partial f}{\partial t} - \frac{\partial f^*}{\partial t} \right) \right) dxdt \geq 0 \quad \forall f \in \mathcal{U}_{ad},$$

where (p, k) is the unique classical solution of the adjoint equation

$$\begin{aligned} (3.6a) \quad & \frac{\partial p}{\partial t} + d_1 \Delta p + rp(1 - 2u^*) - a \frac{v^*}{(1 + bu^*)^2} p + a \frac{v^*}{(1 + bu^*)^2} k = u^* - \bar{u} \text{ in } Q, \\ & \frac{\partial k}{\partial t} + d_2 \Delta k + a \frac{u^*}{1 + bu^*} k - mk - 2 \frac{gf^*v^*}{(1 + g^2v^{*2})^2} k - a \frac{u^*}{1 + bu^*} p = v^* - \bar{v} \text{ in } Q, \\ & \frac{\partial p}{\partial \nu} = \frac{\partial k}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T), \end{aligned}$$

(3.6b)

$$p(x, T) = k(x, T) = 0 \quad \text{in } \Omega.$$

Proof. Let (u^*, v^*, f^*) be an optimal solution. From the optimality condition (3.3) and (3.1) and from (3.6a)–(3.6b), we have after integration by parts

$$\begin{aligned} 0 &\leq \int_Q \left[(u^* - \bar{u})y + (v^* - \bar{v})z + \alpha \frac{\partial f^*}{\partial t} \frac{\partial h}{\partial t} \right] dxdt \\ &= \int_Q \left\{ y \left[\frac{\partial p}{\partial t} + d_1 \Delta p + rp(1 - 2u^*) - a \frac{v^*}{(1 + bu^*)^2} p + a \frac{v^*}{(1 + bu^*)^2} k \right] \right. \\ &\quad \left. + z \left(\frac{\partial k}{\partial t} + d_2 \Delta k + a \frac{u^*}{1 + bu^*} k - mk - \frac{2gf^*v^*}{(1 + g^2v^{*2})^2} k - a \frac{pu^*}{1 + bu^*} \right) + \alpha \frac{\partial f^*}{\partial t} \frac{\partial h}{\partial t} \right\} dxdt \\ &= \int_Q p \left[-\frac{\partial y}{\partial t} + d_1 \Delta y + ry(1 - 2u^*) - a \frac{v^*}{(1 + bu^*)^2} y - a \frac{u^*}{1 + bu^*} z \right] dxdt + \int_\Omega yp \Big|_0^T dx \\ &\quad + \int_Q k \left[-\frac{\partial z}{\partial t} + d_2 \Delta z - \frac{2gf^*v^*}{(1 + g^2v^{*2})^2} z + a \frac{u^*}{1 + bu^*} z - mz + a \frac{v^*}{(1 + bu^*)^2} y \right] dxdt \\ &\quad + \int_\Omega zk \Big|_0^T dx + \alpha \int_\Omega \frac{\partial f^*}{\partial t} \frac{\partial h}{\partial t} dxdt \\ &= \int_Q k \frac{gv^{*2}}{1 + g^2v^{*2}} h dxdt + \alpha \int_\Omega \frac{\partial f^*}{\partial t} \frac{\partial h}{\partial t} dxdt, \end{aligned}$$

and therefore (3.5). \square

4. Numerical results. To illustrate the theoretical results of the optimal control problem in the previous section we present results of numerical experiments in two space dimensions. The state equations and adjoint equations were solved using a “lumped mass,” semi-implicit (in time) Galerkin finite element method with piecewise linear continuous basis functions. We showed previously that this approach was highly effective in solving the forward-in-time equations of a similar predator-prey system [15].

The control was updated using a variable-step gradient algorithm based on the fully discrete optimality condition (see (4.4) below). At each iteration of the gradient algorithm the method requires the sequential solution of the discrete state and adjoint equations (see (4.1) and (4.3) below). In practice one cannot solve these systems simultaneously. The discrete state equations are solved by marching forward in time starting from an initial condition, while the discrete adjoint equations are solved by marching backward in time (from T) starting from a terminal condition. For further details, see [16].

We employed a (uniform) right-angled triangulation Ω^h of the square $\Omega = [0, L] \times [0, L]$, with space steps h , and numerically solved the optimal control problem up to time T with uniform time steps Δt . We introduce S^h , the standard finite element space

$$S^h := \{v \in C(\bar{\Omega}) : v|_\tau \text{ is linear } \forall \tau \in \Omega^h\} \subset H^1(\Omega).$$

Let $\{x_i\}_{i=0}^J$ be the set of nodes of the triangulation. We introduce $\pi^h : C(\bar{\Omega}) \mapsto S^h$, the Lagrange interpolation operator, such that $\pi^h v(x_j) = v(x_j)$ for all $j = 0, \dots, J$. In order to formulate our finite element approximation of the reaction-diffusion system we define a discrete L^2 inner product on $C(\bar{\Omega})$ given by $(u, v)^h := \int_\Omega \pi^h(u(x)v(x)) dx$,

which approximates the usual L^2 inner product (u, v) . Given $\mathbf{f}_h \in \text{Lip}(\Omega)$ and $u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega)$, $(\mathbf{u}_h, \mathbf{v}_h)$ is a solution of the fully discrete, semi-implicit (in time) fish-plankton system if $u_h^{(n)}, v_h^{(n)} \in S^h$ satisfies the system

$$(4.1) \quad \begin{aligned} & \frac{1}{\Delta t} \left(u_h^{(n)} - u_h^{(n-1)}, \chi_h \right)^h + d_1 \left(\nabla u_h^{(n)}, \nabla \chi_h \right) \\ &= \left(r u_h^{(n)} (1 - |u_h^{(n-1)}|) - \frac{a u_h^{(n-1)} v_h^{(n)}}{1 + b |u_h^{(n-1)}|}, \chi_h \right)^h, \\ & \frac{1}{\Delta t} \left(v_h^{(n)} - v_h^{(n-1)}, \chi_h \right)^h + d_2 \left(\nabla v_h^{(n)}, \nabla \chi_h \right) \\ &= \left(\frac{a u_h^{(n-1)} v_h^{(n)}}{1 + b |u_h^{(n-1)}|} - m v_h^{(n)} - \frac{g f_h^{(n)} v_h^{(n-1)2}}{1 + g^2 v_h^{(n-1)2}}, \chi_h \right)^h \end{aligned}$$

for all $\chi_h \in S^h$, $n = 1, 2, \dots, N$, with initial densities $u_h^{(0)} = \pi^h u_0(x)$, $v_h^{(0)} = \pi^h v_0(x)$. The discrete cost functional used in the optimal control problem is given by

$$(4.2) \quad \mathcal{J}_h^N(\mathbf{u}_h, \mathbf{v}_h, \mathbf{f}_h) = \frac{\Delta t}{2} \sum_{n=1}^N \left(\|u_h^{(n)} - \bar{u}^{(n)}\|^2 + \|v_h^{(n)} - \bar{v}^{(n)}\|^2 \right) + \frac{\alpha}{2\Delta t} \sum_{n=1}^N \|f_h^{(n)} - f_h^{(n-1)}\|^2.$$

Thus we can now formulate the fully discrete optimal control problem as follows:

$$(P^{h,\Delta t}) \quad \begin{aligned} & \text{Given } \Delta t = T/N, h = L/J, u_0, v_0 \in H^1(\Omega) \cap L^\infty(\Omega) \\ & \text{and } (\bar{u}, \bar{v}) \in \mathcal{T}_{ad}, \text{ find } (\mathbf{u}_h, \mathbf{v}_h, \mathbf{f}_h) \in S^h \times S^h \times S^h \text{ such} \\ & \text{that (4.1) is satisfied for } n = 1, 2, \dots, N \text{ and the cost} \\ & \text{functional (4.2) is minimized.} \end{aligned}$$

To complete the fully discrete optimality system we also need the following fully discrete adjoint system: The adjoint functions $p_h^{(n)}, k_h^{(n)} \in S^h$ satisfy

$$(4.3) \quad \begin{aligned} & -\frac{1}{\Delta t} \left(p_h^{(n)} - p_h^{(n-1)}, \chi_h \right)^h + d_1 \left(\nabla p_h^{(n-1)}, \nabla \chi_h \right) \\ &= \left(r(1 - 2|u_h^{(n)}|) p_h^{(n-1)} - a \frac{v_h^{(n)}}{(1 + b |u_h^{(n)}|)^2} p_h^{(n-1)} + a \frac{v_h^{(n)}}{1 + b |u_h^{(n)}|^2} k_h^{(n-1)} - u_h^{(n)} - \bar{u}^{(n)}, \chi_h \right)^h \\ & -\frac{1}{\Delta t} \left(k_h^{(n)} - k_h^{(n-1)}, \chi_h \right)^h + d_2 \left(\nabla k_h^{(n-1)}, \nabla \chi_h \right) \\ &= \left(a \frac{u_h^{(n)}}{1 + b |u_h^{(n)}|^2} p_h^{(n-1)} + a \frac{u_h^{(n)}}{1 + b |u_h^{(n)}|^2} k_h^{(n-1)} - m k_h^{(n-1)} - g f_h^{(n)} \frac{2v_h^{(n)}}{1 + g^2 v_h^{(n)2}} - v_h^{(n)} - \bar{v}^{(n)}, \chi_h \right)^h \end{aligned}$$

for all $\chi_h \in S^h$, $n = 1, 2, \dots, N$, with the terminal conditions $p_h^{(N)} = k_h^{(N)} = 0$. The fully discrete optimality condition is

$$(4.4) \quad \begin{aligned} 0 \leq & \Delta t \sum_{n=1}^{N-1} \left(\tilde{f}_h^{(n)}, g \frac{v_h^{(n)2}}{1 + g^2 v_h^{(n)2}} k_h^{(n-1)} - \alpha \frac{f_h^{(n+1)} - 2f_h^{(n)} + f_h^{(n-1)}}{\Delta t} \right) \\ & - \frac{\alpha}{\Delta t} \left(\tilde{f}_h^{(0)}, f_h^{(1)} - f_h^{(0)} \right) + \left(\tilde{f}_h^{(N)}, \Delta t g \frac{v_h^{(N)2}}{1 + g^2 v_h^{(N)2}} k_h^{(N-1)} + \alpha \frac{f_h^{(N)} - f_h^{(N-1)}}{\Delta t} \right) \end{aligned}$$




FIG. 4.1. Uncontrolled (a), optimally controlled (b), target (c), and the control f (d) for phytoplankton densities u at time $T = 100$ for (1.2). Parameter values: $d_1 = d_2 = 0.05$, $r = 1$, $a = b = 20$, $m = 0.8$, $g = 10$, $\alpha = 10^{-5}$. For details of initial data, see text.

for all $\tilde{\mathbf{f}}_h \in \text{Tan}\mathcal{U}_{ad}(\mathbf{f}_h)$.

In the numerical experiments we chose the domain $\Omega = [0, 100] \times [0, 100]$ and initial conditions of the optimal control problem to be the freely evolving ($f = 0$) system (1.2) at time $t = 1000$, with parameter values and initial data corresponding to Figure 1.2(b) (see caption). The target functions are one-armed Archimedean spirals with period of rotation equal to 20 and are given by

$$\begin{aligned}\bar{u}(R, \theta, t) &= 0.495 \cos(\theta + R/12 + \pi t/10) + 0.495, \\ \bar{v}(R, \theta, t) &= 0.21 \sin(\theta + R/12 + \pi t/10) + 0.22,\end{aligned}$$

where $R := \sqrt{(x-50)^2 + (y-50)^2}$ and $\theta := \arctan[(y-50)/(x-50)]$. The numerical results for the optimal control of phytoplankton densities u are shown in Figure 4.1 at time $T = 100$. A plot of the reduction in cost functional with increasing iteration count suggests the near optimality of the system (see [16]). The results show that at time $T = 100$ the controlled system is close to the desired state, and that we were successful in driving the system from a disordered state to an ordered one.

5. Conclusions. The mathematical formulation, analysis, and numerical solution of an optimal control problem for a nonlinear plankton-fish reaction-diffusion

system was presented. The model was discussed in the context of biomanipulation of eutrophic lakes. After considering the local dynamics of the system and proving the global existence and uniqueness of the classical solutions of the state equations, we presented the mathematical analysis of the plankton-fish optimal control problem. Numerical solutions were obtained with the aid of a semi-implicit Galerkin finite element method with piecewise linear continuous basis functions. The numerical results illustrate the ability of a variable step-size gradient algorithm to drive the plankton dynamics from a chaotic regime to an (arbitrary) ordered state. The time taken to achieve the ordered distribution of phytoplankton in nondimensional units was $t = 100$, which with an assumed maximum growth rate for phytoplankton in eutrophic conditions of $R_0 = 0.5$ per day [39] gives the time taken to achieve this state to be $\tau = 200$ days. The theoretical results in this paper provide the basis for a numerical analysis of the optimal control problem [16]. Furthermore, our results can be generalized in numerous ways to include, for example, convection driven flows, forcing via nutrient inputs, and stochastic influences. We leave these tasks for future work.

REFERENCES

- [1] S. ANIȚA, *Optimal control of a nonlinear population dynamics with diffusion*, J. Math. Anal. Appl., 152 (1990), pp. 176–208.
- [2] S. ANIȚA, *A fractional step scheme for the optimal control of a nonlinear population dynamics with diffusion*, An. Științ. Univ. Al. I. Cuza Iași. Mat., 46 (2000), pp. 157–168.
- [3] N. APREUTESEI, *An optimal control problem of Lotka-Volterra system with diffusion*, Bul. Inst. Politeh. Iași. Secț. I. Mat. Mec. Teor. Fiz., 44 (1998), pp. 31–41.
- [4] J. BALL, *Remarks on blow-up and nonexistence theorems for nonlinear evolution equations*, Quart. J. Math. Oxford, 2 (1977), pp. 473–486.
- [5] A. BORZI AND R. GRIESSE, *Experiences with a space-time multigrid method for the optimal control of a chemical turbulence model*, Internat. J. Numer. Methods Fluids, 47 (2005), pp. 879–885.
- [6] S. CARPENTER, D. CHRISTENSEN, J. COLE, K. COTTINGHAM, X. HE, J. HODGSON, J. KITCHELL, S. KNIGHT, M. PACE, D. POST, D. SCHINDLER, AND N. VOICHICK, *Biological control of eutrophication in lakes*, Environ. Sci. Technol., 29 (1995), pp. 784–786.
- [7] S. CERRAL, *Optimal control problems for stochastic reaction-diffusion systems with non-Lipschitz coefficients*, SIAM J. Control Optim., 39 (2001), pp. 1779–1816.
- [8] R. Z. CHEN, D. S. ZHANG, AND J. Q. LI, *Existence and uniqueness of solution and boundary control for population systems with spatial diffusion*, J. Systems Sci. Math. Sci., 22 (2002), pp. 1–13.
- [9] D. S. COHEN, J. C. NEU, AND R. R. ROSALES, *Rotating spiral wave solutions of reaction-diffusion equations*, SIAM J. Appl. Math., 35 (1978), pp. 536–547.
- [10] J. DIEUDONNÉ, *Foundations of Modern Analysis*, Pure Appl. Math., Academic Press, New York, 1960.
- [11] R. DRENNER AND K. HAMBRIGHT, *Biomanipulation of fish assemblages as a lake restoration technique*, Arch. Hydrobiol., 146 (1999), pp. 129–165.
- [12] K. FISTER AND C. M. MCCARTHY, *Optimal control of a chemotaxis system*, Quart. Appl. Math., 61 (2003), pp. 193–211.
- [13] R. FISTER, *Optimal control of harvesting in a predator-prey parabolic system*, Houston J. Math., 23 (1997), pp. 341–355.
- [14] M. GARVIE AND C. TRENCH, *Analysis of two generic spatially extended predator-prey models*, Nonlinear Anal. Real World Appl., submitted.
- [15] M. GARVIE AND C. TRENCH, *Finite element approximation of spatially extended predator-prey interactions with the Holling type II functional response*, Numer. Math., to appear.
- [16] M. GARVIE AND C. TRENCH, *Numerical Analysis of a Nutrient-Plankton Optimal Control Problem*, in preparation, 2006.
- [17] W. GENTLEMAN, A. LEISING, B. FROST, S. STROM, AND J. MURRAY, *Functional responses for zooplankton feeding on multiple resources: A review of assumptions and biological dynamics*, Deep-Sea Res. Pt. II, 50 (2003), pp. 2847–2875.

- [18] L.-A. HANSSON, M. GYLLSTRÖM, A. STAHL-DELBANCO, AND M. SVENSSON, *Responses to fish predation and nutrients by plankton at different levels of taxonomic resolution*, *Freshwater Biol.*, 49 (2004), pp. 1538–1550.
- [19] J. HE AND R. GLOWINSKI, *Neumann control of unstable parabolic systems: Numerical approach*, *J. Optim. Theory Appl.*, 96 (1998), pp. 1–55.
- [20] D. HENRY, *Geometric Theory of Semilinear Parabolic Equations*, *Lecture Notes in Math.* 840, Springer-Verlag, New York, 1981.
- [21] J. HIETALA, K. VAKKILAINEN, AND T. KAIRESALO, *Community resistance and change to nutrient enrichment and fish manipulation in a vegetated lake littoral*, *Freshwater Biol.*, 49 (2004), pp. 1525–1537.
- [22] S. L. HOLLIS, R. H. MARTIN, JR., AND M. PIERRE, *Global existence and boundedness in reaction-diffusion systems*, *SIAM J. Math. Anal.*, 18 (1987), pp. 744–761.
- [23] J. HORPPILA, H. PELTONEN, T. MALINEN, E. LUOKKANEN, AND T. KAIRESALO, *Top-down or bottom-up effects by fish: Issues of concern in biomanipulation of lakes*, *Restor. Ecol.*, 6 (1998), pp. 20–28.
- [24] I. HRINCA, *An optimal control problem for the Lotka-Volterra system with diffusion*, *Panamer. Math. J.*, 12 (2002), pp. 23–46.
- [25] A. KOZAK AND R. GOLDYN, *Zooplankton versus phyto- and bacterioplankton in the Maltański Reservoir (Poland) during an extensive biomanipulation experiment*, *J. Plankton Res.*, 26 (2004), pp. 37–48.
- [26] S. LENHART, M. LIANG, AND V. PROTOPODESCU, *Optimal control of boundary habitat hostility for interacting species*, *Math. Method. Appl. Sci.*, 22 (1999), pp. 1061–1077.
- [27] J. LIONS, *Quelques Méthodes de Résolution des Problèmes aux Limites non Linéaires*, Dunod Gauthier-Villars, Paris, 1969.
- [28] M. LUND AND J. DAVIS, *Seasonal dynamics of plankton communities and water chemistry in a eutrophic wetland (Lake Monger, Western Australia): Implications for biomanipulation*, *Mar. Freshwater Res.*, 51 (2000), pp. 321–332.
- [29] H. MALCHOW, *Motional instabilities in prey-predator systems*, *J. Theoret. Biol.*, 204 (2000), pp. 639–647.
- [30] H. MALCHOW, F. HILKER, AND S. PETROVSKII, *Noise and productivity dependence of spatio-temporal pattern formation in a prey-predator system*, *Discrete Contin. Dyn. Syst. Ser. B*, 4 (2004), pp. 705–711.
- [31] H. MALCHOW, S. PETROVSKII, AND A. MEDVINSKY, *Pattern formation in models of plankton dynamics. A synthesis*, *Oceanol. Acta*, 24 (2001), pp. 479–487.
- [32] H. MALCHOW, S. PETROVSKII, AND A. MEDVINSKY, *Numerical study of plankton-fish dynamics in a spatially structured and noisy environment*, *Ecol. Model.*, 149 (2002), pp. 247–255.
- [33] H. MALCHOW, B. RADTKE, M. KALLACHE, A. MEDVINSKY, D. TIKHONOV, AND S. PETROVSKII, *Spatio-temporal pattern formation in coupled models of plankton dynamics and fish school motion*, *Nonlinear Anal.-Real.*, 1 (2000), pp. 53–67.
- [34] A. B. MEDVINSKY, S. V. PETROVSKII, I. A. TIKHONOVA, H. MALCHOW, AND B.-L. LI, *Spatio-temporal complexity of plankton and fish dynamics*, *SIAM Rev.*, 44 (2002), pp. 311–370.
- [35] J. MORGAN, *Global existence for semilinear parabolic systems*, *SIAM J. Math. Anal.*, 20 (1989), pp. 1128–1144.
- [36] M. PASCUAL, *Diffusion-induced chaos in a spatial predator-prey system*, *Proc. Roy. Soc. London Ser. B*, 251 (1993), pp. 1–7.
- [37] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, *Appl. Math. Sci.* 44, Springer-Verlag, New York, 1983.
- [38] D. REJAS, S. DECLERCK, J. AUWERKERKEN, P. TAK, AND L. MEESTER, *Plankton dynamics in a tropical floodplain lake: Fish, nutrients, and the relative importance of bottom-up and top-down control*, *Freshwater Biol.*, 50 (2005), pp. 52–69.
- [39] M. SCHEFFER, *Fish and nutrients interplay determines algal biomass: A minimal model*, *Oikos*, 62 (1991), pp. 271–282.
- [40] J. SHERRATT, *Invading wave fronts and their oscillatory wakes are linked by a modulated travelling phase resetting wave*, *Phys. D*, 117 (1998), pp. 145–166.
- [41] J. SHERRATT, B. EAGAN, AND M. LEWIS, *Oscillations and chaos behind predator-prey invasion: Mathematical artifact or ecological reality?*, *Philos. Trans. Roy. Soc. London Ser. B*, 352 (1997), pp. 21–38.
- [42] J. SMOLLER, *Shock Waves and Reaction-Diffusion Equations*, *Grundlehren Math. Wiss.* 258, Springer-Verlag, New York, 1983.
- [43] T. STAIB, *Control of a chemical reactor (reaction-diffusion equation)*, *Z. Angew. Math. Mech.*, 76 (1996), pp. 681–682.
- [44] I. TÁTRAI, K. MÁTYÁS, J. KORPONAI, G. PAULOVITS, P. POMOGYI, AND F. PEKÁR, *Manage-*

- ment of fish communities and its impacts on the lower trophic levels in shallow ecosystems in Hungary*, *Hydrobiologia*, 506–509 (2003), pp. 489–496.
- [45] I. TIKHONOVA, B. LI, H. MALCHOW, AND A. MEDVINSKY, *The impact of the phytoplankton growth rate on spatial and temporal dynamics of plankton communities in a heterogeneous environment*, *Biofizika*, 48 (2003), pp. 891–899.
- [46] P. TU, *Dynamical Systems: An Introduction with Applications in Economics and Biology*, Springer-Verlag, Berlin, 1994.
- [47] S. WIGGINS, *Introduction to Applied Nonlinear Dynamical Systems and Chaos*, Texts Appl. Math. 2, Springer-Verlag, New York, 1990.
- [48] A. WISSEL AND J. BENNDORF, *Contrasting effects of the invertebrate predator Chaoborus obscuripes and planktivorous fish on plankton communities of a long term biomanipulation experiment*, *Arch. Hydrobiol.*, 143 (1998), pp. 129–146.
- [49] B. WISSEL, K. FREIER, B. MÜLLER, J. KOOP, AND J. BENNDORF, *Moderate planktivorous fish biomass stabilizes biomanipulation by suppressing large invertebrate predators of Daphnia*, *Arch. Hydrobiol.*, 149 (2000), pp. 177–192.
- [50] K. WYSUJACK AND T. MEHNER, *Comparison of losses of planktivorous fish by predation and seine-fishing in a lake undergoing long-term biomanipulation*, *Freshwater Biol.*, 47 (2002), pp. 2425–2434.